

# Renormalization group and elliptic homogenization in high contrast

Scott Armstrong <sup>\*</sup>

Tuomo Kuusi <sup>†</sup>

May 17, 2024

## Abstract

We prove a quantitative estimate on the homogenization length scale in terms of the ellipticity ratio  $\Lambda/\lambda$  of the coefficient field. This upper bound is applicable to high contrast elliptic equations demonstrating near-critical behavior. Specifically, we show that, given a suitable decay of correlation, the length scale at which homogenization is observed is at most  $\exp(C \log^3(1 + \Lambda/\lambda))$ . The proof introduces the new concept of coarse-grained ellipticity, which measures the effective ellipticity ratio of the equation—and thus the strength of the disorder—after integrating out smaller scales. By a direct analytic argument, we obtain an approximate differential inequality for this coarse-grained ellipticity as a function of the length scale. This approach can be interpreted as a rigorous renormalization group argument, and provides a quantitative framework for homogenization that can be iteratively applied across an arbitrary number of length scales.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>The coarse-grained diffusion matrices</b>	<b>17</b>
<b>3</b>	<b>Renormalization in high contrast</b>	<b>47</b>
<b>4</b>	<b>Renormalization in small contrast</b>	<b>70</b>
<b>5</b>	<b>Equations with weaker mixing assumptions</b>	<b>83</b>
<b>6</b>	<b>Quantitative homogenization</b>	<b>90</b>
<b>A</b>	<b>Besov spaces and functional inequalities</b>	<b>123</b>
<b>B</b>	<b>Geometric means for positive matrices</b>	<b>126</b>
<b>C</b>	<b>Orlicz quasi-norms and concentration inequalities</b>	<b>127</b>
<b>D</b>	<b>Examples of random fields satisfying the assumptions</b>	<b>139</b>
	<b>References</b>	<b>154</b>

---

<sup>\*</sup>Courant Institute of Mathematical Sciences, New York University. [scotta@cims.nyu.edu](mailto:scotta@cims.nyu.edu).

<sup>†</sup>Department of Mathematics and Statistics, University of Helsinki. [tuomo.kuusi@helsinki.fi](mailto:tuomo.kuusi@helsinki.fi).

# 1. Introduction

**1.1. Homogenization in high contrast.** We consider the elliptic equation

$$-\nabla \cdot \mathbf{a}(x) \nabla u = 0 \quad \text{in } U \subseteq \mathbb{R}^d, \quad (1.1)$$

in which  $\mathbf{a}(x)$  is a  $\mathbb{Z}^d$ -stationary random field in dimension  $d \geq 2$ , valued in the set  $\mathbb{R}^{d \times d}$  of  $d \times d$  matrices with real entries. Under appropriate ellipticity and decorrelation assumptions on the coefficient field  $\mathbf{a}(x)$ , the equation (1.1) homogenizes on large scales. This means essentially that its solutions will be close to those of the equation  $-\nabla \cdot \bar{\mathbf{a}} \nabla u_{\text{hom}} = 0$ , for a constant and deterministic matrix  $\bar{\mathbf{a}}$  which depends in a quite complicated way on the law of  $\mathbf{a}(x)$  and describes the macroscopic behavior of the system. The phenomenon of elliptic homogenization is observed across a wide spectrum of statistical physics problems, in which diffusive limits on macroscopic scales are expected to emerge from microscopic disorder.

Given an error tolerance  $\delta > 0$ , the *homogenization length scale*  $\mathcal{X}$  is a random variable which is defined in rough terms as the minimal scale above which the relative homogenization error (the ratio of the size of the error  $u - u_{\text{hom}}$  to the size of  $u_{\text{hom}}$ ) is smaller than  $\delta$ . This length scale characterizes the transition beyond which the local fluctuations in the solutions have averaged out and the macroscopic behavior is dominant. As such, the stochastic moments of  $\mathcal{X}$  measure the extent to which local randomness in the field  $\mathbf{a}(x)$  affects the large-scale properties of the solutions.

In this paper, we establish quantitative upper bounds on the homogenization length scale for very general coefficient fields. These estimates depend explicitly on the ellipticity of the field  $\mathbf{a}(x)$ , as well as its rate of decorrelation, but otherwise require no structural assumption on the law of  $\mathbf{a}(x)$ . Of particular interest is the dependence of  $\mathcal{X}$  on the ellipticity of the coefficient field. We obtain estimates which are completely new even in the special case that the field  $\mathbf{a}(x)$  satisfies (almost surely) a uniform ellipticity condition of the form

$$\exists \lambda, \Lambda > 0, \quad \lambda \leq \Lambda, \quad \lambda |e|^2 \leq e \cdot \mathbf{a}(x) e \quad \text{and} \quad \Lambda^{-1} |e|^2 \leq e \cdot \mathbf{a}^{-1}(x) e, \quad \forall x, e \in \mathbb{R}^d. \quad (1.2)$$

Any upper bound on  $\mathcal{X}$  for general fields satisfying (1.2) must diverge as the *ellipticity ratio*  $\Lambda/\lambda$  tends to infinity—even under the strongest possible mixing assumptions, such as a finite range of dependence. Indeed, as  $\Lambda/\lambda$  increases, the dependence of the solutions on the coefficient field becomes more singular, necessitating, in general, a larger scale separation for the macroscopic behavior to manifest. This divergence of the homogenization length scale mirrors critical phenomena widely observed in statistical physics, in which systems nearing critical points exhibit diverging correlation lengths and increased sensitivity to external parameters.

An example which exhibits a diverging homogenization length scale, and to which our results apply, is a continuum version of the conductance model at criticality. Consider a Poisson point process on  $\mathbb{R}^d$  with intensity  $\gamma > 0$  and let  $A \subseteq \mathbb{R}^d$  be the union of all balls of radius one centered at a point in the point cloud. It is well-known that there is a critical value  $\gamma_c \in (0, \infty)$  such that  $A$  has an infinite connected component (which is necessarily unique) if  $\gamma > \gamma_c$ , and no infinite component if  $\gamma < \gamma_c$ . This is often referred to as the *continuum percolation model*, and we associate an elliptic coefficient field to it by setting  $\mathbf{a} := \mathbf{I}_d \mathbf{1}_A + \lambda \mathbf{1}_{\mathbb{R}^d \setminus A}$ , where  $0 < \lambda \ll 1$  is a small parameter. The scalar field  $\mathbf{a}(x)$  has the physical interpretation of the conductivity of a random material at the point  $x$ . It satisfies (1.2) with  $(\Lambda, \lambda) = (1, \lambda)$  and so its ellipticity ratio is  $\lambda^{-1}$ . Smaller values of  $\lambda$  mean the resistance of the vacant set  $\mathbb{R}^d \setminus A$  is larger, and the flux of the solutions of (1.1) will therefore be more concentrated on the set  $A$ . The connectedness (or lack thereof) of the set  $A$  becomes the main driver of large-scale behavior of solutions, and in the limit  $\lambda \rightarrow 0$  we should

expect the homogenization length scale  $\mathcal{X}$  to be roughly of the same order as the correlation length scale of the underlying percolation problem. However, since  $\gamma = \gamma_c$ , this is infinite. It is therefore natural to wonder how large we should expect  $\mathcal{X}$  to be as a function of  $\lambda^{-1}$ .

The following theorem is the first general quantitative estimate in *high contrast homogenization*. Assuming only the uniform ellipticity condition (1.2) and a unit range of dependence, it provides an upper bound estimate on the homogenization length scale which states roughly that

$$\mathcal{X} \lesssim \exp(C \log^3(1 + \Lambda/\lambda)) \simeq \left(\frac{\Lambda}{\lambda}\right)^{C \log^2(1 + \Lambda/\lambda)} \quad (1.3)$$

for a prefactor constant  $C(\delta, d) < \infty$  which depends only on the dimension  $d$  and tolerance  $\delta$ . This estimate is a special case of the main results in the paper, presented below in Section 1.2, which apply to more general coefficient fields (very singular and/or degenerate fields are allowed, as are those with much weaker decay of correlations) and provide stronger, more extensive quantitative estimates in their conclusions.

**Theorem A** (Quantitative homogenization in high contrast). *Let  $\mathbb{P}$  be the law of a  $\mathbb{Z}^d$ -stationary random field  $\mathbf{a}(\cdot)$ , valued in  $\mathbb{R}^{d \times d}$ , such that:*

- $\mathbf{a}(\cdot)$  satisfies the uniform ellipticity condition (1.2) with constants  $0 < \lambda \leq 1 \leq \Lambda < \infty$ ,  $\mathbb{P}$ -a.s.
- $\mathbf{a}(\cdot)$  has a unit range of dependence.

Let  $\bar{\mathbf{a}}$  denote the corresponding homogenized matrix,  $\bar{\mathbf{s}} := \frac{1}{2}(\bar{\mathbf{a}} + \bar{\mathbf{a}}^t)$  be the symmetric part of  $\bar{\mathbf{a}}$  and  $\bar{\lambda}$  be the smallest eigenvalue of  $\bar{\mathbf{s}}$ . Denote the family  $\{E_r\}_{r \geq 0}$  of ellipsoids adapted to  $\bar{\mathbf{s}}$  by

$$E_r := \{x \in \mathbb{R}^d : x \cdot \bar{\mathbf{s}}^{-1} x \leq \bar{\lambda}^{-1} r^2\}. \quad (1.4)$$

Then, for every  $\delta > 0$ , there exists  $C(\delta, d) < \infty$  and a nonnegative random variable  $\mathcal{X}$  satisfying

$$\mathbb{P}[\mathcal{X} \geq t \exp(C \log^3(1 + \Lambda/\lambda))] \leq \exp(-t^d), \quad \forall t \in [1, \infty), \quad (1.5)$$

such that the following statements are valid:

- Homogenization of the Dirichlet problem. For every  $r \geq \mathcal{X}$ ,  $f \in L^2(E_r)$  and  $g \in H^1(E_r)$ , if we let  $u, u_{\text{hom}} \in H^1(E_r)$  be the solutions of the Dirichlet problems

$$\begin{cases} -\nabla \cdot \mathbf{a} \nabla u = 0 & \text{in } E_r, \\ u = g & \text{on } \partial E_r, \end{cases} \quad \text{and} \quad \begin{cases} -\nabla \cdot \mathbf{a} \nabla u_{\text{hom}} = 0 & \text{in } E_r, \\ u_{\text{hom}} = g & \text{on } \partial E_r, \end{cases} \quad (1.6)$$

then we have the estimate

$$\|u - u_{\text{hom}}\|_{L^2(E_r)} \leq \delta(r \|\nabla g\|_{L^2(E_r)} + r^2 \|f\|_{L^2(E_r)}). \quad (1.7)$$

- Large-scale  $C^{0,1}$  regularity. For every  $R \geq \mathcal{X}$  and solution  $u \in H^1(E_R)$  of the equation

$$-\nabla \cdot \mathbf{a} \nabla u = 0 \quad \text{in } E_R,$$

we have the estimate

$$\sup_{r \in [\mathcal{X}, R]} \int_{E_r} \nabla u \cdot \mathbf{a} \nabla u \leq C \int_{E_R} \nabla u \cdot \mathbf{a} \nabla u. \quad (1.8)$$

It is widely believed that the divergences of correlation lengths near critical points, as well as that of other physical quantities, are described by power laws. That is, one expects a correlation length to be a power of the underlying macroscopic control parameters driving the phase transition. Typical examples include percolation and Ising models, in which one expects the correlation length  $\xi$  to be of order  $|p - p_c|^{-\nu}$  and  $|T/T_c - 1|^{-\nu}$ , respectively, where  $p_c$  is the critical percolation probability,  $T_c$  is the critical temperature for the Ising model, and the occupation probability  $p \in (0, 1)$  and temperature  $T > 0$  are the control parameters. In the continuum percolation model described above, we would expect the correlation length  $\xi$  to be of order  $|\gamma - \gamma_c|^{-\nu}$ . The value of  $\nu$ , as well as that of other critical exponents, is expected to be universal in the sense that it should depend only on the dimension  $d$  and the type of model, but not on the particular microscopic details of the model. For instance, the value of  $\nu$  for a bond percolation model on the lattice  $\mathbb{Z}^d$  is expected to be the same as the exponent for continuum percolation.

The existence of critical exponents is of great physical interest and there is a large literature devoted to estimating and computing them, with predictions of their exact values available for certain models. This is typically accomplished by heuristic renormalization group arguments, with rigorous results being comparatively rare. Some models for which critical exponents are known rigorously include certain two-dimension models in which conformal invariance can be exploited (such as site percolation on the  $2d$  triangular lattice [SW01]), some exactly integrable models, and in sufficiently large dimensions where mean field methods are applicable (see [Har90] in the case of bond percolation). For most models, polynomial upper bounds—much less the existence of critical exponents—have not been rigorously demonstrated. For instance, to our knowledge, the best upper bound for the correlation length for Bernoulli bond percolation on the lattice  $\mathbb{Z}^2$  is

$$\xi \leq \exp(C|p - p_c|^{-2}). \quad (1.9)$$

This result was proved in [DCKT21] for  $\mathbb{Z}^d$  in general dimension  $d \geq 2$ , and is obviously far from the expected polynomial-type dependence.

In the context of the elliptic equation (1.1), the ellipticity ratio  $\Lambda/\lambda$  plays the role of a control parameter, and so the natural conjecture is that the homogenization length scale should satisfy an upper bound of the form  $\mathcal{X} \lesssim (\Lambda/\lambda)^\nu$  for some finite exponent  $\nu(d) < \infty$ . Proving such an upper bound estimate is perhaps the most important open problem in quantitative homogenization. Apart from its intrinsic interest, such an estimate would have immediate and important consequences in mathematical physics and probability. Theorem A does not provide such an estimate. However, the upper bound in (1.3) is close to a polynomial-type bound in the sense that the desired fixed exponent  $\nu = C$  replaced by one that is only logarithmically diverging,  $\nu = C \log^2(1 + \Lambda/\lambda)$ , which is evidently significantly better than an exponential upper bound like the one in (1.9). In fact, it is to our knowledge the best rigorous upper bound which has been obtained for a general class of models in low dimensions.

To prove Theorem A, we study certain coarse-grained diffusion matrices which are defined at a given scale and in a particular region of space. Based on these objects, we introduce the new concept of *coarse-grained ellipticity*, which is a relaxation of the usual uniform ellipticity ratio. This quantity is a softer and more flexible notion of ellipticity compared to uniform ellipticity and, in particular, permits certain degenerate and unbounded coefficient fields. We view the process of homogenization as a *flow* of the coarse-grained ellipticity, from a possibly very large number at small length scales to unity in the large-scale limit. Indeed, as we show in the paper, the homogenization error can be controlled, in a deterministic way, by the coarse-grained ellipticity. The homogenization length scale  $\mathcal{X}$  is then, roughly, the scale at which the coarse-grained ellipticity is smaller than  $1 + \delta$ . At

the heart of this paper is an analytic argument which obtains a differential inequality for coarse-grained ellipticity as a function of (the logarithm of) the length scale, which then implies that the desired bound on the homogenization length scale. This argument is notable for being entirely *renormalizable* in the sense that its outputs (bounds on the coarse-grained ellipticity) are of the same form as its inputs. It is therefore possible to iterate it, and indeed the proof Theorem A relies on such an iteration.

In a concurrent joint work with Bou-Rabee [ABRK23], we prove a superdiffusive central limit theorem for a Brownian particle in a critically correlated, divergence-free drift. The high contrast homogenization estimates proved in this present paper played an important role in the arguments in [ABRK23]. In particular, the fact that the exponent in our estimate for the homogenization length scale in (1.3) grows logarithmically, rather than like a power of  $\Lambda/\lambda$ , is crucial. In the context of that paper, homogenization estimates must iterated an infinite number of times as a way of formalizing a renormalization group argument. One difficulty encountered there is that the ellipticity ratio is also growing as a function of the scale, and there is an apparent “race” between homogenization and an accumulation of disorder. A quantitative estimate like (1.3) is needed to ensure that the randomness at each scale can be integrated out before interacting in an unexpected way with the other, larger scales. Of course, this kind of phenomenon is not unique to this particular problem, and we expect that the methods developed here will find similar applications to other critical models in mathematical physics.

Quantitative estimates for elliptic homogenization problems have been extensively studied in recent years in the regime of *moderate* ellipticity contrast. By this we mean that the ellipticity ratio  $\Lambda/\lambda$  is held fixed and the goal, broadly speaking, is to obtain estimates for the homogenization error as a function of the scale separation ratio as it asymptotically approaches infinity. Originating in the pioneering works [GO11, GO12], this theory has recently reached maturity and there are now very detailed and precise quantitative estimates available (an overview and further references can be found in our monograph [AK24]). Each of the several independent approaches to this theory uses arguments which are constructive and produces constants which are explicitly computable. While the dependence on the ellipticity ratio has been kept implicit in this literature, it is possible to extract an estimate for the homogenization length scale  $\mathcal{X}$  by tracking the dependence of  $\Lambda/\lambda$  through the whole theory. Prior to this work, such a bookkeeping exercise would reveal, in all cases, an exponential upper bound, comparable to (1.9), of the form  $\mathcal{X} \lesssim \exp(C(\Lambda/\lambda)^p)$  for an exponent  $p$  which is at least  $1/2$  and, we expect, typically much larger (see the discussion below (1.45) for more).

This paper is also the first to develop a systematic theory of quantitative homogenization for a wide range of degenerate equations. Previous works on quantitative homogenization in non-uniformly elliptic settings have addressed finite difference equations on supercritical bond percolation clusters [AD18, Dar21, DG21], domains with random inclusions [DG22], and, more recently, log-normal coefficients with an integrable covariance function [CGQ24] (cf. Proposition 1.3, below). These works extend techniques from the moderate contrast, uniformly elliptic theory while managing specific degeneracies in an ad hoc manner. Each considers only “far from critical” cases.<sup>1</sup> In contrast, our introduction of the concept of *coarse-grained ellipticity* leads to a quantitative theory covering a broad class of degenerate equations. Note that, while the estimates stated in this

---

<sup>1</sup>For example, if the parameter  $\gamma$  in the conductance model mentioned above is either very large or very small, then the system does not display critical behavior and can be analyzed quantitatively for all values of  $\lambda$  by arguments which are comparatively much simpler than those deployed here. Similar comments apply to the papers [AD18, Dar21, DG21], which analyze harmonic functions on supercritical bond percolation clusters on  $\mathbb{Z}^d$ , for any  $p > p_c$ , and prove sharp quantitative homogenization estimates. The constants in these estimates however depends on  $p - p_c$  like in (1.9), as the correlation length scale for supercritical percolation is used an input to the homogenization argument. As such, the results in these papers should not be seen as analyzing near-critical phenomena.

paper do not give sharp scaling exponents for the homogenization error in the regime of large scale separation, such estimates can be straightforwardly obtained by combining known techniques from the moderate contrast theory with Theorem B below.

In the next subsection we state our main results, which are a great deal more general than Theorem A. Since our methods are based on renormalizing the coarse-grained ellipticity, the uniform ellipticity condition can be replaced by the assumption that the coarse-grained ellipticity ratio is bounded on sufficiently large scales. This allows us to consider very general coefficient fields which may be very degenerate and/or singular, including some of those mentioned in the previous paragraph and others which have not been previously analyzed.

**1.2. Coarse-grained ellipticity and the statement of the main results.** In this subsection, we state the main result, which is a general quantitative homogenization result for elliptic equations with high contrast coefficients.

We begin by introducing some notation. The set of  $m$ -by- $n$  matrices with real entries is denoted by  $\mathbb{R}^{m \times n}$ . The transpose of a matrix  $A$  is denoted by  $A^t$ . The sets of  $m$ -by- $m$  symmetric and anti-symmetric matrices are denoted, respectively, by

$$\mathbb{R}_{\text{sym}}^{d \times d} := \{A \in \mathbb{R}^{d \times d} : A = A^t\} \quad \text{and} \quad \mathbb{R}_{\text{skew}}^{d \times d} := \{A \in \mathbb{R}^{d \times d} : A = -A^t\}.$$

We also define the cone of matrices with positive symmetric part by

$$\mathbb{R}_+^{d \times d} := \{A \in \mathbb{R}^{d \times d} : e \cdot Ae > 0, \forall e \in \mathbb{R}^d\}.$$

Let  $\Omega$  be the collection of all Lebesgue measurable maps  $\mathbf{a} : \mathbb{R}^d \rightarrow \mathbb{R}_+^{d \times d}$  such that, if we split  $\mathbf{a}(\cdot)$  into its symmetric and anti-symmetric parts by writing  $\mathbf{a}(x) = \mathbf{s}(x) + \mathbf{k}(x)$ , where we define

$$\mathbf{s}(x) := \frac{1}{2}(\mathbf{a}(x) + \mathbf{a}^t(x)) \in \mathbb{R}_{\text{sym}}^{d \times d} \quad \text{and} \quad \mathbf{k}(x) := \frac{1}{2}(\mathbf{a}(x) - \mathbf{a}^t(x)) \in \mathbb{R}_{\text{skew}}^{d \times d}, \quad x \in \mathbb{R}^d, \quad (1.10)$$

then we have that

$$\mathbf{s}, \mathbf{s}^{-1}, \mathbf{k}^t \mathbf{s}^{-1} \mathbf{k} \in L_{\text{loc}}^1(\mathbb{R}^d; \mathbb{R}^{d \times d}). \quad (1.11)$$

The condition (1.11) represents the minimal qualitative ellipticity we require of our coefficient fields.

It is convenient to arrange the symmetric and anti-symmetric parts of the field  $\mathbf{a}(\cdot)$  as the block entries of an  $\mathbb{R}_{\text{sym}}^{2d \times 2d}$ -valued random field  $\mathbf{A}$ , which is defined by

$$\mathbf{A}(x) := \begin{pmatrix} (\mathbf{s} + \mathbf{k}^t \mathbf{s}^{-1} \mathbf{k})(x) & -(\mathbf{k}^t \mathbf{s}^{-1})(x) \\ -(\mathbf{s}^{-1} \mathbf{k})(x) & \mathbf{s}^{-1}(x) \end{pmatrix}. \quad (1.12)$$

The field  $\mathbf{A}(x)$  defined in (1.12) arises naturally in the variational interpretation of (1.1) for coefficient fields  $\mathbf{a}(x)$  which may not be symmetric, and it consequently plays an essential role in coarse-graining. Notice that the assumption (1.11) is just the requirement that  $\mathbf{A}$  belong to  $L_{\text{loc}}^1(\mathbb{R}^d; \mathbb{R}_{\text{sym}}^{2d \times 2d})$ . We may equivalently regard the set  $\Omega$  as being the collection of fields  $\mathbf{A}(\cdot)$  having the form of (1.12), with  $\mathbf{s} \in \mathbb{R}_{\text{sym}}^{d \times d}$  and  $\mathbf{k} \in \mathbb{R}_{\text{skew}}^{d \times d}$ , and with entries belonging to  $L_{\text{loc}}^1(\mathbb{R}^d)$ . By abuse of notation, we will sometimes consider either  $\mathbf{a}(\cdot)$  or  $\mathbf{A}(\cdot)$  as the canonical element of  $\Omega$ , whichever happens to be more convenient. Throughout, the random fields  $\mathbf{s}(\cdot)$  and  $\mathbf{k}(\cdot)$  always refer to those defined in (1.10).

We define a  $\sigma$ -field  $\mathcal{F}(U)$ , for each Borel subset  $U \subseteq \mathbb{R}^d$ , as the one generated by the family random variables of the form

$$\mathbf{a} \mapsto \int_{\mathbb{R}^d} e' \cdot \mathbf{a}(x) e \varphi(x) dx \quad e, e' \in \mathbb{R}^d, \varphi \in C_c^\infty(U).$$



We also denote  $\mathcal{F} := \mathcal{F}(\mathbb{R}^d)$ . We let  $\{T_y : y \in \mathbb{R}^d\}$  denote the group of  $\mathbb{R}^d$  translations acting on  $\Omega$ . That is,  $T_y : \Omega \rightarrow \Omega$  is given by  $T_y \mathbf{a} = \mathbf{a}(\cdot + y)$ . We extend this group action to  $\mathcal{F}$  by defining  $T_y F := \{T_y \mathbf{a} : \mathbf{a} \in F\}$  for  $F \in \mathcal{F}$ .

We consider a probability measure  $\mathbb{P}$  on  $(\Omega, \mathcal{F})$  satisfying the three assumptions (P1), (P2) and (P3) stated below. The first is that  $\mathbb{P}$  is statistically homogeneous.

(P1) *Stationarity with respect to  $\mathbb{Z}^d$ -translations:*

$$\mathbb{P} \circ T_z = \mathbb{P}, \quad \forall z \in \mathbb{Z}^d. \quad (1.13)$$

We turn next to the ellipticity assumption. Conceptually, this assumption requires that the field behaves elliptically only in a suitable coarse-grained sense, a much less rigid condition than uniform ellipticity. We formulate this assumption in terms of the *coarse-grained matrices* which are objects that play a central role in this paper. They are denoted, for each bounded Lipschitz domain  $U \subseteq \mathbb{R}^d$ , by  $\mathbf{A}(U)$ . These are a random elements of  $\mathbb{R}_{\text{sym}}^{2d \times 2d}$  which depend only on the restriction  $\mathbf{a}|_U$  of the field  $\mathbf{a}(\cdot)$  to  $U$  and are to be understood as a coarse-graining of the field  $\mathbf{A}(x)$  in (1.12) with respect to  $U$ . They can be represented in block matrix form as

$$\mathbf{A}(U) := \begin{pmatrix} (\mathbf{s} + \mathbf{k}^t \mathbf{s}_*^{-1} \mathbf{k})(U) & -(\mathbf{k}^t \mathbf{s}_*^{-1})(U) \\ -(\mathbf{s}_*^{-1} \mathbf{k})(U) & \mathbf{s}_*^{-1}(U) \end{pmatrix}, \quad (1.14)$$

As the notation suggests, we think of the matrices  $\mathbf{s}(U)$  and  $\mathbf{s}_*(U)$  as a coarse-graining of the symmetric part  $\mathbf{s}(\cdot)$  of the field  $\mathbf{a}(\cdot)$ , and  $\mathbf{k}(U)$  as a coarse-graining of the anti-symmetric part. If the field  $\mathbf{a}(x)$  is symmetric, then  $\mathbf{k}(U)$  vanishes and the expression in (2.1) simplifies into a block diagonal form, and  $\mathbf{s}(U)$  and  $\mathbf{s}_*(U)$  are equivalent to the “dual” pair of subadditive quantities introduced in [AS16], given by the energy of Dirichlet and Neumann problems in  $U$  with affine boundary data on  $\partial U$ . The coarse-grained matrices were generalized to the case of nonsymmetric coefficient fields in [AM16], and the matrix  $\mathbf{A}(U)$  we work with in this paper is equivalent to the quantities analyzed in that paper and later in [AKM16, AKM17, AKM19] (see [AK24] for the latest exposition). The matrices  $\mathbf{s}(U)$ ,  $\mathbf{s}_*(U)$  and  $\mathbf{k}(U)$  do not have a simple interpretation in terms of Dirichlet and Neumann problems, in general. We postpone their definitions to Section 2.2.

The ellipticity assumption roughly says that the coarse-grained matrices, on scales larger than a sufficiently large (random) scale, are bounded by a deterministic constant;<sup>2</sup> and moreover, on such scales, the coarse-grained matrices for smaller subcubes are also upper-bounded by a power  $\gamma < 1$  of the scale separation ratio. Throughout the paper, we denote triadic cubes by

$$\square_m := \left( -\frac{1}{2}3^m, \frac{1}{2}3^m \right)^d.$$

(P2) *Ellipticity above a minimal scale.* There exist a matrix  $\mathbf{E}_0 \in \mathbb{R}_{\text{sym}}^{2d \times 2d}$ , an exponent  $\gamma \in [0, 1)$ , an increasing function  $\Psi_{\mathcal{S}} : \mathbb{R}_+ \rightarrow [1, \infty)$ , a constant  $K_{\Psi_{\mathcal{S}}} \in (1, \infty)$  satisfying the growth condition

$$t\Psi_{\mathcal{S}}(t) \leq \Psi_{\mathcal{S}}(K_{\Psi_{\mathcal{S}}}t), \quad \forall t \in [1, \infty), \quad (1.15)$$

and a nonnegative random variable  $\mathcal{S}$  which satisfies the bound

$$\mathbb{P}[\mathcal{S} > t] \leq \frac{1}{\Psi_{\mathcal{S}}(t)}, \quad \forall t \in (0, \infty), \quad (1.16)$$

---

<sup>2</sup>An upper bound for  $\mathbf{A}(x)$  in (1.12) is equivalent to an upper bound for  $\mathbf{a}(x)$  and a lower bound for its symmetric part  $\mathbf{s}(x)$ . Likewise, an upper bound for  $\mathbf{A}(U)$  is analogous to a double-sided ellipticity bound in a coarse-grained sense.

such that, for every  $m \in \mathbb{Z}$ ,

$$3^m \geq \mathcal{S} \implies \mathbf{A}(z + \square_n) \leq 3^{\gamma(m-n)} \mathbf{E}_0, \quad \forall n \in \mathbb{Z} \cap (-\infty, m], \quad z \in 3^n \mathbb{Z}^d \cap \square_m. \quad (1.17)$$

We will have more to say on the motivation behind (P2) in Section 1.3 below, as well as some examples of coefficient fields satisfying it. For the moment, we mention that the above assumption with  $\mathcal{S} = 0$  and  $\gamma = 0$  is equivalent to the classical uniform ellipticity condition. This is the assumption that (1.2) is satisfied with full  $\mathbb{P}$ -probability. To see this, observe that the condition (1.2) is equivalent to

$$\mathbf{s}^{-1}(x) \leq \lambda^{-1} \mathbf{I}_d \quad \text{and} \quad (\mathbf{s} + \mathbf{k} \mathbf{s}^{-1} \mathbf{k}^t)(x) \leq \Lambda \mathbf{I}_d, \quad \forall x \in \mathbb{R}^d. \quad (1.18)$$

This in turn implies that the matrix  $\mathbf{A}(x)$  in (1.12) satisfies

$$\mathbf{A}(x) = \begin{pmatrix} (\mathbf{s} + \mathbf{k}^t \mathbf{s}^{-1} \mathbf{k})(x) & -(\mathbf{k}^t \mathbf{s}^{-1})(x) \\ -(\mathbf{s}^{-1} \mathbf{k})(x) & \mathbf{s}^{-1}(x) \end{pmatrix} \leq \begin{pmatrix} 2\Lambda \mathbf{I}_d & 0 \\ 0 & 2\lambda^{-1} \mathbf{I}_d \end{pmatrix}.$$

It turns out that the coarse-grained matrix  $\mathbf{A}(U)$  in any bounded domain  $U$  is bounded from above by the mean in  $U$  of the matrix  $\mathbf{A}(x)$  defined in (1.12):

$$\mathbf{A}(U) \leq \int_U \mathbf{A}(x) dx \leq 2 \int_U \begin{pmatrix} (\mathbf{s} + \mathbf{k}^t \mathbf{s}^{-1} \mathbf{k})(x) & 0 \\ 0 & \mathbf{s}^{-1}(x) \end{pmatrix} dx. \quad (1.19)$$

It therefore follows that the classical uniform ellipticity assumption (1.2) implies the assumption (P2) with  $\gamma = 0$ ,  $\mathcal{S} = 0$ , and

$$\mathbf{E}_0 = \begin{pmatrix} 2\Lambda \mathbf{I}_d & 0 \\ 0 & 2\lambda^{-1} \mathbf{I}_d \end{pmatrix}. \quad (1.20)$$

Conversely, in view of the equivalence of (1.2) and (1.18), the assumption (P2) with  $\gamma = 0$  and  $\mathcal{S} = 0$  implies (by the Lebesgue differentiation theorem) the uniform ellipticity condition (1.2) where  $\Lambda$  is the largest eigenvalue of the upper  $d$ -by- $d$  block of  $\mathbf{E}_0$  and  $\lambda^{-1}$  is the largest eigenvalue of the lower  $d$ -by- $d$  block of  $\mathbf{E}_0$ . Therefore, up to changing the ellipticity constants by at most a factor of two, the assumption (P2), with  $\mathbf{E}_0$  as in the previous display, is equivalent to the uniform ellipticity condition (1.2).

The purpose of the exponent  $\gamma$  and minimal scale  $\mathcal{S}$  is precisely to allow for the non-uniformity of the ellipticity condition. The random scale  $\mathcal{S}$  allows us to consider fields which are “elliptic” only on sufficiently large scales, with the function  $\Psi_{\mathcal{S}}$  quantifying the distribution of the random variable  $\mathcal{S}$ . When computing the “ellipticity” of a given scale, the exponent  $\gamma$  allows us to be more forgiving of bad behavior on small scales by discounting these scales by a power of the scale separation. This flexibility will turn out to be very useful, even in the analysis of the uniformly elliptic case. This is because this ellipticity assumption (P2) is *renormalizable*: as we will show, the pushforward of  $\mathbb{P}$  of the coefficients under a dilation map  $\mathbf{a} \mapsto \mathbf{a}(R \cdot)$ , with  $R > 1$ , will also satisfy the same ellipticity condition—but with a possible improvement of the ellipticity constants (in the sense that the matrix  $\mathbf{E}_0$  may be smaller). This is the way we renormalize the equation.

Associated to the matrix  $\mathbf{E}_0$  in assumption (P2) are some important objects and parameters. We start from the block form

$$\mathbf{E}_0 = \begin{pmatrix} \mathbf{E}_{11} & \mathbf{E}_{12} \\ \mathbf{E}_{21} & \mathbf{E}_{22} \end{pmatrix}, \quad (1.21)$$



where  $\mathbf{E}_{ij} \in \mathbb{R}^{d \times d}$  for  $i, j \in \{1, 2\}$ , and we define

$$\begin{cases} \mathbf{s}_{*,0} := \mathbf{E}_{22}^{-1}, \\ \mathbf{k}_0 := -\mathbf{E}_{22}^{-1} \mathbf{E}_{21}, \\ \mathbf{s}_0 := \mathbf{E}_{11} - \mathbf{E}_{12} \mathbf{E}_{22}^{-1} \mathbf{E}_{21}, \\ \mathbf{b}_0 := \mathbf{E}_{11}. \end{cases} \quad (1.22)$$

In other words, we have given names to the block entries of  $\mathbf{E}_0$  so that

$$\mathbf{E}_0 = \begin{pmatrix} \mathbf{s}_0 + \mathbf{k}_0^t \mathbf{s}_{*,0}^{-1} \mathbf{k}_0 & -\mathbf{k}_0^t \mathbf{s}_{*,0}^{-1} \\ -\mathbf{s}_{*,0}^{-1} \mathbf{k}_0 & \mathbf{s}_{*,0}^{-1} \end{pmatrix} = \begin{pmatrix} \mathbf{b}_0 & -\mathbf{k}_0^t \mathbf{s}_{*,0}^{-1} \\ -\mathbf{s}_{*,0}^{-1} \mathbf{k}_0 & \mathbf{s}_{*,0}^{-1} \end{pmatrix}. \quad (1.23)$$

It follows that

$$\mathbf{E}_0 \leq 2 \begin{pmatrix} \mathbf{s}_0 + \mathbf{k}_0^t \mathbf{s}_{*,0}^{-1} \mathbf{k}_0 & 0 \\ 0 & \mathbf{s}_{*,0}^{-1} \end{pmatrix}. \quad (1.24)$$

We next define the *ellipticity ratio*  $\Theta$  by<sup>3</sup>

$$\Theta := \min_{\mathbf{h} \in \mathbb{R}_{\text{skew}}^{d \times d}} |\mathbf{s}_{*,0}^{-1/2} (\mathbf{s}_0 + (\mathbf{k}_0 - \mathbf{h})^t \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{h})) \mathbf{s}_{*,0}^{-1/2}|. \quad (1.25)$$

We let  $\mathbf{h}_0 \in \mathbb{R}_{\text{skew}}^{d \times d}$  to be a minimizer of the above quantity, so that

$$\Theta = |\mathbf{s}_{*,0}^{-1/2} (\mathbf{s}_0 + (\mathbf{k}_0 - \mathbf{h}_0)^t \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{h}_0)) \mathbf{s}_{*,0}^{-1/2}|.$$

We also define the ellipticity constants  $0 < \lambda \leq \Lambda < \infty$  by

$$\lambda := |\mathbf{s}_{*,0}^{-1}|^{-1} \quad \text{and} \quad \Lambda := \min_{\mathbf{h} \in \mathbb{R}_{\text{skew}}^{d \times d}} |(\mathbf{s}_0 + (\mathbf{k}_0 - \mathbf{h})^t \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{h}))| \quad (1.26)$$

and the *aspect ratio*  $\Pi$  by

$$\Pi := \frac{\Lambda}{\lambda}. \quad (1.27)$$

Here, and in the rest of the paper,  $|\mathbf{B}|$  denotes the spectral norm of a square matrix  $\mathbf{B}$ , that is, the square root of the largest eigenvalue of  $\mathbf{B}^t \mathbf{B}$ . We will discover that, since  $\mathbf{E}_0$  dominates the coarse-grained matrices by (P2), we must have the ordering

$$\mathbf{s}_0 \geq \mathbf{s}_{*,0}.$$

It follows that

$$1 \leq \Theta \leq \Pi = \frac{\Lambda}{\lambda}.$$

Why do we have two competing notions of ellipticity ratio,  $\Theta$  and  $\Pi$ ? The classical ellipticity assumption (1.2) simultaneously controls two different things, which we need to keep separate: (i) the ratio of the size of  $\mathbf{a}(x)$  to its smallest eigenvalue at each point  $x$ ; and (ii) the ratio of matrices  $\mathbf{a}(x)$  and  $\mathbf{a}(y)$  at two different points  $x$  and  $y$ . It is important in our setting to distinguish these two, because obviously homogenization should be concerned with (ii) but not with (i). Here it is the aspect ratio  $\Pi$  which measures (i), while the ellipticity ratio  $\Theta$  measures (ii).

---

<sup>3</sup>The motivation for defining  $\Theta$  modulo the subtraction of a constant anti-symmetric matrix can be found below in Section 2.3.

The third and final assumption we need is a quantitative ergodicity condition. The one we use here is formulated in terms of *concentration for sums*, a general mixing condition we previously introduced in [AK24]. It is a linear concentration inequality which is flexible enough to contain all of the different quantitative ergodic assumptions used in stochastic homogenization literature, but still strong enough to yield optimal quantitative homogenization estimates for the most important examples. In particular, it contains finite range of dependence as well as the assumption of a logarithmic Sobolev inequality as special cases.

In order to state this condition, we require some terminology. We first introduce a measure of the “sensitivity” of a random variable with respect to perturbations of the coefficients in a given subset  $U \subseteq \mathbb{R}^d$ . Given an  $\mathcal{F}$ -measurable random variable  $X$  on  $\Omega$ , we define another random variable  $|D_U X|$  by setting, for each  $\mathbf{A} \in \Omega$ ,

$$\begin{aligned} |D_U X|(\mathbf{A}) \\ := \limsup_{t \rightarrow 0} \frac{1}{2t} \sup \left\{ X(\mathbf{A}_1) - X(\mathbf{A}_2) : \mathbf{A}_1, \mathbf{A}_2 \in \Omega, |\mathbf{A}^{-1/2} \mathbf{A}_i \mathbf{A}^{-1/2} - \mathbf{I}_{2d}| \leq t \mathbf{1}_U, \forall i \in \{1, 2\} \right\}. \end{aligned} \quad (1.28)$$

In contrast to the usual notion of “Malliavin derivative” which measures sensitivity with respect to uniformly elliptic fields, we measure our perturbations to  $\mathbf{a}(\cdot)$  *multiplicatively* rather than *additively*. Of course, in the uniformly elliptic case this is a distinction without a difference, but our choice is the more natural one from the point of view of degenerate equations.

(P3) *Concentrations for sum (CFS)*. There exist  $\beta \in [0, 1)$  and  $\nu \in (\gamma, \frac{d}{2}]$  and an increasing function  $\Psi : \mathbb{R}_+ \rightarrow [1, \infty)$  and a constant  $K_\Psi \in [3, \infty)$  satisfying the growth condition

$$t\Psi(t) \leq \Psi(K_\Psi t), \quad \forall t \in [1, \infty), \quad (1.29)$$

such that, for every  $m, n \in \mathbb{N}$  with  $\beta m < n < m$  and family  $\{X_z : z \in 3^n \mathbb{Z}^d \cap \square_m\}$  of random variables satisfying, for every  $z \in 3^n \mathbb{Z}^d \cap \square_m$ ,

$$\begin{cases} \mathbb{E}[X_z] = 0, \\ |X_z| \leq 1, \\ |D_{z+\square_n} X_z| \leq 1, \\ X_z \text{ is } \mathcal{F}(z + \square_n)\text{-measurable}, \end{cases} \quad (1.30)$$

we have the estimate

$$\mathbb{P} \left[ \left| \sum_{z \in 3^n \mathbb{Z}^d \cap \square_m} X_z \right| \geq t 3^{-\nu(m-n)} \right] \leq \frac{1}{\Psi(t)}, \quad \forall t \in [1, \infty). \quad (1.31)$$

The mixing condition (P3) is discussed further in [AK24, Chapter 3] as well as in Section 1.3, below, where we also give some explicit examples satisfying it. We remark that, in the case of finite range of dependence or LSI, the assumption (P3) is satisfied with  $(\beta, \nu) = (0, \frac{d}{2})$  and  $\Psi(t) = \exp(ct^2)$  for some constant  $c(d) > 0$ : see [AK24, Chapter 3].

The following is the main result of the paper. It gives an explicit estimate for the length scale at which we first see homogenization. Some of the notation appearing in the statement is defined below. For instance, the space  $H_s^1$  is defined below in Section 2.1, see (2.2). The underlines in the norms  $\|\cdot\|_{\underline{L}^2}$  and  $\|\cdot\|_{\underline{H}^{-1}}$  denote volume normalization; see (1.49) and (1.50).

A summary of the main ideas and key steps in the proof of the following theorem appears below in Section 1.4.

**Theorem B** (Homogenization in high contrast). *Assume that  $\mathbb{P}$  is a probability measure on  $(\Omega, \mathcal{F})$  satisfying assumptions (P1), (P2) and (P3) above. Define a parameter  $\alpha := (\min\{\nu, 1\} - \gamma)(1 - \beta)$ . There exists a constant  $C(d) < \infty$ , a nonnegative random variable  $\mathcal{X}$  and a matrix  $\bar{\mathbf{a}} \in \mathbb{R}_+^{d \times d}$  such that, if we let  $\bar{\mathbf{s}}$ ,  $\bar{\lambda}$  and  $\{E_r\}_{r \geq 0}$  be as in Theorem A, let  $\kappa := C^{-1}\alpha$  and define a length scale  $L$  by*

$$L := \exp\left(\frac{C}{\alpha^6} \log(\Pi K_\Psi K_{\Psi_S}) \log^2(1 + \Theta)\right), \quad (1.32)$$

then the following statements are valid:

- Estimate of the homogenization length scale. For every  $t \geq 1$  and for  $\mu := (\nu - \gamma)(1 - \beta)$ ,

$$\mathbb{P}[\mathcal{X} \geq CLt] \leq \frac{1}{\Psi(t^\mu)} + \frac{1}{\Psi_S(CLt)}. \quad (1.33)$$

- Harmonic approximation. For every  $r \geq \mathcal{X}$ , and  $u \in H_s^1(E_{3\sqrt{dr}})$  satisfying  $-\nabla \cdot \mathbf{a} \nabla u = 0$  in  $E_{3\sqrt{dr}}$ , there exists  $u_{\text{hom}} \in H^1(E_r)$  satisfying  $-\nabla \cdot \bar{\mathbf{a}} \nabla u_{\text{hom}} = 0$  in  $E_r$  such that we have the estimate

$$\begin{aligned} \frac{\bar{\lambda}^{1/2}}{r} \|u - u_{\text{hom}}\|_{\underline{L}^2(E_r)} + \|\bar{\mathbf{s}}^{1/2}(\nabla u - \nabla u_{\text{hom}})\|_{\underline{H}^{-1}(E_r)} + \|\bar{\mathbf{s}}^{-1/2}(\mathbf{a} \nabla u - \bar{\mathbf{a}} \nabla u_{\text{hom}})\|_{\underline{H}^{-1}(E_r)} \\ \leq C \left(\frac{r}{\mathcal{X}}\right)^{-\kappa} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(E_{3\sqrt{dr}})}. \end{aligned} \quad (1.34)$$

Conversely, for every  $r \geq \mathcal{X}$  and  $u_{\text{hom}} \in H^1(E_{6\sqrt{dr}})$  satisfying  $-\nabla \cdot \bar{\mathbf{a}} \nabla u_{\text{hom}} = 0$  in  $E_{6\sqrt{dr}}$ , there exists  $u \in H^1(E_r)$  satisfying  $-\nabla \cdot \mathbf{a} \nabla u = 0$  in  $E_r$  such that

$$\text{Left Side of (1.34)} \leq C \left(\frac{r}{\mathcal{X}}\right)^{-\kappa} \frac{\bar{\lambda}^{1/2}}{r} \|u_{\text{hom}}\|_{\underline{L}^2(E_{6\sqrt{dr}})}. \quad (1.35)$$

- First-order corrector estimates. There exist  $\mathbb{Z}^d$ -stationary gradient fields  $\{\nabla \phi_e : e \in \mathbb{R}^d\}$  satisfying the equation

$$-\nabla \cdot \mathbf{a}(e + \nabla \phi_e) = 0 \quad \text{in } \mathbb{R}^d. \quad (1.36)$$

and these satisfy, for every  $e \in \mathbb{R}^d$ ,

$$\|\bar{\mathbf{s}}^{1/2} \nabla \phi_e\|_{\underline{H}^{-1}(E_r)} + \|\bar{\mathbf{s}}^{-1/2}(\mathbf{a}(e + \nabla \phi_e) - \bar{\mathbf{a}} e)\|_{\underline{H}^{-1}(E_r)} \leq C |\bar{\mathbf{s}}^{1/2} e| \left(\frac{r}{\mathcal{X}}\right)^{-\kappa}, \quad \forall r \geq \mathcal{X}. \quad (1.37)$$

Moreover, for every  $\theta \in (0, 1)$ , if we define

$$\mathcal{A}_1(\mathbb{R}^d) := \left\{v \in H_{\mathbf{s}, \text{loc}}^1(\mathbb{R}^d) : -\nabla \cdot \mathbf{a} \nabla v = 0 \text{ in } \mathbb{R}^d, \limsup_{r \rightarrow \infty} r^{-(1+\theta)} \|v\|_{\underline{L}^2(B_r)} = 0\right\}, \quad (1.38)$$

then  $\mathcal{A}_1(\mathbb{R}^d)$  coincides with the set  $\{x \mapsto e \cdot x + \phi_e(x) + c : e \in \mathbb{R}^d, c \in \mathbb{R}\}$ .

- Large-scale regularity. For every  $R \geq \mathcal{X}$  and solution  $u \in H^1(E_R)$  of the equation

$$-\nabla \cdot \mathbf{a} \nabla u = 0 \quad \text{in } E_R, \quad (1.39)$$

we have the estimate

$$\sup_{r \in [\mathcal{X}, R]} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(E_r)} \leq C \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(E_R)}. \quad (1.40)$$

Moreover, there exists  $v \in \mathcal{A}_1(\mathbb{R}^d)$  such that, for every  $\theta \in (0, 1)$ ,

$$\|\mathbf{s}^{1/2} \nabla(u - v)\|_{\underline{L}^2(E_r)} \leq C \left(\frac{r}{R}\right)^\theta \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(E_R)}, \quad (1.41)$$

where  $C_{(1.41)}$  depends on  $\theta$  in addition to  $d$ .

**1.3. Examples satisfying the hypotheses of Theorem B.** The results in this paper have been formulated with near-critical, high contrast problems in mind. However, our general ellipticity assumption (P2) enables us to derive new quantitative homogenization estimates for degenerate and/or unbounded coefficient fields. This assumption also reveals that certain “large contrast” cases can be reinterpreted within our framework as having small contrast. To demonstrate these points, we present three fundamental examples of random fields that satisfy our assumptions (P1), (P2), and (P3). For each example, the results of Theorem B are novel.

The first example is a scalar coefficient field with *Poisson inclusions*. We consider two Poisson point clouds  $\omega_1$  and  $\omega_2$  on  $\mathbb{R}^d$  with intensities  $\rho_1 \geq 0$  and  $\rho_2 \geq 0$ , respectively. Let  $\lambda \in (0, 1]$ ,  $\Lambda \in [1, \infty)$  and define the scalar matrix-valued field

$$\mathbf{a} := (1 + (\Lambda - 1)\mathbf{1}_{B_1} * \omega_1 + (\lambda - 1)\mathbf{1}_{B_1} * \omega_2)\mathbf{I}_d. \quad (1.42)$$

This field clearly satisfies (P1). Since it has a finite range of dependence, it also satisfies (P3) with  $(\beta, \nu) = (0, \frac{d}{2})$  and  $\Psi(t) = \exp(ct^2)$  for some constant  $c(d) > 0$ . The interest is in the ellipticity assumption (P2). As mentioned above, regardless of the values of intensities  $\rho_1$  and  $\rho_2$ , this coefficient field satisfies the uniform ellipticity assumption (1.2) with constants  $\lambda$  and  $\Lambda$ , and therefore (P2) with  $\mathcal{S} = 0$ ,  $\gamma = 0$  and  $\mathbf{E}_0$  is in (1.20).

However, in the case that  $\rho_1$  and  $\rho_2$  are small (perhaps  $10^{-2}$ ) and both  $\Lambda$  and  $\lambda^{-1}$  are very large, we can do better than use the uniform ellipticity condition to check (P2). In this case, the random inclusions are rare and the connected components of their union will be far from percolating. Therefore, while the uniform ellipticity ratio  $\Lambda\lambda^{-1}$  is very large, Theorem B will give a pessimistic bound for the homogenization length scale.

To get a more effective bound, we use the flexibility of the condition (P2). Rather than relying on the uniform ellipticity of the field as a way of checking it, we argue instead that, on a sufficiently large (random) scale (the typical size of which is a power of  $\Lambda\lambda^{-1}$ ), the coefficient field has a coarse-grained ellipticity contrast close to one. Precisely, we have that the following statement, the proof of which appears in Appendix D.1.

**Proposition 1.1** (Poisson inclusions). *There exist constants  $c(d) \in (0, 1]$  and  $C(d) \in [1, \infty)$  such that, if  $\max\{\rho_1, \rho_2\} \leq c$  and  $\gamma \in (0, 1)$ , then the random field  $\mathbf{a}(\cdot)$  defined in (1.42) above satisfies assumptions (P1), (P2) and (P3) with the following parameters:*

$$\mathbf{E}_0 = (1 + C|\log \rho|^{-2})\mathbf{I}_{2d}, \quad \Psi_{\mathcal{S}}(t) = \exp\left(c \max\{\Lambda, \lambda^{-1}\}^{-\frac{1}{d+2}-\frac{\gamma}{d}} t^{\frac{\gamma}{d+2}} - 1\right), \quad \Psi(t) = \exp(ct^2).$$

In particular,  $\Theta \leq \Pi \leq 1 + C|\log \rho|^{-2}$ ,  $K_{\Psi} = C$  and  $K_{\Psi_{\mathcal{S}}} = (C\gamma^{-1})^{\frac{d+2}{\gamma}} \max\{\Lambda, \lambda^{-1}\}^{\frac{1}{\gamma}+1+\frac{2}{d}}$ .

Proposition 1.1 says that, if the intensities of the Poisson processes are small enough, then this seemingly “high contrast” homogenization problem can be treated as a small contrast problem. Consequently, an application of Theorem B implies that, in this case, the length scale for homogenization is proportional to a power of  $\max\{\Lambda, \lambda^{-1}\}$  (with stretched exponential moments).

For the second example, we consider the *advection-diffusion equation*

$$-\lambda\Delta u + \mathbf{b}(x) \cdot \nabla u = 0. \quad (1.43)$$

Here  $\mathbf{b}(x)$  is divergence-free, random vector field which can be written as

$$\mathbf{b} = -\nabla \cdot \mathbf{k}, \quad (1.44)$$

for a stream matrix  $\mathbf{k}$  which is a Gaussian random field taking values in the set  $\mathbb{R}_{\text{skew}}^{d \times d}$  of anti-symmetric matrices. Specifically, assume that each of the entries of  $\mathbf{k}$  is given by the convolution of a fractional Gaussian field with Hurst parameter  $-\sigma \in (-d/2, 0)$  and the standard mollifier (see Appendix D.2 for the definition and explicit construction of the fractional Gaussian fields). Here we do not make any restriction on the covariance structure of the different entries in  $\mathbf{k}$ .

The identity (1.44) allows us to write the equation (1.43) as

$$-\nabla \cdot (\lambda \mathbf{I}_d + \mathbf{k}(x)) \nabla u = 0.$$

Since  $\mathbf{k}$  is a Gaussian random field, it does not belong to  $L^\infty(\mathbb{R}^d)$  and so the equation is not literally uniformly elliptic. However, we show that it satisfies (P2) with an ellipticity ratio of order  $\sigma^{-3} \lambda^{-2}$ .

**Proposition 1.2** (Gaussian stream matrices). *Consider the random field  $\mathbf{a} = \lambda \mathbf{I}_2 + \mathbf{k}$ , where each of the entries of  $\mathbf{k} \in \mathbb{R}_{\text{skew}}^{d \times d}$  is given by the convolution of a fractional Gaussian field with Hurst parameter  $-\sigma \in (-d/2, 0)$  and the standard mollifier. There exists  $C(d) < \infty$  such that the field  $\mathbf{a}(\cdot)$  satisfies the assumptions (P1), (P2) and (P3) with the following parameters:*

$$\begin{cases} \gamma \in (0, \sigma \wedge 1), \\ \mathbf{E}_0 = \begin{pmatrix} 2(\lambda + C\lambda^{-1}\sigma^{-3})\mathbf{I}_d & 0 \\ 0 & 2\lambda^{-1}\mathbf{I}_d \end{pmatrix}, \\ \Psi_S(t) = (\sigma - \gamma) \exp(C^{-1}t^\gamma - C\gamma^{-1}|\log \gamma|), \\ \beta = 1 - 2\sigma/d, \\ \Psi(t) = \Gamma_2(c(\frac{d}{2} - \sigma)t). \end{cases}$$

The proof of Proposition 1.2 appears in Appendix D.2.3.

If the Hurst parameter  $\sigma$  is equal to zero, the  $L^2$  oscillation of the stream matrix  $\mathbf{k}$  is no longer bounded as a function of the scale and the ellipticity is infinite (even in the sense of (P2)). In this case, the equation does not have a diffusive limit and rather exhibits *superdiffusivity*. As we show in [ABRK23], the techniques introduced in this paper are nevertheless up to the task of analyzing it.

We turn next to our third example: *log-normal coefficient fields* which are of the form

$$\mathbf{a}(x) = \exp(h\mathbf{g}(x)),$$

where  $h > 0$  and  $\mathbf{g}$  is a Gaussian field valued in the set  $\mathbb{R}^{d \times d}$  of (not necessarily symmetric) real  $d$ -by- $d$  matrices. For concreteness, we assume that each of the entries of  $\mathbf{g}$  is given by the convolution of a fractional Gaussian field with Hurst parameter  $-\sigma \in (-d/2, 0)$  and the standard mollifier.

**Proposition 1.3** (Log-normal fields). *Consider the random field  $\mathbf{a} = \exp(h\mathbf{g})$ , where each of the entries of  $\mathbf{g} \in \mathbb{R}^{d \times d}$  is given by the convolution of a fractional Gaussian field with Hurst parameter  $-\sigma \in (-d/2, 0)$  and the standard mollifier. There exists  $C(d) < \infty$  such that the field  $\mathbf{a}(\cdot)$  satisfies the assumptions (P1), (P2) and (P3) with the following parameters:*

$$\begin{cases} \gamma \in (0, 1), \\ \mathbf{E}_0 = \exp(C h^2 \sigma^{-2}) \mathbf{I}_{2d} \\ \Psi_S(t) = \exp(C^{-1} h^{-2} \sigma^2 \log^2 t - C h^2 \sigma^{-2} \gamma^{-2}), \\ \beta = 1 - 2\sigma/d, \\ \Psi(t) = \Gamma_2(c(\frac{d}{2} - \sigma)t). \end{cases}$$

The proof of Proposition 1.3 appears in Appendix D.3.

**1.4. An overview of the proof of Theorem B.** In this subsection we explain the main ideas comprising the proof of Theorem B, and explain where they are formalized in the paper. We break the argument into five informal “assertions.”

**Assertion 1.** *The ellipticity condition (P2) is sufficiently strong that it implies basic  $L^2$  elliptic theory on large scales. That is, on scales larger than  $S$ , we get the same basic  $L^2$ -type estimates as in the uniformly elliptic case—with the ellipticity constants  $\lambda$  and  $\Lambda$  in defined in (1.26) taking the place of the usual constants of uniform ellipticity appearing in (1.2).*

This is the main purpose of Section 2. There we introduce the coarse-grained matrices and explore their basic properties, including the basic *coarse-graining inequalities* in (2.35), (2.38), (2.39) and (2.40). These basic estimates allow us to control the spatial averages (and thus the negative Sobolev norms) of the gradients and fluxes of solutions of the equation. Later in the section, we show how these estimates may be combined with the assumption (P2) to obtain coarse-grained versions of the Poincaré and Caccioppoli inequalities—the two basic estimates needed in elliptic regularity theory. See Lemmas 2.11, 2.12 and 2.13.

**Assertion 2.** *The quantity  $\Theta - 1$  quantifies, in a deterministic way, the difference of the solutions of the equation (1.1) and those of the constant-coefficient equation*

$$-\nabla \cdot \mathbf{a}_0 \nabla u = 0,$$

*on scales larger than  $S$ , where  $\mathbf{a}_0 := \mathbf{s}_0 + \mathbf{k}_0$  and  $\mathbf{s}_0$  and  $\mathbf{k}_0$  are as defined in (1.22).*

This is an extension of the previous assertion; here we are saying that the parameter  $\Theta$  defined in (1.25) is a sufficiently good measure of the ellipticity ratio that, if it is close to unity, then (P2) does behave like we expect equations with (uniformly) small contrast to behave. In particular, they are close to solutions of a constant-coefficient equation. This assertion is formalized rigorously in Section 6, using purely deterministic arguments. See Lemma 6.5 for the coarse-graining of the energy density, and Propositions 6.7 and 6.8 for homogenization estimates.

**Assertion 3.** *The ellipticity assumption (P2) is renormalizable. If we let  $\mathbb{P}_{n_0}$  be the pushforward of  $\mathbb{P}$  under the dilation map*

$$\mathbf{a} \mapsto (x \mapsto \mathbf{a}(3^{n_0}x)),$$

*then  $\mathbb{P}_{n_0}$  satisfies (P2) with  $\mathbf{E}_0$  replaced by the expectation  $\overline{\mathbf{A}}(\square_{n_0-l_0})$  of  $\mathbf{A}(\square_{n_0-l_0})$ , where  $l_0$  is a constant which is roughly  $C[\log \Theta]$ . In other words, if we “zoom out” and view the equation from a larger scale, then we have the same assumptions as before, except that the mean of the coarse-grained coefficients (on a slightly smaller scale) take the role of the ellipticity upper bounds.*

The rigorous version of Assertion 3 is stated and proved in Section 2.6: see in particular Proposition 2.6 and Lemma 2.7. It is a relatively simple consequence of the subadditivity of the coarse-grained matrices, combined with an application of assumption (P3).

It is Assertion 3 that gives rise to the renormalization (semi)group. It is natural then to define a scale-dependent notion of ellipticity ratio; we do this by defining  $\Theta_n$  to be the quantity defined analogously to (1.25), but with  $\overline{\mathbf{A}}(\square_n)$  in place of  $\mathbf{E}_0$ : see (2.84).

The subadditivity property of the coarse-grained matrices implies that  $n \mapsto \Theta_n$  is monotone decreasing, and qualitative homogenization implies that it does converge to 1 as  $n \rightarrow \infty$ . Meanwhile, Assertion 2 says that quantitative homogenization estimates will immediately follow once we can give an upper bound on the scale  $n$  such that the quantity  $\Theta_n - 1$  is small.

This leads very naturally to the problems of estimating the size of the scale  $n$  such that  $\Theta_n - 1$  is no larger than  $\frac{1}{2}(\Theta - 1)$ . Such an estimate could then be iterated many times, with the help of Assertion 3 above, to obtain an estimate of the scale on which the renormalized ellipticity ratio is at most  $1 + \delta$ , for  $\delta > 0$  as small as desired.

**Assertion 4.** *If  $\Theta \geq 2$ , then for every  $\sigma \in (0, \frac{1}{2}]$  and  $n \in \mathbb{N}$  satisfying  $n \geq C \log(1 + \sigma^{-1}\Pi) \log \Theta$ ,*

$$\Theta_n - 1 \leq \sigma(\Theta - 1).$$

The precise version of Assertion 4 is stated in Proposition 3.2, and the proof of this proposition is the analytic heart of the paper. It is here that we see the full power of the renormalization and coarse-graining arguments.

The proof is inspired by ideas which originate in [AS16]. That paper, and subsequent works, obtain an inequality which (substantially simplified) states roughly that

$$\Theta_{m+10} - 1 \leq (1 - C_0(d, \Lambda/\lambda)^{-1})(\Theta_m - 1). \quad (1.45)$$

The constant  $C_0(d, \Lambda/\lambda)$  comes from various applications of elliptic estimates and the Poincaré inequality, so it has the form  $C_0(d, \Lambda/\lambda) = C(d) \cdot (\Lambda/\lambda)^p$  for some exponent  $p$ .<sup>4</sup> It is not hard to see that this inequality must be iterated approximately  $C_0(d, \Lambda/\lambda) \log(\Lambda/\lambda)$  many times before the error  $\Theta_m - 1$  is smaller than  $1/2$ . Therefore the upper bound for the length scale of homogenization that this argument gives is roughly

$$\mathcal{X} \lesssim 3^{C_0(d, \Lambda/\lambda) \log(\Lambda/\lambda)} \simeq \exp(CC_0(d, \Lambda/\lambda) \log(\Lambda/\lambda)) \simeq \exp(C(\Lambda/\lambda)^p \log(\Lambda/\lambda)). \quad (1.46)$$

If there is any hope to improve this bound to a sub-exponential bound in the ellipticity ratio, it seems that we need to *remove all dependence on the ellipticity constants* from our argument! This may seem quite hopeless, since elliptic estimates come with dependence on  $\Lambda/\lambda$ , and ellipticity is obviously an important assumption that we need to use.

But an estimate with no dependence on the ellipticity constants is precisely what Assertion 4 gives—with the caveat that we must take a bigger step, say from  $m$  to  $m + C \log(1 + \sigma^{-1}\Pi) \log \Theta$  rather than from  $m$  to  $m + 10$  like in (1.45). This is the only way that dependence on  $\Theta$  or  $\Pi$  is allowed to enter into the proof of Assertion 4: via the scale restriction (the lower bound on  $n$ ). Note that here  $n$  is size of the step in the iteration since, by renormalization (Assertion 3), we can assume  $m = 0$  without loss of generality.

To get rid of the ellipticity dependence, we rely on the coarse-grained version of elliptic estimates summarized in Assertion 1. The idea is to look for a sequence of consecutive scales  $\{n_1, \dots, n_1 + k\}$  such that  $\mathbb{E}[\mathbf{A}(\square_m)]$  does not change much for  $m \in [n_1, n_1 + k]$ . Since this quantity is monotone decreasing in  $m$ , it can only change in one direction and therefore a suitable sequence of consecutive scales can be found by a simple pigeonhole argument. (This pigeonhole argument is the one place where the scale restriction is needed in the proof.) We then argue that, along this finite sequence of scales, the optimizing functions in certain variational formulas for the coarse-grained matrices *must be flat*: that is, their gradients and fluxes must be close to constant functions. This implies, by means of a new *coarse-grained div-curl argument*, that the expectations of the two coarse-grained matrices  $\mathbf{s}(\square)$  and  $\mathbf{s}_*(\square)$  are close to each other (with  $\square$  being the cube on the largest scale in this range of scales). Since the difference  $\mathbb{E}[\mathbf{s}(\square_n) - \mathbf{s}_*(\square_n)]$  actually upper bounds the quantity  $\Theta_n$ , this yields the desired conclusion.

---

<sup>4</sup>Even a single application of an energy estimate such as Caccioppoli's inequality will produce a factor of  $(\Lambda/\lambda)^{1/2}$ , so we would have  $p \geq 1/2$  in (1.46).



If the ellipticity  $\Theta$  is sufficiently close to one, then the statement of Assertion 4 can be improved and the convergence of the renormalized diffusivities to one can be sped up. The idea is essentially that, if  $\Theta - 1$  is small, then we can reduce  $\Theta - 1$  by a factor of two by zooming out only a fixed finite number of scales—provided we are working in a suitable geometry. An iteration then yields an algebraic rate of decay, summarized in the following informal statement.

**Assertion 5.** *There exist  $\sigma_0(d), \alpha(d) \in (0, 1/2]$  and  $C(d) < \infty$  such that, if  $\Theta - 1 \leq \sigma_0$  and*

$$\frac{1}{2}\mathbf{I}_d \leq \mathbf{s}_0 \leq 2\mathbf{I}_d, \quad (1.47)$$

*then we have the estimate*

$$\Theta_m - 1 \leq C3^{-m\alpha}(\Theta - 1), \quad \forall m \in \mathbb{N}. \quad (1.48)$$

Assertion 5 is proved in Section 4.1: see Proposition 4.2.

The five assertions above are assembled into a proof of Theorem B in Section 6.4. The statement of Theorem A is not quite a corollary of Theorem B, since the latter does not state a homogenization result for the Dirichlet problem. We need to use the boundedness of the coefficients (in the context of Theorem A) to handle boundary layers which the general hypotheses of Theorem B give us no means to control. This argument, and the proof of Theorem A, appears in Section 6.2.

In Section 5 we demonstrate that the hypotheses of Theorem B can be further relaxed. We anticipate that such a generalization will be important for applications, and indeed it is necessary for the arguments in [ABRK23].

**1.5. Notation.** We denote  $r \wedge s := \min\{r, s\}$  and  $r \vee s := \max\{r, s\}$ . The Hölder conjugate of an exponent  $p \in [1, \infty]$  is denoted by  $p'$ , where  $p' := p(p-1)^{-1}$  if  $p \neq \infty$  and  $p' := 1$  if  $p = \infty$ . The Euclidean norm on  $\mathbb{R}^m$  is denoted by  $|\cdot|$ . We let  $\mathbb{R}^{m \times n}$  denote the set of  $m$ -by- $n$  matrices with real entries. We let  $B^t$  denote the transpose of a matrix  $B$ . The  $n$ -by- $n$  identity matrix is  $\mathbf{I}_n$ . The symmetric and anti-symmetric  $n$ -by- $n$  matrices are denoted respectively by  $\mathbb{R}_{\text{sym}}^{n \times n}$  and  $\mathbb{R}_{\text{skew}}^{n \times n}$ . The Loewner ordering on  $\mathbb{R}_{\text{sym}}^{n \times n}$  is denoted by  $\leq$ ; that is, if  $A, B \in \mathbb{R}_{\text{sym}}^{n \times n}$  then we write  $A \leq B$  if  $B - A$  has nonnegative eigenvalues. Unless otherwise indicated, the norm we use for  $\mathbb{R}^{m \times n}$ , denoted by  $|A|$ , is the spectral norm, that is, the square root of the largest eigenvalue of  $A^t A$ . The Lebesgue measure of a (measurable) subset  $U \subseteq \mathbb{R}^d$  is  $|U|$ . If  $V$  is a subset of  $\mathbb{R}^d$  of codimension 1 (like the boundary  $\partial U$  of a nice domain  $U$ ), then  $|V|$  is instead the  $(d-1)$ -dimensional Hausdorff measure of  $V$ . Volume-normalized integrals and  $L^p$  norms are denoted, for  $p \in [1, \infty)$ , by

$$(f)_U := \int_U f(x) dx := \frac{1}{|U|} \int_U f(x) dx \quad \text{and} \quad \|f\|_{L^p(U)} := \left( \int_U |f(x)|^p dx \right)^{1/p}. \quad (1.49)$$

We denote by  $|A|$  the cardinality of a finite set  $A$ . A slash through the sum symbol  $\sum$  denotes the average of a finite sequence: for every  $f : A \rightarrow \mathbb{R}$ ,

$$\sum_{a \in A} f(a) := \frac{1}{|A|} \sum_{a \in A} f(a).$$

We denote indicator functions (of events and of sets) by  $\mathbf{1}$ . The standard Hölder spaces are denoted by  $C^{k,\alpha}(U)$  for every  $k \in \mathbb{N}$ ,  $\alpha \in (0, 1]$  and domain  $U \subseteq \mathbb{R}^d$  and Sobolev spaces are denoted by  $W^{s,p}(U)$  for  $s \in \mathbb{R}$  and  $p \in [1, \infty]$ . The fractional Sobolev spaces (for  $s \notin \mathbb{Z}$ ) are defined in [AKM19, Appendix B]. The classical Sobolev space  $W^{1,p}(U)$  is defined by the norm

$$\|f\|_{W^{1,p}(U)} := \left( \|\nabla f\|_{L^p(U)}^p + \|f\|_{L^p(U)}^p \right)^{\frac{1}{p}}.$$

In the case  $p = 2$  this is denoted by  $H^1(U)$ . If  $|U| < \infty$ , we denote the volume-normalized norms  $\|\cdot\|_{\underline{W}^{1,p}(U)}$  by

$$\|f\|_{\underline{W}^{1,p}(U)} := \left( \|\nabla f\|_{\underline{L}^p(U)}^p + |U|^{-\frac{p}{d}} \|f\|_{\underline{L}^p(U)}^p \right)^{\frac{1}{p}}.$$

The negative, dual seminorms are defined by

$$[f]_{\underline{W}^{-1,p'}(U)} := \sup \left\{ \int_U f(x)g(x) dx : g \in C_c^\infty(U), [g]_{\underline{W}^{1,p}(U)} \leq 1 \right\} \quad (1.50)$$

and

$$[f]_{\widehat{\underline{W}}^{-1,p'}(U)} := \sup \left\{ \int_U f(x)g(x) dx : [g]_{\underline{W}^{1,p}(U)} \leq 1, (g)_U = 0 \right\}.$$

If  $p = p' = 2$ , then we write  $H^{-1}$  in place of  $W^{-1,p}$ . We let  $W_0^{1,p}(U)$  denote the closure of  $C_c^\infty(U)$  in  $W^{1,p}(U)$  with respect to the norm  $\|\cdot\|_{W^{1,p}(U)}$ . If  $X(U)$  is a function space defined for every domain  $U \subseteq \mathbb{R}^d$ , then  $X_{\text{loc}}(U)$  denotes the set of functions on  $U$  which belong to  $X(U \cap B_R)$  for every  $R \in [1, \infty)$ . We let  $C_0(\mathbb{R}^d)$  denote the space of continuous functions  $u : \mathbb{R}^d \rightarrow \mathbb{R}$  such that  $\lim_{|x| \rightarrow \infty} u(x) = 0$ , and  $C_c^k(\mathbb{R}^d)$  denotes the subspace of  $C^k(\mathbb{R}^d)$  with compact support in  $\mathbb{R}^d$ . We use the  $\mathcal{O}_\Psi(\cdot)$  notation defined in Section C to keep track of the stochastic integrability of our random variables. Throughout, for  $\sigma \in (0, \infty)$  we denote  $\Gamma_\sigma(t) := \exp(t^\sigma)$ . The bold symbol  $\mathbf{\Gamma}$  denotes the gamma function  $\mathbf{\Gamma}(s) := \int_0^\infty t^{s-1} \exp(-t) dt$ .

## 2. The coarse-grained diffusion matrices

In this section we introduce the coarse-grained diffusion matrices that form the basis of our approach in this paper. These quantities are not new here, and go back to the works [AS16, AM16]. The novelty in this paper lies in the precise way they are used to renormalize the equation. (For historical context and a more complete presentation of some of the material covered in this section, we refer to [AK24, Chapters 4 & 5].)

There are various possible definitions for the “coarse-grained diffusion matrix,” and the ones we introduce are not the only plausible choices. Particularly in the general nonsymmetric case, our definitions may initially seem counterintuitive. However, these specific quantities are crucial for proving results such as Theorem A. They exhibit a complex algebraic structure and possess essential properties that facilitate coarse-graining. Attempting to substitute alternative notions of “box diffusivity” into our arguments would result in failure. To Paraphrase Steven Weinberg’s Third Law of Progress [Wei83], *you may use any quantities you like to study elliptic homogenization, but if you use the wrong ones, you’ll be sorry.*

Given a bounded domain  $U \subseteq \mathbb{R}^d$ , we will define two symmetric matrices  $\mathbf{s}_*(U)$  and  $\mathbf{s}(U)$ , which we think of as two competing coarse-grained versions of the symmetric part  $\mathbf{s}(\cdot)$  of the coefficient field, and another matrix  $\mathbf{k}(U)$  which may not be antisymmetric but we still consider to be the coarse-grained version of the anti-symmetric part. The two symmetric matrices satisfy the ordering  $\mathbf{s}_*(U) \leq \mathbf{s}(U)$ , as we will show, and we think of the pair as giving us lower and upper bounds for the coarse-grained symmetric part—with their difference representing the uncertainty (or error) in the coarse-graining procedure.

There are several equivalent ways of defining them and thinking about these coarse-grained matrices. They can first be arranged in a  $2d$ -by- $2d$  matrix as

$$\mathbf{A}(U) := \begin{pmatrix} (\mathbf{s} + \mathbf{k}^t \mathbf{s}_*^{-1} \mathbf{k})(U) & -(\mathbf{k}^t \mathbf{s}_*^{-1})(U) \\ -(\mathbf{s}_*^{-1} \mathbf{k})(U) & \mathbf{s}_*^{-1}(U) \end{pmatrix}. \quad (2.1)$$

This matrix  $\mathbf{A}(U)$  can be considered to be a coarse-graining of the field  $\mathbf{A}(x)$  defined in (1.12). This may seem at first strange and unfamiliar, but thinking in terms of  $\mathbf{A}(U)$  (as opposed to  $\mathbf{s}(U)$ ,  $\mathbf{s}_*(U)$  and  $\mathbf{k}(U)$ ) has many algebraic and analytic advantages. It is also very natural from the *variational* point of view, as indeed coarse-grained matrices have a variational interpretation. In fact they have several variational interpretations: in terms of the quantity  $J$  and  $J^*$  in (2.10) and (2.11), and the “double variable” analogue in (2.16).

The coarse-grained matrices have a rich structure and many interesting properties. In this section, we will list the facts that are used in this paper, while omitting some of the proofs of the more basic properties (each of which can be found in [AK24, Section 5]). We also prove an important statement about renormalizations of the assumptions (see Section 2.6) as well as provide some functional inequalities which indicate that the assumption of (P2) is a good notion of a scale-dependent ellipticity condition (see Section 2.8).

**2.1. Basic Sobolev space framework.** Recall that  $\mathbf{a} \in \Omega$  means that  $\mathbf{s}$ ,  $\mathbf{s}^{-1}$  and  $\mathbf{k}^t \mathbf{s}^{-1} \mathbf{k}$  have entries belonging to  $L^1_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^d \times \mathbb{R}^d)$ , where  $\mathbf{s}$  and  $\mathbf{k}$  are respectively the symmetric and anti-symmetric parts of  $\mathbf{a}$ . Equivalent, we can consider  $\mathbf{A}$  to be the canonical element, in which case of  $\mathbf{A} \in \Omega$  means simply that  $\mathbf{A} \in L^1_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^{2d \times 2d})$ . It is important to keep in mind that this is the minimal *qualitative* requirement for our coefficients fields. Our main results require, of course, quantitative ellipticity assumptions, namely (P2). As we will show, this ensures that the solutions of the equation are much better behaved compared to what we can show under the qualitative assumption. However, to even define the coarse-grained coefficients appearing in the quantitative ellipticity assumption we must introduce some basic notions from elliptic theory, which are somewhat nonstandard due to the general qualitative setting which allows for unbounded and highly degenerate equations. For this reason we give a thorough (if succinct) presentation.

For each  $\mathbf{a} \in \Omega$  and subset  $U \subseteq \mathbb{R}^d$ , we define the function spaces  $H^1_{\mathbf{s}}(U)$  as the completion of  $C^\infty(U)$  with respect to the norm

$$\|u\|_{H^1_{\mathbf{s}}(U)} := \left( \|u\|_{L^1(U)}^2 + \int_U \nabla u \cdot \mathbf{s} \nabla u \right)^{1/2}. \quad (2.2)$$

Observe that, by Hölder’s inequality, we have that

$$u \in H^1_{\mathbf{s}}(U) \implies \nabla u, \mathbf{a} \nabla u \in L^1(U). \quad (2.3)$$

Indeed,  $u \in H^1_{\mathbf{s}}(U)$  implies that  $\mathbf{s}^{1/2} \nabla u \in L^2(U)$  and the assumption of  $\mathbf{a} \in \Omega$  implies that  $\mathbf{s}^{1/2}$ ,  $\mathbf{s}^{-1/2}$  and  $\mathbf{k} \mathbf{s}^{-1/2}$  also belong to  $L^2(U)$ . This together with Cauchy-Schwarz give the implication (2.3). According to [KO84, Theorem 1.11]<sup>5</sup>, the space  $H^1_{\mathbf{s}}(U)$  is a complete (Hilbert) space for every  $\mathbf{a} \in \Omega$  and  $U \subseteq \mathbb{R}^d$ . It is clear that

$$C_c^\infty(U) \subseteq H^1_{\mathbf{s}}(U).$$

We also define the subspace of “trace zero” functions by

$$H^1_{\mathbf{s},0}(U) := \text{closure of } C_c^\infty(U) \text{ with respect to } \|\cdot\|_{H^1_{\mathbf{s}}(U)}.$$

The linear subspace of  $H^1_{\mathbf{s},0}(U)$  consisting of solutions of the equation  $\nabla \cdot \mathbf{a} \nabla u = 0$  is denoted by

$$\mathcal{A}(U) := \{u \in H^1_{\mathbf{s}}(U) : \nabla \cdot \mathbf{a} \nabla u = 0 \text{ in } U\}.$$

---

<sup>5</sup>The paper [KO84] considers the case of scalar  $\mathbf{s}$ , but the proof easily generalizes to general (matrix-valued) case.

Here the equation  $\nabla \cdot \mathbf{a} \nabla u = 0$  is to be understood in the sense of distributions; that is,

$$\int_U \nabla \psi \cdot \mathbf{a} \nabla u = 0, \quad \forall \psi \in C_c^\infty(U).$$

Things are now set up correctly for the application of the Riesz representation theorem to the Dirichlet problem

$$\begin{cases} -\nabla \cdot \mathbf{s} \nabla v = f & \text{in } U, \\ v = 0 & \text{on } \partial U. \end{cases} \quad (2.4)$$

We deduce that, for every bounded domain  $U \subseteq \mathbb{R}^d$  and element  $f$  of the dual space of  $H_{\mathbf{s},0}^1(U)$ , the boundary-value problem (2.4) has a unique solution  $v \in H_{\mathbf{s},0}^1(U)$ . This means that  $v$  belongs to  $H_{\mathbf{s},0}^1(U)$  and, for every  $u \in H_{\mathbf{s},0}^1(U)$ ,

$$\int_U \nabla u \cdot \mathbf{s} \nabla v = \langle u, f \rangle,$$

with the brackets  $\langle \cdot, \cdot \rangle$  denoting the pairing between  $H_{\mathbf{s},0}^1(U)$  and its dual. We deduce that the dual space of  $H_{\mathbf{s},0}^1(U)$ , which we denote by  $H_{\mathbf{s}}^{-1}(U)$ , can be characterized as

$$H_{\mathbf{s}}^{-1}(U) := \{ \nabla \cdot \mathbf{s}^{1/2} \mathbf{f} : \mathbf{f} \in L^2(U)^d \}.$$

Indeed, the inclusion  $\supseteq$  is obvious and the reverse inclusion  $\subseteq$  follows from the solvability of (2.4). We define the dual norm  $\| \cdot \|_{H_{\mathbf{s}}^{-1}(U)}$  by

$$\|f\|_{H_{\mathbf{s}}^{-1}(U)} := \sup \{ \langle u, f \rangle : u \in H_{\mathbf{s},0}^1(U), \|u\|_{H_{\mathbf{s},0}^1(U)} \leq 1 \}$$

We often abuse notation by writing  $\int_U u f$  in place of  $\langle u, f \rangle$  when  $u \in H_{\mathbf{s},0}^1(U)$  and  $f \in H_{\mathbf{s}}^{-1}(U)$ .

We next discuss the solvability of the (non necessarily self-adjoint) equation  $-\nabla \cdot \mathbf{a} \nabla u = 0$  in every bounded domain  $U \subseteq \mathbb{R}^d$ . For this purpose we introduce the norm

$$\|u\|_{H_{\mathbf{a}}^1(U)} := \left( \|u\|_{H_{\mathbf{s}}^1(U)}^2 + \|\nabla \cdot \mathbf{k} \nabla u\|_{H_{\mathbf{s}}^{-1}(U)}^2 \right)^{1/2}, \quad (2.5)$$

and we let  $H_{\mathbf{a}}^1(U)$  denote the closure of  $C^\infty(U)$  with respect to the norm  $\| \cdot \|_{H_{\mathbf{a}}^1(U)}$ . We also let  $H_{\mathbf{a},0}^1(U)$  denote the closure of  $C_c^\infty(U)$  with respect to  $\| \cdot \|_{H_{\mathbf{a}}^1(U)}$ .

The Lions-Lax-Milgram lemma (see [Sho97, Theorem 2.1, page 109]) implies, for every  $f \in H_{\mathbf{s}}^{-1}(U)$ , the existence of a unique solution  $u \in H_{\mathbf{a},0}^1(U)$  of the Dirichlet problem

$$\begin{cases} -\nabla \cdot \mathbf{a} \nabla u = f & \text{in } U, \\ u = 0 & \text{on } \partial U. \end{cases} \quad (2.6)$$

Being a solution of (2.6) means that  $u \in H_{\mathbf{a},0}^1(U)$  and  $u$  satisfies

$$\int_U \nabla w \cdot \mathbf{s} \nabla u = \langle \nabla \cdot \mathbf{k} \nabla u + f, w \rangle, \quad \forall w \in H_{\mathbf{s},0}^1(U).$$

We interpret this simply as

$$\int_U \nabla w \cdot \mathbf{a} \nabla u = \langle w, f \rangle, \quad \forall w \in H_{\mathbf{s},0}^1(U).$$

Similarly, the Lions-Lax-Milgram lemma implies the well-posedness of the Neumann problem. We introduce the space

$$L^2_{\text{sol},\mathbf{s}}(U) := \{\mathbf{g} : \mathbf{s}^{-1/2}\mathbf{g} \in L^2(U)^d, \nabla \cdot \mathbf{g} = 0\}$$

and we let  $\hat{H}_{\mathbf{s}}^{-1}(U)$  be the closure of  $C_c^\infty(U)$  with respect to the norm

$$\|f\|_{\hat{H}_{\mathbf{s}}^{-1}(U)} := \sup \left\{ \int_U f u : u \in H_{\mathbf{s}}^1(U), \|u\|_{H_{\mathbf{s}}^1(U)} \leq 1 \right\}.$$

Since constant functions belong to  $H_{\mathbf{s}}^1(U)$ , each element  $f \in \hat{H}_{\mathbf{s}}^{-1}(U)$  has a well-defined mean value on  $U$  which we denote by  $(f)_U$ . For every  $f \in \hat{H}_{\mathbf{s}}^{-1}(U)$  with  $(f)_U = 0$  and  $\mathbf{g} \in L^2_{\text{sol},\mathbf{s}}(U)$ , there exists a unique  $u \in H_{\mathbf{a}}^1(U)$  satisfying  $(u)_U = 0$  and

$$\begin{cases} -\nabla \cdot \mathbf{a} \nabla u = f & \text{in } U, \\ \mathbf{n} \cdot (\mathbf{a} \nabla u - \mathbf{g}) = 0 & \text{on } \partial U, \end{cases} \quad (2.7)$$

where  $\mathbf{n}$  is the outward-pointing unit normal vector on  $\partial U$ . The interpretation of (2.7) is that

$$\int_U \nabla w \cdot (\mathbf{a} \nabla u - \mathbf{g}) = \langle w, f \rangle, \quad \forall w \in H_{\mathbf{s}}^1(U).$$

The soft analysis discussed above, which assumes only  $\mathbf{a} \in \Omega$ , is very limited. We are unable to perform basic energy estimates, or even test the equation with the solutions multiplied by a cutoff function, because we are unable to show that the product  $\varphi u$  belongs to  $H_{\mathbf{s}}^1(U)$ , even if  $u \in H_{\mathbf{s}}^1(U)$  and  $\varphi \in C_c^\infty(U)$ .

In order to address this issue and proceed further, we will need some basic Sobolev-type embeddings for our space  $H_{\mathbf{s}}^1$ , and this requires a stronger assumption on the coefficient field  $\mathbf{a}(\cdot)$  beyond that  $\mathbf{a} \in \Omega$ . As it turns out, certain bounds on the coarse-grained coefficients—which are implied by assumption (P2)—provide exactly what is needed. These Sobolev-type embeddings are presented below in Section 2.8 (see Lemma 2.11 for the embeddings and Lemma 2.13 for the justification of testing). However, we must first introduce the coarse-grained matrices and explore their basic properties.

**2.2. The coarse-grained matrices: definitions and basic properties.** For every realization of the coefficients  $\mathbf{a} \in \Omega$  and bounded Lipschitz domain  $U \subseteq \mathbb{R}^d$  we associate three matrices  $\mathbf{s}(U)$ ,  $\mathbf{s}_*(U)$  and  $\mathbf{k}(U)$ . The matrices  $\mathbf{s}(U)$  and  $\mathbf{s}_*(U)$  are symmetric, invertible and satisfy the ordering  $\mathbf{s}_*(U) \leq \mathbf{s}(U)$ . Together this pair represents the symmetric part of coarse-grained field, and the is a quantification of the “uncertainty” of the coarse-graining, as we will see below. The matrix  $\mathbf{k}(U)$  represents the anti-symmetric part of the coarse-grained field. It is not necessarily anti-symmetric, in general, but its symmetric part  $\frac{1}{2}(\mathbf{k} + \mathbf{k}^t)(U)$  is bounded by the gap  $\mathbf{s}(U) - \mathbf{s}_*(U)$ , as will be shown below, and is thus bounded by the uncertainty.

There are several equivalent ways to define the matrices  $\mathbf{s}(U)$ ,  $\mathbf{s}_*(U)$  and  $\mathbf{k}(U)$  and here we opt for a variational formulation. We first introduce the quantity  $J(U, p, q)$ , which is defined for and  $p, q \in \mathbb{R}^d$  by

$$J(U, p, q) := \sup_{u \in \mathcal{A}(U)} \int_U \left( -\frac{1}{2} \nabla u \cdot \mathbf{s} \nabla u - p \cdot \mathbf{a} \nabla u + q \cdot \nabla u \right). \quad (2.8)$$

Notice that  $J(U, p, q)$  is well-defined by the discussion in the previous subsection. In particular, the integrand in (2.8) belongs to  $L^1(U)$  for each  $\mathbf{a} \in \Omega$  and  $u \in \mathcal{A}(U)$ . We also define the analogue of this quantity for the adjoint operator by

$$J^*(U, p, q) := \sup_{u \in \mathcal{A}^*(U)} \int_U \left( -\frac{1}{2} \nabla u \cdot \mathbf{s} \nabla u - p \cdot \mathbf{a}^t \nabla u + q \cdot \nabla u \right) \quad (2.9)$$

where

$$\mathcal{A}^*(U) := \{u \in H_{\mathbf{a}}^1(U) : -\nabla \cdot \mathbf{a}^t \nabla u = 0 \text{ in } U\}$$

denotes the set of solutions to the adjoint equation in the domain  $U$ . The supremums in the variational problems on the right sides of (2.8) and (2.9) are achieved, and the maximizers belong to  $H_{\mathbf{a}}^1(U)$  and are unique up to additive constants. We denote them by  $v(\cdot, U, p, q)$  and  $v^*(\cdot, U, p, q)$ , respectively.

The mapping  $(p, q) \mapsto J(U, p, q)$  is quadratic, and it is convenient to write it using matrices. We let  $\mathbf{s}(U), \mathbf{s}_*(U) \in \mathbb{R}_{\text{sym}}^{d \times d}$  and  $\mathbf{k}(U) \in \mathbb{R}^{d \times d}$  be defined in a such a way that the following relation is satisfied:

$$J(U, p, q) = \frac{1}{2} p \cdot \mathbf{s}(U) p + \frac{1}{2} (q + \mathbf{k}(U) p) \cdot \mathbf{s}_*^{-1}(U) (q + \mathbf{k}(U) p) - p \cdot q. \quad (2.10)$$

It turns out that  $J^*$  can be written by means of the same matrices; it satisfies

$$J^*(U, p, q) = \frac{1}{2} p \cdot \mathbf{s}(U) p + \frac{1}{2} (q - \mathbf{k}(U) p) \cdot \mathbf{s}_*^{-1}(U) (q - \mathbf{k}(U) p) - p \cdot q. \quad (2.11)$$

In other words, when we coarse-grain the adjoint  $\mathbf{a}^t(\cdot)$  of the field  $\mathbf{a}(\cdot)$ , we leave  $\mathbf{s}(U)$  and  $\mathbf{s}_*(U)$  unchanged and exchange  $\mathbf{k}(U)$  with  $-\mathbf{k}(U)$ . This is a non-obvious fact that cannot be deduced using algebra alone: it follows from the dual variational principles (2.12) and (2.19) and below: see [AK24, Lemma 5.4] for a proof.

We collect the coarse-grained matrices into a single  $2d$ -by- $2d$  matrix by defining  $\mathbf{A}(U)$  as in (2.1). This larger matrix can be thought of as a coarse-graining of the matrix in (1.12), and it has the following variational interpretation (see [AK24, Lemma 5.3]), which gives an alternative way of defining the coarse-grained matrices. We have the formula

$$\frac{1}{2} P \cdot \mathbf{A}(U) P = \inf \left\{ \int_U \frac{1}{2} (X + P) \cdot \mathbf{A} (X + P) : X \in L_{\mathbf{a}, \text{pot}, 0}^2(U) \times L_{\mathbf{a}, \text{sol}, 0}^2(U) \right\}. \quad (2.12)$$

where  $L_{\mathbf{a}, \text{pot}, 0}^2(U)$  is defined as the closure of the set  $\{\nabla \phi : \phi \in C_c^\infty(U)\}$  of smooth, compactly supported gradients with respect to the norm  $\mathbf{f} \mapsto (\int_U \mathbf{f} \cdot \mathbf{s} \mathbf{f})^{1/2}$ , and  $L_{\mathbf{a}, \text{sol}, 0}^2(U)$  is the closure of the set  $\{\mathbf{f} : \mathbf{f} \in C_c^\infty(U; \mathbb{R}^d), \nabla \cdot \mathbf{f} = 0\}$  of smooth, compactly supported divergence-free fields with respect to the norm  $\mathbf{f} \mapsto (\int_U \mathbf{f} \cdot \mathbf{s}^{-1} \mathbf{f})^{1/2}$ .

The coarse-grained quantity  $\mathbf{A}(U)$  has the same information as  $J$  and  $J^*$ , or equivalently the coarse-grained matrices  $\mathbf{s}(U)$ ,  $\mathbf{s}_*(U)$  and  $\mathbf{k}(U)$ . By straightforward algebraic manipulations, we observe that the identities (2.10) and (2.11) are equivalent to

$$J(U, p, q) = \frac{1}{2} \begin{pmatrix} -p \\ q \end{pmatrix} \cdot \mathbf{A}(U) \begin{pmatrix} -p \\ q \end{pmatrix} - p \cdot q \quad \text{and} \quad J^*(U, p, q) = \frac{1}{2} \begin{pmatrix} p \\ q \end{pmatrix} \cdot \mathbf{A}(U) \begin{pmatrix} p \\ q \end{pmatrix} - p \cdot q. \quad (2.13)$$

It is sometimes helpful to refer to the top-left  $d$ -by- $d$  block of  $\mathbf{A}(U)$ , so we define

$$\mathbf{b}(U) := (\mathbf{s} + \mathbf{k}^t \mathbf{s}_*^{-1} \mathbf{k})(U). \quad (2.14)$$

We “double the variables” by combining  $J$  and  $J^*$  into a single quantity by defining

$$\mathbf{J}\left(U, \begin{pmatrix} p \\ q \end{pmatrix}, \begin{pmatrix} q^* \\ p^* \end{pmatrix}\right) := \frac{1}{2}J(U, p - p^*, q^* - q) + \frac{1}{2}J^*(U, p^* + p, q^* + q). \quad (2.15)$$

By (2.13) and some straightforward algebraic manipulations, the definition (2.15) is equivalent to

$$\mathbf{J}(U, P, Q) = \frac{1}{2}P \cdot \mathbf{A}(U)P + \frac{1}{2}Q \cdot \mathbf{A}_*^{-1}(U)Q - P \cdot Q, \quad \forall P, Q \in \mathbb{R}^{2d}, \quad (2.16)$$

where the matrix  $\mathbf{A}_*^{-1}(U)$  is defined by swapping the rows and columns of  $\mathbf{A}(U)$ :

$$\mathbf{A}_*^{-1}(U) := \begin{pmatrix} \mathbf{s}_*^{-1}(U) & -(\mathbf{s}_*^{-1}\mathbf{k})(U) \\ -(\mathbf{k}^t\mathbf{s}_*^{-1})(U) & (\mathbf{s} + \mathbf{k}^t\mathbf{s}_*^{-1}\mathbf{k})(U) \end{pmatrix}. \quad (2.17)$$

The following formulas for the inverses of  $\mathbf{A}(U)$  and  $\mathbf{A}_*^{-1}(U)$  are obtained by a direct computation:

$$\begin{cases} \mathbf{A}^{-1}(U) = \begin{pmatrix} \mathbf{s}^{-1}(U) & (\mathbf{s}^{-1}\mathbf{k}^t)(U) \\ (\mathbf{k}\mathbf{s}^{-1})(U) & (\mathbf{s}_* + \mathbf{k}\mathbf{s}^{-1}\mathbf{k}^t)(U) \end{pmatrix}, \\ \mathbf{A}_*(U) = \begin{pmatrix} (\mathbf{s}_* + \mathbf{k}\mathbf{s}^{-1}\mathbf{k}^t)(U) & (\mathbf{k}\mathbf{s}^{-1})(U) \\ (\mathbf{s}^{-1}\mathbf{k}^t)(U) & \mathbf{s}^{-1}(U) \end{pmatrix}. \end{cases} \quad (2.18)$$

The quantity  $\mathbf{J}(U, P, Q)$  also has the following variational formulation, which is easy to check (or see [AK24, Lemma 5.2]): for every  $P, Q \in \mathbb{R}^{2d}$ ,

$$\mathbf{J}(U, P, Q) = \sup_{X \in \mathcal{S}(U)} \int_U \left( -\frac{1}{2}X \cdot \mathbf{A}X - P \cdot \mathbf{A}X + Q \cdot X \right), \quad (2.19)$$

where the space  $\mathcal{S}(U)$  is defined by

$$\mathcal{S}(U) := \left\{ \begin{pmatrix} \nabla v + \nabla v^* \\ \mathbf{a}\nabla v - \mathbf{a}^t\nabla v^* \end{pmatrix} : v \in \mathcal{A}(U), v^* \in \mathcal{A}^*(U) \right\}. \quad (2.20)$$

The coarse-grained objects defined above have a rich structure. The properties above, as well as those we list below, can be found in [AK24, Section 5.1].

We continue by discussing basic upper and lower bounds. The matrices  $\mathbf{s}(U)$  and  $\mathbf{s}_*(U)$  are ordered, and the coarse-grained matrices  $\mathbf{s}_*(U)$  and  $\mathbf{b}(U)$  are bounded from above and below by the averages of the field. We have that

$$\left( \int_U \mathbf{s}^{-1}(x) dx \right)^{-1} \leq \mathbf{s}_*(U) \leq \mathbf{s}(U) \leq \mathbf{b}(U) \leq \int_U (\mathbf{s} + \mathbf{k}^t\mathbf{s}_*^{-1}\mathbf{k})(x) dx. \quad (2.21)$$

Each of the inequalities in (2.21) is very easy to prove with the exception of  $\mathbf{s}_*(U) \leq \mathbf{s}(U)$ , which is a consequence of the identities (2.10), (2.11) and a duality argument: see [AK24, Lemma 5.4] and the discussion following it. The other bounds (2.21) can be written more compactly and also more generally in terms of the  $2d$  block matrices:

$$\left( \int_U \mathbf{A}^{-1}(x) dx \right)^{-1} \leq \mathbf{A}_*(U) \leq \mathbf{A}(U) \leq \int_U \mathbf{A}(x) dx. \quad (2.22)$$

The matrix  $\mathbf{A}(U)$  and its inverse also satisfy, for every  $\eta > 0$ ,

$$\begin{cases} \mathbf{A}(U) \leq \begin{pmatrix} (\mathbf{s} + (1 + \eta^{-1})\mathbf{k}^t\mathbf{s}_*^{-1}\mathbf{k})(U) & 0 \\ 0 & (1 + \eta)\mathbf{s}_*^{-1}(U) \end{pmatrix}, \\ \mathbf{A}^{-1}(U) \leq \begin{pmatrix} (1 + \eta)\mathbf{s}^{-1}(U) & 0 \\ 0 & (\mathbf{s}_* + (1 + \eta^{-1})\mathbf{k}\mathbf{s}^{-1}\mathbf{k}^t)(U) \end{pmatrix}. \end{cases} \quad (2.23)$$



Another important property of the quantity  $J(U, p, q)$  is subadditivity. We write this in terms of the  $2d$  block matrices as follows. For every bounded Lipschitz domain  $U \subseteq \mathbb{R}^d$  and disjoint partition  $\{U_i\}_{i=1, \dots, N}$  of  $U$  (up to a zero Lebesgue measure set), we have

$$\mathbf{A}(U) \leq \sum_{i=1}^N \frac{|U_i|}{|U|} \mathbf{A}(U_i) \quad \text{and} \quad \mathbf{A}_*^{-1}(U) \leq \sum_{i=1}^N \frac{|U_i|}{|U|} \mathbf{A}_*^{-1}(U_i). \quad (2.24)$$

The bounds in (2.24) should be regarded as a generalization of (2.22), a coarse-grained version of the latter. Note that while each of  $\mathbf{A}(U)$ ,  $\mathbf{A}_*^{-1}(U)$ ,  $\mathbf{b}(U)$  and  $\mathbf{s}_*^{-1}(U)$  is subadditive, but neither  $\mathbf{s}(U)$  nor  $\mathbf{k}(U)$  is subadditive in any sense.

By [AK24, Remark 5.7], we have that the symmetric part of  $\mathbf{k}$  is controlled by the gap between  $\mathbf{s}(U)$  and  $\mathbf{s}_*(U)$ :

$$(\mathbf{k} + \mathbf{k}^t)(U) \leq (\mathbf{s} - \mathbf{s}_*)(U) \quad \text{and} \quad -(\mathbf{k} + \mathbf{k}^t)(U) \leq (\mathbf{s} - \mathbf{s}_*)(U). \quad (2.25)$$

This is also proved below in (2.58). The difference of  $\mathbf{s}(U)$  and  $\mathbf{s}_*(U)$  can also be expressed by means of  $J$  and  $J^*$  via the identity

$$e \cdot (\mathbf{s} - \mathbf{s}_*)(U)e = J(U, e, (\mathbf{s}_* - \mathbf{k})(U)e) + J^*(U, e, (\mathbf{s}_* + \mathbf{k})(U)e). \quad (2.26)$$

More generally, we have that by [AK24, Lemma 5.2], for every  $\tilde{\mathbf{s}} \in \mathbb{R}_{\text{sym}}^{d \times d}$  and  $\tilde{\mathbf{k}} \in \mathbb{R}^{d \times d}$ , we have

$$\begin{aligned} e \cdot (\mathbf{s}(U) - \mathbf{s}_*(U))e + (\tilde{\mathbf{s}} - \mathbf{s}_*(U))e \cdot \mathbf{s}_*^{-1}(U)(\tilde{\mathbf{s}} - \mathbf{s}_*(U))e + (\tilde{\mathbf{k}} - \mathbf{k}(U))e \cdot \mathbf{s}_*^{-1}(U)(\tilde{\mathbf{k}} - \mathbf{k}(U))e \\ = J(U, e, (\tilde{\mathbf{s}} - \tilde{\mathbf{k}})e) + J^*(U, e, (\tilde{\mathbf{s}} + \tilde{\mathbf{k}})e). \end{aligned} \quad (2.27)$$

We next explore properties of the coarse-grained matrices that give us information about general solutions. The first variation of the optimization problem in (2.8) is

$$q \cdot \oint_U \nabla w - p \cdot \oint_U \mathbf{a} \nabla w = \oint_U \nabla w \cdot \mathbf{s} \nabla v(\cdot, U, p, q), \quad \forall w \in \mathcal{A}(U). \quad (2.28)$$

The second variation says that

$$\begin{aligned} J(U, p, q) - \oint_U \left( -\frac{1}{2} \nabla w \cdot \mathbf{s} \nabla w - p \cdot \mathbf{a} \nabla w + q \cdot \nabla w \right) \\ = \oint_U \frac{1}{2} (\nabla v(\cdot, U, p, q) - \nabla w) \cdot \mathbf{s} (\nabla v(\cdot, U, p, q) - \nabla w), \quad \forall w \in \mathcal{A}(U). \end{aligned} \quad (2.29)$$

By taking  $w = 0$  in (2.29), it follows that  $J$  can be expressed as the energy of its maximizer:

$$J(U, p, q) = \oint_U \frac{1}{2} \nabla v(\cdot, U, p, q) \cdot \mathbf{s} \nabla v(\cdot, U, p, q). \quad (2.30)$$

We can read off the spatial averages of the gradient and flux of the maximizer  $v(\cdot, U, p, q)$  from the quantity  $J$  itself. We have

$$\begin{cases} \oint_U \nabla v(\cdot, U, p, q) = -p + \mathbf{s}_*^{-1}(U)(q + \mathbf{k}(U)p) \\ \oint_U \mathbf{a} \nabla v(\cdot, U, p, q) = (\mathbf{I}_d - \mathbf{k}^t \mathbf{s}_*^{-1})(U)q - \mathbf{b}(U)p. \end{cases} \quad (2.31)$$

These identities play a central role in the analysis in the paper. It will be convenient to write them in a more compact way, using matrix notation. We introduce the matrix

$$\mathbf{R} := \begin{pmatrix} 0 & \mathbf{I}_d \\ \mathbf{I}_d & 0 \end{pmatrix} \quad (2.32)$$

and then observe that (2.33) is equivalent to

$$\oint_U \begin{pmatrix} \nabla v \\ \mathbf{a} \nabla v \end{pmatrix} (\cdot, U, p, q) = (\mathbf{R} \mathbf{A}(U) + \mathbf{I}_{2d}) \begin{pmatrix} -p \\ q \end{pmatrix}. \quad (2.33)$$

The quantity  $J$  allows us to relate the spatial averages of gradients and fluxes of arbitrary solutions: for every  $p, q \in \mathbb{R}^d$  and  $w \in \mathcal{A}(U)$ , we have by (2.28), (2.30) and Hölder's inequality that

$$\left| \oint_U (p \cdot \mathbf{a} \nabla w - q \cdot \nabla w) \right| = \left| \oint_U \nabla w \cdot \mathbf{s} \nabla v(\cdot, U, p, q) \right| \leq (2J(U, p, q))^{1/2} \left( \oint_U \nabla w \cdot \mathbf{s} \nabla w \right)^{1/2}. \quad (2.34)$$

This inequality is useful when  $J(U, p, q)$  is small, which requires  $q$  and  $p$  to be related and the gap between  $\mathbf{s}(U)$  and  $\mathbf{s}_*(U)$  to be small. Indeed, choosing  $q = (\mathbf{s}_* - \mathbf{k}^t)(U)p$  and taking the supremum over  $|p| = 1$  yields, in view of (2.26),

$$\left| \oint_U \mathbf{a} \nabla w - (\mathbf{s}_* - \mathbf{k}^t)(U) \oint_U \nabla w \right| \leq 2^{1/2} |(\mathbf{s} - \mathbf{s}_*)(U)|^{1/2} \left( \oint_U \nabla w \cdot \mathbf{s} \nabla w \right)^{1/2}. \quad (2.35)$$

This motivates the definition

$$\mathbf{a}_*(U) := \mathbf{s}_*(U) - \mathbf{k}^t(U). \quad (2.36)$$

We can then write the previous inequality as

$$\left| \oint_U (\mathbf{a}_*(U) - \mathbf{a}) \nabla w \right| \leq 2^{1/2} |(\mathbf{s} - \mathbf{s}_*)(U)|^{1/2} \left( \oint_U \nabla w \cdot \mathbf{s} \nabla w \right)^{1/2}. \quad (2.37)$$

The coarse-grained matrix  $\mathbf{s}_*(U)$  gives a lower bound for the spatial average of the gradient of an arbitrary solution in terms of its energy:

$$\frac{1}{2} \left( \oint_U \nabla u \right) \cdot \mathbf{s}_*(U) \left( \oint_U \nabla u \right) \leq \oint_U \frac{1}{2} \nabla u \cdot \mathbf{s} \nabla u, \quad \forall u \in \mathcal{A}(U). \quad (2.38)$$

Similarly, the coarse-grained matrix  $\mathbf{b}(U)$  gives a lower bound for the spatial average of the flux of an arbitrary solution in terms of its energy:

$$\frac{1}{2} \left( \oint_U \mathbf{a} \nabla u \right) \cdot \mathbf{b}^{-1}(U) \left( \oint_U \mathbf{a} \nabla u \right) \leq \oint_U \frac{1}{2} \nabla u \cdot \mathbf{s} \nabla u, \quad \forall u \in \mathcal{A}(U). \quad (2.39)$$

In more generality, we have that

$$\frac{1}{2} (X)_U \cdot \mathbf{A}_*(U) (X)_U \leq \frac{1}{2} \oint_U X \cdot \mathbf{A} X \quad \forall X \in \mathcal{S}(U). \quad (2.40)$$

The proof of (2.40) is simple: we compute

$$\begin{aligned} \frac{1}{2} (X)_U \cdot \mathbf{A}_*(U) (X)_U &= \sup_{Q \in \mathbb{R}^{2d}} (Q \cdot (X)_U - \frac{1}{2} Q \cdot \mathbf{A}_*^{-1}(U) Q) \\ &= \sup_{Q \in \mathbb{R}^{2d}} (Q \cdot (X)_U - \mathbf{J}(U, 0, Q)) \\ &= \sup_{Q \in \mathbb{R}^{2d}} \inf_{Z \in \mathcal{S}(U)} \left( \oint_U (Q \cdot (X - Z)_U + \frac{1}{2} Z \cdot \mathbf{A} Z) \right) \leq \oint_U \frac{1}{2} X \cdot \mathbf{A} X. \end{aligned}$$

We refer to inequalities like (2.35), (2.38), (2.39) and (2.40) as *coarse-graining inequalities*. They give strong evidence that the coarse-grained matrices are aptly named, and they play a central role in the arguments in this paper.

**2.3. Centering the anti-symmetric part of the coefficient field.** The set of solutions of the equation

$$-\nabla \cdot \mathbf{a} \nabla u = 0$$

does not change when we add a constant anti-symmetric matrix to the coefficient field  $\mathbf{a}(\cdot)$ . We may even consider that the field  $\mathbf{a}(\cdot)$  and its anti-symmetric part are defined only modulo a constant anti-symmetric matrix. This is an important invariance which is reflected in properties of the coarse-grained matrices.

For convenience, we extend the definition of the quantity  $J$  by defining, for each given constant anti-symmetric matrix  $\mathbf{h}_0 \in \mathbb{R}_{\text{skew}}^{d \times d}$ ,

$$J_{\mathbf{h}_0}(U, p, q) := \sup_{u \in \mathcal{A}(U)} \int_U \left( -\frac{1}{2} \nabla u \cdot \mathbf{s} \nabla u - p \cdot (\mathbf{a} - \mathbf{h}_0) \nabla u + q \cdot \nabla u \right).$$

In other words, the quantity  $J_{\mathbf{h}_0}$  this is the same as  $J$  if we replace the coefficient field  $\mathbf{a}(\cdot)$  with  $\mathbf{a} - \mathbf{h}_0$ . The observation is then that the  $J$ 's for different  $\mathbf{h}_0$ 's are equivalent in the sense that

$$J_{\mathbf{h}_0}(U, p, q) = J(U, p, q - \mathbf{h}_0 p). \quad (2.41)$$

Indeed, the solution space  $\mathcal{A}(U)$  is unchanged by the subtraction of  $\mathbf{h}_0$  and therefore, if we let  $v_{\mathbf{h}_0}(\cdot, U, p, q) \in \mathcal{A}(U)$  denote the maximizer of  $J_{\mathbf{h}_0}(U, p, q)$ , it follows immediately from the definitions that

$$\nabla v_{\mathbf{h}_0}(\cdot, U, p, q) = \nabla v(\cdot, U, p, q - \mathbf{h}_0 p), \quad \forall p, q \in \mathbb{R}^d. \quad (2.42)$$

From this and a routine computation, we obtain (2.41). The quantity  $J_{\mathbf{h}_0}(U, \cdot, \cdot)$  can be represented by a matrix  $\mathbf{A}_{\mathbf{h}_0}(U)$  which is computed in terms of  $\mathbf{A}(U)$  as follows:

$$\begin{aligned} \begin{pmatrix} -p \\ q \end{pmatrix} \cdot \mathbf{A}_{\mathbf{h}_0}(U) \begin{pmatrix} -p \\ q \end{pmatrix} &= 2J_0(U, p, q - \mathbf{h}_0 p) + 2p \cdot q \\ &= \begin{pmatrix} -p \\ q - \mathbf{h}_0 p \end{pmatrix} \cdot \mathbf{A}(U) \begin{pmatrix} -p \\ q - \mathbf{h}_0 p \end{pmatrix} + 2p \cdot \mathbf{h}_0 p \\ &= \begin{pmatrix} -p \\ q \end{pmatrix} \cdot \begin{pmatrix} \mathbf{I}_d & 0 \\ \mathbf{h}_0 & \mathbf{I}_d \end{pmatrix}^t \mathbf{A}(U) \begin{pmatrix} \mathbf{I}_d & 0 \\ \mathbf{h}_0 & \mathbf{I}_d \end{pmatrix} \begin{pmatrix} -p \\ q \end{pmatrix}. \end{aligned} \quad (2.43)$$

Here we used the fact that  $p \cdot \mathbf{h}_0 p = 0$  since  $\mathbf{h}_0$  is anti-symmetric. Therefore we deduce that

$$\mathbf{A}_{\mathbf{h}_0}(U) = \mathbf{G}_{\mathbf{h}_0}^t \mathbf{A}(U) \mathbf{G}_{\mathbf{h}_0} = \begin{pmatrix} \mathbf{s} + (\mathbf{k} - \mathbf{h}_0)^t \mathbf{s}_*^{-1} (\mathbf{k} - \mathbf{h}_0) & -(\mathbf{k} - \mathbf{h}_0)^t \mathbf{s}_*^{-1} \\ -\mathbf{s}_*^{-1} (\mathbf{k} - \mathbf{h}_0) & \mathbf{s}_*^{-1} \end{pmatrix} (U), \quad (2.44)$$

where we denote

$$\mathbf{G}_{\mathbf{h}_0} := \begin{pmatrix} \mathbf{I}_d & 0 \\ \mathbf{h}_0 & \mathbf{I}_d \end{pmatrix}, \quad (2.45)$$

Comparing (2.44) with (2.1), we see that the subtraction of a constant anti-symmetric matrix  $\mathbf{h}_0$  “commutes” with the coarse-graining operation in the sense that it leaves  $\mathbf{s}(U)$  and  $\mathbf{s}_*(U)$  unchanged and simply subtracts  $\mathbf{h}_0$  from  $\mathbf{k}(U)$ . In particular, for an anti-symmetric matrix  $\mathbf{h}_0$ , we have

$$J(U, p, q - \mathbf{h}_0 p) = \frac{1}{2} \begin{pmatrix} -p \\ q \end{pmatrix} \cdot \mathbf{A}_{\mathbf{h}_0}(U) \begin{pmatrix} -p \\ q \end{pmatrix} - p \cdot q. \quad (2.46)$$

We will use this in our analysis to “center” the quantity  $J$ . We define  $\mathbf{A}_{\mathbf{h}_0,*}$  analogously and we also denote  $\mathbf{b}_{\mathbf{h}_0}(U) := (\mathbf{s} + (\mathbf{k} - \mathbf{h}_0)^t \mathbf{s}_*^{-1}(\mathbf{k} - \mathbf{h}_0))(U)$ .

All of the properties of the coarse-grained matrices given in the previous section can of course be applied to the field  $\mathbf{a} - \mathbf{h}_0$  and then written in terms of  $\mathbf{A}_{\mathbf{h}_0}$ ,  $\mathbf{b}_{\mathbf{h}_0}$ , and so forth. For instance, (2.39) implies that, for any constant anti-symmetric matrix  $\mathbf{h}_0 \in \mathbb{R}_{\text{skew}}^{d \times d}$ ,

$$\frac{1}{2} \left( \int_U (\mathbf{a} - \mathbf{h}_0) \nabla u \right) \cdot \mathbf{b}_{\mathbf{h}_0}^{-1}(U) \left( \int_U (\mathbf{a} - \mathbf{h}_0) \nabla u \right) \leq \int_U \frac{1}{2} \nabla u \cdot \mathbf{s} \nabla u, \quad \forall u \in \mathcal{A}(U). \quad (2.47)$$

In view of (2.41) and (2.42), we have

$$\begin{aligned} \int_U \left( \begin{array}{c} \nabla v \\ (\mathbf{a} - \mathbf{h}_0) \nabla v \end{array} \right) (\cdot, U, p, q - \mathbf{h}_0 p) &= \begin{pmatrix} 0 & \mathbf{I}_d \\ \mathbf{I}_d & 0 \end{pmatrix} \mathbf{A}_{\mathbf{h}_0}(U) \begin{pmatrix} -p \\ q \end{pmatrix} + \begin{pmatrix} -p \\ q \end{pmatrix} \\ &= \begin{pmatrix} -p + \mathbf{s}_*^{-1}(U)(q + (\mathbf{k} - \mathbf{h}_0)(U)p) \\ (\mathbf{I}_d - (\mathbf{k} - \mathbf{h}_0)^t \mathbf{s}_*^{-1})(U)q - \mathbf{b}_{\mathbf{h}_0}(U)p \end{pmatrix}. \end{aligned} \quad (2.48)$$

Throughout the paper, we will freely use the identities and inequalities in the previous subsection, after shifting by any anti-symmetric matrix  $\mathbf{h}_0$  of our choosing.

Since the “centering” operation maps  $\mathbf{A}(U)$  to  $\mathbf{G}_{\mathbf{h}_0}^t \mathbf{A}(U) \mathbf{G}_{\mathbf{h}_0}$ , it leaves the eigenvalues of ratios of pairs of coarse-grained matrices unchanged. Indeed, for any matrix  $\mathbf{h}_0 \in \mathbb{R}^{d \times d}$  (not necessarily skew symmetric) and pair of symmetric matrices  $\mathbf{D}, \mathbf{E} \in \mathbb{R}^{2d \times 2d}$  with  $\mathbf{D}$  being positive definite, if we denote

$$\mathbf{D}_{\mathbf{h}_0} := \mathbf{G}_{\mathbf{h}_0}^t \mathbf{D} \mathbf{G}_{\mathbf{h}_0} \quad \text{and} \quad \mathbf{E}_{\mathbf{h}_0} := \mathbf{G}_{\mathbf{h}_0}^t \mathbf{E} \mathbf{G}_{\mathbf{h}_0},$$

then

$$\mathbf{D}_{\mathbf{h}_0}^{-1} \mathbf{E}_{\mathbf{h}_0} = \mathbf{G}_{\mathbf{h}_0}^{-1} \mathbf{D}^{-1} \mathbf{E} \mathbf{G}_{\mathbf{h}_0}.$$

The matrix  $\mathbf{G}_{\mathbf{h}_0}$  is invertible with the inverse  $\mathbf{G}_{\mathbf{h}_0}^{-1} = \mathbf{G}_{-\mathbf{h}_0}$ . Thus  $\mathbf{D}_{\mathbf{h}_0}^{-1} \mathbf{E}_{\mathbf{h}_0}$  and  $\mathbf{D}^{-1} \mathbf{E}$  are similar. It follows that  $\mathbf{D}^{-1/2} \mathbf{E} \mathbf{D}^{-1/2}$  and  $\mathbf{D}_{\mathbf{h}_0}^{-1/2} \mathbf{E}_{\mathbf{h}_0} \mathbf{D}_{\mathbf{h}_0}^{-1/2}$  have the same set of eigenvalues. In particular,

$$|\mathbf{D}^{-1/2} \mathbf{E} \mathbf{D}^{-1/2} - \mathbf{I}_{2d}| = |\mathbf{D}_{\mathbf{h}_0}^{-1/2} \mathbf{E}_{\mathbf{h}_0} \mathbf{D}_{\mathbf{h}_0}^{-1/2} - \mathbf{I}_{2d}|. \quad (2.49)$$

Suppose next that  $\mathbf{E}_1, \mathbf{E}_2 \in \mathbb{R}^{2d \times 2d}$  are such that, for some  $\theta \in [1, \infty)$ ,

$$\mathbf{E}_1 \leq \theta \mathbf{E}_2 \quad \text{and} \quad \mathbf{E}_j := \mathbf{G}_{-\mathbf{k}_j}^t \begin{pmatrix} \mathbf{s}_j & 0 \\ 0 & \mathbf{s}_{*,j}^{-1} \end{pmatrix} \mathbf{G}_{-\mathbf{k}_j}, \quad \mathbf{s}_j, \mathbf{s}_{*,j}^{-1} \in \mathbb{R}_{\text{sym}}^{d \times d}, \quad \mathbf{k}_j \in \mathbb{R}^{d \times d}, \quad j \in \{1, 2\}. \quad (2.50)$$

Then we have

$$\theta \mathbf{s}_2 \geq \mathbf{s}_1 + (\mathbf{k}_1 - \mathbf{k}_2)^t \mathbf{s}_{*,1}^{-1} (\mathbf{k}_1 - \mathbf{k}_2) \quad \text{and} \quad \mathbf{s}_{*,2} \leq \theta \mathbf{s}_{*,1}. \quad (2.51)$$

To see this, we observe that, for every  $\mathbf{h} \in \mathbb{R}^{d \times d}$ ,

$$0 \leq \mathbf{G}_{\mathbf{h}}^t (\theta \mathbf{E}_2 - \mathbf{E}_1) \mathbf{G}_{\mathbf{h}} = \begin{pmatrix} \theta \mathbf{b}_{2,\mathbf{h}} - \mathbf{b}_{1,\mathbf{h}} & (\mathbf{k}_1 - \mathbf{h})^t \mathbf{s}_{*,1}^{-1} - \theta (\mathbf{k}_2 - \mathbf{h})^t \mathbf{s}_{*,2}^{-1} \\ \mathbf{s}_{*,1}^{-1} (\mathbf{k}_1 - \mathbf{h}) - \theta \mathbf{s}_{*,2}^{-1} (\mathbf{k}_2 - \mathbf{h}) & \theta \mathbf{s}_{*,2}^{-1} - \mathbf{s}_{*,1}^{-1} \end{pmatrix}, \quad (2.52)$$

and thus

$$\mathbf{b}_{1,\mathbf{h}} \leq \theta \mathbf{b}_{2,\mathbf{h}} \quad \forall \mathbf{h} \in \mathbb{R}^{d \times d} \quad \text{and} \quad \mathbf{s}_{*,2} \leq \theta \mathbf{s}_{*,1}. \quad (2.53)$$

Taking  $\mathbf{h} = \mathbf{k}_2$  yields (2.51). In particular, since  $\overline{\mathbf{A}}(\square_m) \leq \overline{\mathbf{A}}(\square_n)$  for every  $m, n \in \mathbb{N}$  with  $m \geq n$  and they are both of the above form, we have that

$$\overline{\mathbf{s}}(\square_m) \leq \overline{\mathbf{s}}(\square_n) \quad \text{and} \quad \overline{\mathbf{s}}_*(\square_m) \geq \overline{\mathbf{s}}_*(\square_n). \quad (2.54)$$

Notice also that, by (1.23), the matrix  $\mathbf{E}_0$  can be written in the form (2.50).

In view of the above discussion, for any fixed  $\mathbf{h}_0 \in \mathbb{R}^{d \times d}$ , the assumption (1.17) is equivalent to

$$3^m \geq \mathcal{S} \implies \mathbf{A}_{\mathbf{h}_0}(z + \square_n) \leq 3^{\gamma(m-n)} \mathbf{E}_{0, \mathbf{h}_0}, \quad \forall n \in \mathbb{Z} \cap (-\infty, m], \quad z \in 3^n \mathbb{Z}^d \cap \square_m, \quad (2.55)$$

where  $\mathbf{E}_{0, \mathbf{h}_0} := \mathbf{G}_{\mathbf{h}_0}^t \mathbf{E}_0 \mathbf{G}_{\mathbf{h}_0}$ . Since the transformation  $\mathbf{E}_0 \mapsto \mathbf{E}_{0, \mathbf{h}_0}$  leaves  $\mathbf{s}_0$  and  $\mathbf{s}_{*,0}$  unchanged, the ellipticity contrast  $\Theta$  is invariant under this transformation, while the new value of  $\Pi$  is bounded above by

$$|\mathbf{s}_0 + (\mathbf{h}_0 - \mathbf{h}_0)^t \mathbf{s}_{*,0}^{-1} (\mathbf{h}_0 - \mathbf{h}_0)| |\mathbf{s}_{*,0}^{-1}| \leq 2\Pi + 2|\mathbf{h}_0^t \mathbf{s}_{*,0}^{-1} \mathbf{h}_0| |\mathbf{s}_{*,0}^{-1}|.$$

**2.4. Two-sided bounds from one-sided bounds.** In the next two lemmas, we formalize an important observation, which is that if  $\Theta - 1$  is small and  $\mathbf{A}(U) \leq \mathbf{E}_0$ , then in fact the difference  $\mathbf{E}_0 - \mathbf{A}(U)$  must also be small. That is, we get a two-sided bound from a one-sided bound for free if the ellipticity contrast is small. This is related to the idea that the difference (or ratio) of  $\mathbf{s}(U)$  and  $\mathbf{s}_*(U)$  should represent the “uncertainty” in the coarse-graining map. Since this is an essentially algebraic fact, we present a slightly more general statement which will prove to be useful.

**Lemma 2.1.** *Suppose that  $\mathbf{E}_1, \mathbf{E}_{*,1} \in \mathbb{R}_{\text{sym}}^{2d \times 2d}$  are symmetric matrices having the form*

$$\mathbf{E}_1 = \begin{pmatrix} \mathbf{s}_1 + \mathbf{k}_1^t \mathbf{s}_{*,1}^{-1} \mathbf{k}_1 & -\mathbf{k}_1^t \mathbf{s}_{*,1}^{-1} \\ -\mathbf{s}_{*,1}^{-1} \mathbf{k}_1 & \mathbf{s}_{*,1}^{-1} \end{pmatrix} \quad \text{and} \quad \mathbf{E}_{*,1} = \begin{pmatrix} \mathbf{s}_{*,1} + \mathbf{k}_1 \mathbf{s}_1^{-1} \mathbf{k}_1^t & \mathbf{k}_1 \mathbf{s}_1^{-1} \\ \mathbf{s}_1^{-1} \mathbf{k}_1^t & \mathbf{s}_1^{-1} \end{pmatrix}, \quad (2.56)$$

satisfying the inequality

$$\mathbf{E}_{*,1} \leq \mathbf{E}_1, \quad (2.57)$$

where  $\mathbf{s}_1, \mathbf{s}_{*,1}, \mathbf{k}_1 \in \mathbb{R}^{d \times d}$  with  $\mathbf{s}_1$  and  $\mathbf{s}_{*,1}$  positive definite. Then  $\mathbf{s}_{*,1} \leq \mathbf{s}_1$  and, by denoting

$$\tilde{\Theta}_1 := |\mathbf{s}_{*,1}^{-1/2} \mathbf{s}_1 \mathbf{s}_{*,1}^{-1/2}|,$$

we have the inequalities

$$|\mathbf{s}_{*,1}^{-1/2} (\mathbf{k}_1 + \mathbf{k}_1^t) \mathbf{s}_{*,1}^{-1/2}| \leq \tilde{\Theta}_1 - 1 \quad (2.58)$$

and

$$|\mathbf{E}_{*,1}^{-1/2} \mathbf{E}_1 \mathbf{E}_{*,1}^{-1/2} - \mathbf{I}_{2d}| \leq 3\tilde{\Theta}_1^{1/2} (\tilde{\Theta}_1 - 1). \quad (2.59)$$

*Proof.* The inequality  $\mathbf{s}_1 \geq \mathbf{s}_{*,1}$  is immediate from (2.57), since the the bottom right matrices in the block forms in (2.56) must be ordered. Consequently,

$$|\mathbf{s}_{*,1}^{-1/2} \mathbf{s}_1 \mathbf{s}_{*,1}^{-1/2} - \mathbf{I}_d| = \tilde{\Theta}_1 - 1 \quad \text{and} \quad |\mathbf{s}_{*,1}^{1/2} (\mathbf{s}_{*,1}^{-1} - \mathbf{s}_1^{-1}) \mathbf{s}_{*,1}^{1/2}| \leq \tilde{\Theta}_1 - 1. \quad (2.60)$$

Let  $\mathbf{h}_0 \in \mathbb{R}^{d \times d}$  and recall the definition of  $\mathbf{G}_{\mathbf{h}_0}$  from (2.45). Observe that

$$\tilde{\mathbf{E}}_1 = \tilde{\mathbf{E}}_{1, \mathbf{h}_0} := \mathbf{G}_{\mathbf{h}_0}^t \mathbf{E}_1 \mathbf{G}_{\mathbf{h}_0} = \begin{pmatrix} \mathbf{s}_1 + (\mathbf{k}_1 - \mathbf{h}_0)^t \mathbf{s}_{*,1}^{-1} (\mathbf{k}_1 - \mathbf{h}_0) & -(\mathbf{k}_1 - \mathbf{h}_0)^t \mathbf{s}_{*,1}^{-1} \\ -\mathbf{s}_{*,1}^{-1} (\mathbf{k}_1 - \mathbf{h}_0) & \mathbf{s}_{*,1}^{-1} \end{pmatrix}$$

and

$$\tilde{\mathbf{E}}_{*,1} = \tilde{\mathbf{E}}_{*,1, \mathbf{h}_0} := \mathbf{G}_{\mathbf{h}_0}^t \mathbf{E}_{*,1} \mathbf{G}_{\mathbf{h}_0} = \begin{pmatrix} \mathbf{s}_{*,1} + (\mathbf{k}_1 + \mathbf{h}_0) \mathbf{s}_1^{-1} (\mathbf{k}_1 + \mathbf{h}_0)^t & (\mathbf{k}_1 + \mathbf{h}_0) \mathbf{s}_1^{-1} \\ \mathbf{s}_1^{-1} (\mathbf{k}_1 + \mathbf{h}_0)^t & \mathbf{s}_1^{-1} \end{pmatrix}.$$

Then  $\tilde{\mathbf{E}}_1$  and  $\tilde{\mathbf{E}}_{*,1}$  are positive and (2.57) is equivalent to  $\tilde{\mathbf{E}}_{*,1} \leq \tilde{\mathbf{E}}_1$ . Moreover, as discussed after (2.45), the matrices  $\tilde{\mathbf{E}}_1 \tilde{\mathbf{E}}_{*,1}^{-1}$  and  $\mathbf{E}_1 \mathbf{E}_{*,1}^{-1}$  are similar. In particular, since both  $\tilde{\mathbf{E}}_1$  and  $\tilde{\mathbf{E}}_{*,1}$  are symmetric,

$$|\tilde{\mathbf{E}}_{*,1}^{-1/2} \tilde{\mathbf{E}}_1 \tilde{\mathbf{E}}_{*,1}^{-1/2} - \mathbf{I}_{2d}| = |\mathbf{E}_{*,1}^{-1/2} \mathbf{E}_1 \mathbf{E}_{*,1}^{-1/2} - \mathbf{I}_{2d}|. \quad (2.61)$$

We next make a reduction to the case that  $\mathbf{k}_1$  is symmetric. Let  $\mathbf{k}_{1,s}$  and  $\mathbf{k}_{1,a}$  denote, respectively, the symmetric and anti-symmetric parts of  $\mathbf{k}_1$  and take  $\mathbf{h}_0 = \mathbf{k}_{1,a}$  in the above definitions. We then find that

$$\tilde{\mathbf{E}}_1 = \begin{pmatrix} \mathbf{s}_1 + \mathbf{k}_{1,s} \mathbf{s}_{*,1}^{-1} \mathbf{k}_{1,s} & -\mathbf{k}_{1,s} \mathbf{s}_{*,1}^{-1} \\ -\mathbf{s}_{*,1}^{-1} \mathbf{k}_{1,s} & \mathbf{s}_{*,1}^{-1} \end{pmatrix} \quad \text{and} \quad \tilde{\mathbf{E}}_{*,1} = \begin{pmatrix} \mathbf{s}_{*,1} + \mathbf{k}_{1,s} \mathbf{s}_1^{-1} \mathbf{k}_{1,s} & \mathbf{k}_{1,s} \mathbf{s}_1^{-1} \\ \mathbf{s}_1^{-1} \mathbf{k}_{1,s} & \mathbf{s}_1^{-1} \end{pmatrix}.$$

These matrices satisfy the same assumptions as  $\mathbf{E}_1$  and  $\mathbf{E}_{*,1}$ , and the symmetric part of  $\mathbf{k}_1$  is unchanged, but the anti-symmetric part of  $\mathbf{k}_1$  has been removed. In view of (2.61), we assume without loss of generality that  $\mathbf{k}_1$  is symmetric; otherwise we replace the pair  $(\mathbf{E}_1, \mathbf{E}_{*,1})$  by  $(\tilde{\mathbf{E}}_1, \tilde{\mathbf{E}}_{*,1})$ .

We next take  $\mathbf{h}_0 = \eta \mathbf{k}_1$  for  $\eta > 0$  in the definition of  $\mathbf{G}_{\mathbf{h}_0}$  and eventually optimize over the parameter  $\eta$ . The inequality  $\tilde{\mathbf{E}}_{*,1} \leq \tilde{\mathbf{E}}_1$  reads as

$$\begin{pmatrix} \mathbf{s}_1 - \mathbf{s}_{*,1} + (1 - \eta)^2 \mathbf{k}_1 \mathbf{s}_{*,1}^{-1} \mathbf{k}_1 - (1 + \eta)^2 \mathbf{k}_1 \mathbf{s}_1^{-1} \mathbf{k}_1 & -(1 - \eta) \mathbf{k}_1 \mathbf{s}_{*,1}^{-1} + (1 + \eta) \mathbf{k}_1 \mathbf{s}_1^{-1} \\ -(1 - \eta) \mathbf{s}_{*,1}^{-1} \mathbf{k}_1 + (1 + \eta) \mathbf{s}_1^{-1} \mathbf{k}_1 & \mathbf{s}_{*,1}^{-1} - \mathbf{s}_1^{-1} \end{pmatrix} \geq 0.$$

The nonnegativity of the top left block says that

$$(1 + \eta)^2 \mathbf{k}_1 \mathbf{s}_1^{-1} \mathbf{k}_1 \leq \mathbf{s}_1 - \mathbf{s}_{*,1} + (1 - \eta)^2 \mathbf{k}_1 \mathbf{s}_{*,1}^{-1} \mathbf{k}_1 \leq (\tilde{\Theta}_1 - 1) \mathbf{s}_{1,*} + \tilde{\Theta}_1 (1 - \eta)^2 \mathbf{k}_1 \mathbf{s}_1^{-1} \mathbf{k}_1.$$

Rearranging, we obtain

$$((1 + \eta)^2 - \tilde{\Theta}_1 (1 - \eta)^2) \mathbf{k}_1 \mathbf{s}_1^{-1} \mathbf{k}_1 \leq (\tilde{\Theta}_1 - 1) \mathbf{s}_{*,1}.$$

We now optimize in  $\eta$  by taking  $\eta := (\tilde{\Theta}_1 + 1)(\tilde{\Theta}_1 - 1)^{-1}$  to get

$$\mathbf{s}_{*,1}^{-1/2} \mathbf{k}_1 \mathbf{s}_1^{-1} \mathbf{k}_1 \mathbf{s}_{*,1}^{-1/2} \leq \frac{(\tilde{\Theta}_1 - 1)^2}{4\tilde{\Theta}_1} \mathbf{I}_d. \quad (2.62)$$

This yields (2.58).

To obtain (2.59), we use the above factorization with  $\mathbf{h}_0 = -\mathbf{k}_1^t = -\mathbf{k}_1$ . This gives us

$$\tilde{\mathbf{E}}_{1,-\mathbf{k}_1} = \begin{pmatrix} \mathbf{s}_1 + 4\mathbf{k}_1 \mathbf{s}_1^{-1} \mathbf{k}_1 & -2\mathbf{k}_1 \mathbf{s}_{*,1}^{-1} \\ -2\mathbf{s}_{*,1}^{-1} \mathbf{k}_1 & \mathbf{s}_{*,1}^{-1} \end{pmatrix} \quad \text{and} \quad \tilde{\mathbf{E}}_{*,1,-\mathbf{k}_1} = \begin{pmatrix} \mathbf{s}_{*,1} & 0 \\ 0 & \mathbf{s}_1^{-1} \end{pmatrix},$$

and therefore, using (2.60) and (2.62),

$$\begin{aligned} & \left| \tilde{\mathbf{E}}_{*,1,-\mathbf{k}_1}^{-1/2} \tilde{\mathbf{E}}_{1,-\mathbf{k}_1} \tilde{\mathbf{E}}_{*,1,-\mathbf{k}_1}^{-1/2} - \mathbf{I}_{2d} \right| \\ &= \left| \begin{pmatrix} \mathbf{s}_{*,1}^{-1/2} \mathbf{s}_1^{-1/2} - \mathbf{I}_d + 4\mathbf{s}_{*,1}^{-1/2} \mathbf{k}_1 \mathbf{s}_1^{-1} \mathbf{k}_1 \mathbf{s}_{*,1}^{-1/2} & -2\mathbf{s}_{*,1}^{-1/2} \mathbf{k}_1 \mathbf{s}_{*,1}^{-1} \mathbf{s}_1^{1/2} \\ -2\mathbf{s}_1^{1/2} \mathbf{s}_{*,1}^{-1} \mathbf{k}_1 \mathbf{s}_{*,1}^{-1/2} & \mathbf{s}_1^{1/2} \mathbf{s}_{*,1}^{-1} \mathbf{s}_1^{1/2} - \mathbf{I}_d \end{pmatrix} \right| \\ &\leq \tilde{\Theta}_1 - 1 + 4|\mathbf{s}_{*,1}^{-1/2} \mathbf{k}_1 \mathbf{s}_1^{-1} \mathbf{k}_1 \mathbf{s}_{*,1}^{-1/2}| + 2|\mathbf{s}_{*,1}^{-1/2} \mathbf{k}_1 \mathbf{s}_{*,1}^{-1} \mathbf{s}_1^{1/2}| \\ &\leq (2 + \tilde{\Theta}_1^{1/2})(\tilde{\Theta}_1 - 1). \end{aligned}$$

In view of (2.61) this completes the proof of (2.59).  $\square$

**Lemma 2.2.** Let  $\mathbf{E}_1$  and  $\tilde{\Theta}_1$  satisfy the hypotheses of Lemma 2.1 and let  $U$  be a bounded Lipschitz domain such that

$$\mathbf{A}(U) \leq \mathbf{E}_1. \quad (2.63)$$

Then

$$|\mathbf{E}_1^{-1/2} \mathbf{A}(U) \mathbf{E}_1^{-1/2} - \mathbf{I}_{2d}| \leq (2 + \tilde{\Theta}_1^{1/2})(\tilde{\Theta}_1 - 1). \quad (2.64)$$

*Proof.* The hypothesis of (2.63) implies that

$$\mathbf{E}_{*,1} \leq \mathbf{A}_*(U) \leq \mathbf{A}(U) \leq \mathbf{E}_1.$$

Indeed, the first inequality above is equivalent to (2.63), since  $\mathbf{E}_{*,1}^{-1}$  and  $\mathbf{A}_*(U)^{-1}$  are obtained, respectively, by flipping the rows and columns in the block matrix representation of  $\mathbf{E}_1$  and  $\mathbf{A}(U)$ . The inequality (2.64) then follows from (2.59).  $\square$

**2.5. Stochastic bounds for the coarse-grained matrices.** We show first that the assumption of (P2) implies control across a range of mesoscopic scales.

**Lemma 2.3** (Improving ellipticity on large mesoscales). *Assume that  $\mathbb{P}$  satisfies (P1) and (P2). For every  $h \in \mathbb{N}$ , there exists a random scale  $\mathcal{S}_h$  satisfying*

$$\mathcal{S}_h \leq \mathcal{O}_{\Psi_S}(K_{\Psi_S}^{4(d+1)} 3^h) \quad (2.65)$$

such that, for every  $m \in \mathbb{Z}$ ,

$$3^m \geq \mathcal{S}_h \implies \mathbf{A}(z + \square_n) \leq 3^{\gamma(m-h-n)+} \mathbf{E}_0, \quad \forall n \in \mathbb{Z} \cap (-\infty, m], \quad z \in 3^n \mathbb{Z}^d \cap \square_m. \quad (2.66)$$

*Proof.* Fix  $h \in \mathbb{N}$ . For every  $m \in \mathbb{N}$  with  $m \geq h$ , we have that

$$\begin{aligned} & \mathbb{P} \left[ \sup_{n \in \mathbb{Z} \cap (-\infty, m]} 3^{-\gamma(m-h-n)+} \sup_{z \in 3^n \mathbb{Z}^d \cap \square_m} |\mathbf{E}_0^{-1/2} \mathbf{A}(z + \square_n) \mathbf{E}_0^{-1/2}| > 1 \right] \\ &= \mathbb{P} \left[ \sup_{n \in \mathbb{Z} \cap (-\infty, m-h]} 3^{-\gamma(m-h-n)} \sup_{z \in 3^n \mathbb{Z}^d \cap \square_m} |\mathbf{E}_0^{-1/2} \mathbf{A}(z + \square_n) \mathbf{E}_0^{-1/2}| > 1 \right] \\ &\leq \sum_{z' \in 3^{m-h} \mathbb{Z}^d \cap \square_m} \mathbb{P} \left[ \sup_{n \in \mathbb{Z} \cap (-\infty, m-h]} 3^{-\gamma(m-h-n)} \sup_{z \in z' + 3^n \mathbb{Z}^d \cap \square_{m-h}} |\mathbf{E}_0^{-1/2} \mathbf{A}(z + \square_n) \mathbf{E}_0^{-1/2}| > 1 \right] \\ &= 3^{d(m-h)} \mathbb{P} \left[ \sup_{n \in \mathbb{Z} \cap (-\infty, m-h]} 3^{-\gamma(m-h-n)} \sup_{z \in 3^n \mathbb{Z}^d \cap \square_{m-h}} |\mathbf{E}_0^{-1/2} \mathbf{A}(z + \square_n) \mathbf{E}_0^{-1/2}| > 1 \right] \\ &\leq 3^{d(m-h)} \mathbb{P}[\mathcal{S} > 3^{m-h}] \\ &\leq 3^{-(m-h)} (\Psi_S(K_{\Psi_S}^{-4(d+1)} 3^{m-h}))^{-1}. \end{aligned}$$

In the above display, subadditivity was used to get the first equality, a union bound gives the next line, then stationarity in the following line, and finally assumption (P2) and (C.2) in the last line. Define

$$\mathcal{S}_h := \left\{ 3^m : m \in \mathbb{Z}, \sup_{n \in \mathbb{Z} \cap (-\infty, m]} 3^{-\gamma(m-h-n)+} \sup_{z \in 3^n \mathbb{Z}^d \cap \square_m} |\mathbf{E}_0^{-1/2} \mathbf{A}(z + \square_n) \mathbf{E}_0^{-1/2}| > 1 \right\}.$$

By another union bound and using (1.15), we obtain, for every  $m \in \mathbb{N}$  with  $m \geq h$ ,

$$\mathbb{P}[\mathcal{S}_h \geq 3^m] \leq \sum_{n=m}^{\infty} 3^{-(n-h)} (\Psi_S(K_{\Psi_S}^{-4(d+1)} 3^{n-h}))^{-1} \leq \frac{1}{\Psi_S(K_{\Psi_S}^{-4(d+1)} 3^{m-h})}.$$

This completes the proof of the lemma.  $\square$



The assumption (P2) gives us control over all finite moments of the coarse-grained matrices.

**Lemma 2.4** (Upper bounds for coarse-grained matrices). *For every  $m \in \mathbb{N}$  and  $n \in \mathbb{Z}$  with  $n \leq m$ ,*

$$\sup_{z \in 3^n \mathbb{Z}^d \cap \square_m} |\mathbf{E}_0^{-1/2} \mathbf{A}(z + \square_n) \mathbf{E}_0^{-1/2}| \leq 3^{\gamma(m-n)} \left(1 + \mathcal{O}_{\Psi_S}(3^{\gamma-m})\right). \quad (2.67)$$

In particular, for every  $n \in \mathbb{N}$ ,

$$|\mathbf{E}_0^{-1/2} \mathbf{A}(\square_n) \mathbf{E}_0^{-1/2}| \leq 1 + \mathcal{O}_{\Psi_S}(3^{\gamma-n}). \quad (2.68)$$

*Proof.* The assumption (P2) implies that, for every  $m \in \mathbb{N}$  and  $n \in \mathbb{Z}$  with  $n \leq m$ ,

$$\begin{aligned} \sup_{z \in 3^n \mathbb{Z}^d \cap \square_m} |\mathbf{E}_0^{-1/2} \mathbf{A}(z + \square_n) \mathbf{E}_0^{-1/2}| &\leq 3^{\gamma(m-n)} + \left(\frac{3\mathcal{S}}{3^n}\right)^\gamma \mathbf{1}_{\{S > 3^m\}} \\ &\leq 3^{\gamma(m-n)} + 3^{\gamma(m-n+1)} \left(\frac{\mathcal{S}}{3^m}\right) \\ &\leq 3^{\gamma(m-n)} \left(1 + \mathcal{O}_{\Psi_S}(3^{\gamma-m})\right). \end{aligned}$$

This completes the proof.  $\square$

In view of (2.68) and (C.5), we have the boundedness of all finite moments of  $|\mathbf{A}(\square_n)|$ . In fact, by (2.67), this can be extended to  $|\mathbf{A}(U)|$  for any bounded Lipschitz domain  $U \subseteq \mathbb{R}^d$  by partitioning the domain into triadic cubes and using subadditivity and the fact that  $\gamma < 1$ . This kind of argument can be found in proof of Lemma 2.8, below, so we do not give it here.

According to Lemma 2.4, the assumption of (P2) implies that  $\mathbf{A}(U)$  have all finite moments bounded. We may therefore define

$$\overline{\mathbf{A}}(U) := \mathbb{E}[\mathbf{A}(U)]. \quad (2.69)$$

We let  $\overline{\mathbf{s}}(U)$ ,  $\overline{\mathbf{s}}_*(U)$ ,  $\overline{\mathbf{k}}(U)$  and  $\overline{\mathbf{b}}(U)$  denote deterministic matrices which satisfy:

$$\begin{cases} \overline{\mathbf{s}}_*(U) = \mathbb{E}[\mathbf{s}_*^{-1}(U)]^{-1}, \\ \overline{\mathbf{s}}_*(U) \overline{\mathbf{k}}(U) = \mathbb{E}[\mathbf{s}_*^{-1}(U) \mathbf{k}(U)], \\ \overline{\mathbf{b}}(U) := \overline{\mathbf{s}}(U) + \overline{\mathbf{k}}^t(U) \overline{\mathbf{s}}_*^{-1}(U) \overline{\mathbf{k}}(U) = \mathbb{E}[\mathbf{s}(U) + \mathbf{k}^t(U) \mathbf{s}_*^{-1}(U) \mathbf{k}(U)]. \end{cases} \quad (2.70)$$

We see immediately that the first line of (2.70) defines  $\overline{\mathbf{s}}_*(U)$ , the second line defines  $\overline{\mathbf{k}}(U)$ , and the third line defines  $\overline{\mathbf{s}}(U)$  and  $\overline{\mathbf{b}}(U)$ . In other words, these matrices are defined in such a way that

$$\overline{\mathbf{A}}(U) = \begin{pmatrix} (\overline{\mathbf{s}} + \overline{\mathbf{k}}^t \overline{\mathbf{s}}_*^{-1} \overline{\mathbf{k}})(U) & -(\overline{\mathbf{k}}^t \overline{\mathbf{s}}_*^{-1})(U) \\ -(\overline{\mathbf{s}}_*^{-1} \overline{\mathbf{k}})(U) & \overline{\mathbf{s}}_*^{-1}(U) \end{pmatrix} \quad (2.71)$$

and taking the expectation of most natural expressions involving the coarse-grained matrices amounts to putting bars over each matrix.

We next show that, for every  $n \in \mathbb{N}$ ,

$$(1 + 3^{3-n} K_{\Psi_S}^2)^{-1} \overline{\mathbf{A}}(\square_n) \leq \mathbf{E}_0 \leq (1 + 3^{6-n} K_{\Psi_S}^2) (1 + 32(\Theta - 1)) \overline{\mathbf{A}}(\square_n). \quad (2.72)$$

We compute

$$\begin{aligned} \mathbb{E}[\mathbf{E}_0^{-1/2} \mathbf{A}(\square_n) \mathbf{E}_0^{-1/2}] &= \mathbb{E}[\mathbf{E}_0^{-1/2} \mathbf{A}(\square_n) \mathbf{E}_0^{-1/2} \mathbf{1}_{\{3^n \geq S\}}] + \mathbb{E}[\mathbf{E}_0^{-1/2} \mathbf{A}(\square_n) \mathbf{E}_0^{-1/2} \mathbf{1}_{\{3^n < S\}}] \\ &\leq (1 + 3^\gamma \mathbb{E}[\mathcal{S}^\gamma \mathbf{1}_{\{3^n < S\}}] 3^{-\gamma n}) \mathbf{I}_{2d} \leq (1 + 3^{1-n} \mathbb{E}[S]) \mathbf{I}_{2d}. \end{aligned} \quad (2.73)$$

The expectation of  $\mathcal{S}$  can be crudely estimated using (C.5) as

$$\mathbb{E}[\mathcal{S}] \leq \left(1 + 2K_{\Psi_S}(1 + \log K_{\Psi_S})\right) \leq 5K_{\Psi_S}^2.$$

We therefore obtain the lower bound in (2.72). By taking  $\mathbf{h}_0$  be the minimizing matrix in (1.25), by (2.53), the above estimates also yield that

$$\bar{\mathbf{s}}_*^{-1}(\square_n) \leq (1 + 3^{1-n}\mathbb{E}[\mathcal{S}])\mathbf{s}_{*,0}^{-1} \quad \text{and} \quad \mathbf{b}_{\mathbf{h}_0}(\square_n) \leq (1 + 3^{1-n}\mathbb{E}[\mathcal{S}])\mathbf{b}_{0,\mathbf{h}_0}.$$

Therefore, using (2.23), (2.68) and  $\bar{\mathbf{s}}_*(\square_n) \leq \bar{\mathbf{s}}(\square_n)$ , we then also deduce by the above display that

$$\begin{aligned} |\bar{\mathbf{A}}(\square_n)^{-1/2} \mathbf{E}_0 \bar{\mathbf{A}}(\square_n)^{-1/2}| &\leq 4(|\bar{\mathbf{s}}^{-1}(\square_n)\mathbf{b}_{0,\mathbf{h}_0}| \vee |(\bar{\mathbf{s}}_* + (\bar{\mathbf{k}} - \mathbf{h}_0)\bar{\mathbf{s}}^{-1}(\bar{\mathbf{k}} - \mathbf{h}_0)^t)(\square_n)\mathbf{s}_{*,0}^{-1}|) \\ &\leq 4\Theta(1 + 3^{3-n}K_{\Psi_S}^2). \end{aligned} \quad (2.74)$$

This proves the upper bound in (2.72) if  $\Theta \geq 9/8$ . If, on the other hand,  $\Theta \leq 9/8$ , then the upper bound follows by Lemma 2.2

We next discuss sensitivity estimates for the random matrix  $\mathbf{A}(U)$ . It is immediate from the variational characterization of  $\mathbf{A}(U)$  in (2.12) and (2.16) that, with  $D_U$  defined in (1.28) above,

$$|D_U(P \cdot \mathbf{A}(U)P)| \leq P \cdot \mathbf{A}(U)P, \quad \forall P \in \mathbb{R}^{2d}. \quad (2.75)$$

Indeed, if we fix a bounded Lipschitz domain  $U \subseteq \mathbb{R}^d$  and let  $\mathbf{a}_1$  and  $\mathbf{a}_2$  be two coefficient fields in  $\Omega$  with corresponding  $2d$  matrices  $\mathbf{A}_1$  and  $\mathbf{A}_2$ , then we find that

$$\begin{aligned} P \cdot \mathbf{A}_1(U)P &= \inf \left\{ \int_U (X + P) \cdot \mathbf{A}_1(X + P) : X \in L_{\mathbf{a},\text{pot},0}^2(U) \times L_{\mathbf{a},\text{sol},0}^2(U) \right\} \\ &\leq \|\mathbf{A}_2^{-1/2} \mathbf{A}_1 \mathbf{A}_2^{-1/2}\|_{L^\infty(U)} \inf \left\{ \int_U (X + P) \cdot \mathbf{A}_2(X + P) : X \in L_{\mathbf{a},\text{pot},0}^2(U) \times L_{\mathbf{a},\text{sol},0}^2(U) \right\} \\ &= \|\mathbf{A}_2^{-1/2} \mathbf{A}_1 \mathbf{A}_2^{-1/2}\|_{L^\infty(U)} \cdot P \cdot \mathbf{A}_2(U)P \end{aligned}$$

This implies (2.75). It is immediate from the definitions that

$$\mathbf{A}(U) \text{ is } \mathcal{F}(U)\text{-measurable.} \quad (2.76)$$

The sensitivity estimate (2.75) and the locality (2.76) of  $\mathbf{A}(U)$  will allow us to apply our mixing assumption (P3) to sums of coarse-grained matrices.

We next apply the CFS condition (P3) to sums of the coarse-grained matrix  $\mathbf{A}(U)$ . Since these random variables are not bounded, we need to apply a cutoff function and use the previous lemma to control the error this causes.

**Lemma 2.5** (Concentration for sums of  $\mathbf{A}$ 's). *For every  $k, m, n \in \mathbb{N}$  with  $\beta m < n < k \leq m$  and  $z \in 3^k \mathbb{Z}^d \cap \square_m$ ,*

$$\sum_{z' \in 3^n \mathbb{Z}^d \cap (z + \square_k)} (\mathbf{A}(z' + \square_n) - \bar{\mathbf{A}}(\square_n)) \mathbf{1}_{\{\mathcal{S} \leq 3^m\}} \leq \mathcal{O}_\Psi \left( 4 \cdot 3^{\gamma(m-n)} 3^{-\nu(k-n)} \mathbf{E}_0 \right). \quad (2.77)$$

*Proof.* Denote  $T := 3^{\gamma(m-n)}$ . We take a smooth cutoff function  $\varphi : \mathbb{R}_+ \rightarrow [0, 1]$  satisfying

$$\mathbf{1}_{[0,T]} \leq \varphi \leq \mathbf{1}_{[0,2T]}, \quad |\varphi'| \leq 2T^{-1}. \quad (2.78)$$

Denote, for each  $z \in 3^n \mathbb{Z}^d \cap \square_m$ ,

$$\tilde{\mathbf{A}}_z := \varphi(|\mathbf{E}_0^{-1/2} \mathbf{A}(z + \square_n) \mathbf{E}_0^{-1/2}|) \mathbf{E}_0^{-1/2} \mathbf{A}(z + \square_n) \mathbf{E}_0^{-1/2}.$$

It is clear that  $|\tilde{\mathbf{A}}_z| \leq 2T$ . According to (2.75), we have that

$$|D_{z+\square_n} \tilde{\mathbf{A}}_z| \leq |\tilde{\mathbf{A}}_z| (1 + \varphi'(|\mathbf{E}_0^{-1/2} \mathbf{A}(z + \square_n) \mathbf{E}_0^{-1/2}|)) \leq 4T. \quad (2.79)$$

By (2.76), it is clear that  $\tilde{\mathbf{A}}_z$  is  $\mathcal{F}(z + \square_n)$ -measurable. We may therefore apply (P3) to obtain, for every  $k \in \mathbb{N} \cap (n, m]$  and  $z \in 3^k \mathbb{Z}^d \cap \square_m$ ,

$$\left| \sum_{z' \in 3^n \mathbb{Z}^d \cap (z + \square_k)} (\tilde{\mathbf{A}}_{z'} - \mathbb{E}[\tilde{\mathbf{A}}_{z'}]) \right| \leq \mathcal{O}_\Psi(4T 3^{-\nu(k-n)}). \quad (2.80)$$

We also have that

$$\mathbf{A}(z + \square_n) \mathbf{1}_{\{\mathcal{S} \leq 3^m\}} = \tilde{\mathbf{A}}_z \mathbf{1}_{\{\mathcal{S} \leq 3^m\}} \quad \text{and} \quad \overline{\mathbf{A}}(\square_n) \geq \mathbb{E}[\tilde{\mathbf{A}}_z], \quad (2.81)$$

and therefore

$$\sum_{z' \in 3^n \mathbb{Z}^d \cap (z + \square_k)} \mathbf{E}_0^{-1/2} (\mathbf{A}(z' + \square_n) - \overline{\mathbf{A}}(\square_n)) \mathbf{E}_0^{-1/2} \mathbf{1}_{\{\mathcal{S} \leq 3^m\}} \leq \sum_{z' \in 3^n \mathbb{Z}^d \cap (z + \square_k)} (\tilde{\mathbf{A}}_{z'} - \mathbb{E}[\tilde{\mathbf{A}}_{z'}]).$$

Therefore, (2.77) follows by (2.80). The proof is complete.  $\square$

**2.6. Renormalization of the ellipticity assumption.** We next show that the assumptions  $\mathbb{P}$  satisfies (P1), (P2) and (P3) can be *renormalized*. To formalize this, we introduce the mapping  $D_{n_0} : \Omega \rightarrow \Omega$  given by dilation by  $3^{n_0}$ ,

$$(D_{n_0} \mathbf{a})(x) = \mathbf{a}(3^{n_0} x) \quad (2.82)$$

and we define  $\mathbb{P}_{n_0}$  by

$$\mathbb{P}_{n_0} := \text{the pushforward of } \mathbb{P} \text{ under } D_{n_0}. \quad (2.83)$$

We show that  $\mathbb{P}_{n_0}$  satisfies the same assumptions as  $\mathbb{P}$ —with the ellipticity matrix  $\mathbf{E}_0$  replaced by  $\overline{\mathbf{A}}(\square_{n_0-l_0})$  for a sufficiently large scale separation parameter  $l_0$ —and some suitable modifications to the other parameters (we must also slightly enlarge  $\gamma$  and replace  $\mathcal{S}$  by a new minimal scale  $\mathcal{S}'$  which has integrability quantified by a new function  $\Psi_{\mathcal{S}'}$  given in terms of the  $\Psi_{\mathcal{S}}$  and  $\Psi$ ).

The main point is that the ellipticity ratio for  $\overline{\mathbf{A}}(\square_{n_0-l_0})$  may be much smaller than  $\mathbf{E}_0$ . It is natural therefore to define, for each  $n \in \mathbb{N}$ , a new parameter  $\Theta_n$ , which we call the *renormalized ellipticity ratio*  $\Theta_n \in [1, \infty)$  at scale  $3^n$ , which is the ellipticity ratio for  $\overline{\mathbf{A}}(\square_n)$ . In view of (1.25) and (2.1), we define it by

$$\Theta_n := \min_{\mathbf{h}_0 \in \mathbb{R}_{\text{skew}}^{d \times d}} |(\bar{\mathbf{s}}_*^{-1/2} \bar{\mathbf{b}}_{\mathbf{h}_0} \bar{\mathbf{s}}_*^{-1/2})(\square_n)|. \quad (2.84)$$

Note that  $n \mapsto \Theta_n$  is monotone decreasing. For convenience, we define an exponent  $\mu$ , used throughout the rest of the paper, by

$$\mu := (\nu - \gamma)(1 - \beta). \quad (2.85)$$

**Proposition 2.6** (Renormalization of the assumptions). *Suppose  $\mathbb{P}$  satisfies (P1), (P2) and (P3). Let  $\rho \in (\gamma, \min\{\nu, 1\})$  and  $\delta > 0$ . Suppose that  $l_0 \in \mathbb{N}$  satisfies*

$$l_0 \geq \frac{1}{\rho - \gamma} \left(1 + \frac{d}{\mu}\right) (5 + \log(\delta^{-1}\Theta)) + \frac{6}{\mu} (1 + \log K_\Psi). \quad (2.86)$$

*For every  $n_0 \in \mathbb{N}$  with  $n_0 \geq l_0 + \log K_\Psi$ , the pushforward  $\mathbb{P}_{n_0}$  of  $\mathbb{P}$  under the dilation map given in (2.82) satisfies the assumptions (P1), (P2) and (P3), where the parameters  $(\gamma, \Psi_S, \mathbf{E}_0)$  in assumption (P2) are replaced by  $(\rho, \Psi_{S'}, (1 + \delta)\overline{\mathbf{A}}(\square_{n_0 - l_0}))$  and  $\Psi_{S'}$  is defined by*

$$\Psi_{S'}(t) := \frac{1}{2} \min\{\Psi_S(3^{n_0}t), \Psi(t^\mu)\}. \quad (2.87)$$

*Proof.* The conditions (P1) and (P3) for  $\mathbb{P}_{n_0}$  are immediate from their validity for  $\mathbb{P}$ , as the dilation causes no harm; the only condition which needs to be checked is therefore (P2), and this is the content of Lemma 2.7, below.  $\square$

The function  $\Psi_{S'}$  satisfies  $t\Psi_{S'}(t) \leq \Psi_{S'}(K_{\Psi_{S'}}t)$  for all  $t \geq 1$  with  $K_{\Psi_{S'}}$  given by

$$K_{\Psi_{S'}} := \max\{K_{\Psi_S}, K_\Psi^{\lceil 1/\mu \rceil}\}. \quad (2.88)$$

This follows from the definition of  $\Psi_{S'}$  in (2.87) and (C.8). The new value of the ellipticity ratios  $\Theta$  and  $\Pi$  are at most  $(1 + \delta)^2\Theta_{n_0 - l_0}$  and  $256(1 + \delta)^2\Pi$ , respectively. This follows immediately from the definition (2.84) of  $\Theta_n$ , and, respectively, (2.72) and the bound  $n_0 - l_0 \geq \log K_\Psi$ .

We turn to the proof of the main part of Proposition 2.6, which we put in the following lemma.

**Lemma 2.7** (Renormalizing of ellipticity). *Let  $\rho \in (\gamma, 1)$  and  $\delta > 0$ . Suppose that  $l_0 \in \mathbb{N}$  satisfies*

$$l_0 \geq \frac{1}{\rho - \gamma} \left(1 + \frac{d}{\mu}\right) (5 + \log(\delta^{-1}\Theta)) + \frac{6}{\mu} (1 + \log K_\Psi).$$

*For every  $n \in \mathbb{N}$  with  $n \geq l_0 + \log K_\Psi$ , there exists a minimal scale  $S' \geq S$  satisfying*

$$S' = \mathcal{O}_{\Psi_{S'}}(3^n) \quad \text{where} \quad \Psi_{S'}(t) := \frac{1}{2} \min\{\Psi_S(3^n t), \Psi(t^\mu)\}, \quad (2.89)$$

*such that, for every  $m \in \mathbb{N}$  with  $m \geq n$ ,*

$$3^m \geq S' \implies \mathbf{A}(z + \square_k) \leq (1 + \delta 3^{\rho(m-k)}) \overline{\mathbf{A}}(\square_{n-l_0}), \quad \forall k \in \mathbb{Z} \cap (-\infty, m], \quad z \in 3^k \mathbb{Z}^d \cap \square_m. \quad (2.90)$$

*Proof.* Let  $h \in \mathbb{N}$  be a parameter to be selected below. Let  $m, n, l_0 \in \mathbb{N}$  with  $m \geq n + h$  and  $n - l_0 \geq \log K_{\Psi_S}$ . Fix  $k, l \in \mathbb{N}$  with  $m - h < k \leq m$  and  $\max\{n - l_0, \beta k\} < l < k$ . Using (2.72), (2.77) and subadditivity, we find that, for every  $z \in 3^k \mathbb{Z}^d \cap \square_m$ ,

$$\begin{aligned} \mathbf{A}(z + \square_k) \mathbf{1}_{\{S \leq 3^m\}} &\leq \sum_{y \in 3^l \mathbb{Z}^d \cap (z + \square_k)} \mathbf{A}(y + \square_l) \mathbf{1}_{\{S \leq 3^m\}} \\ &\leq \overline{\mathbf{A}}(\square_l) + \mathcal{O}_\Psi(4 \cdot 3^{\gamma(m-l)} 3^{-\nu(k-l)} \mathbf{E}_0) \\ &\leq (1 + \mathcal{O}_\Psi(128\Theta(1 + 3^{-l}K_{\Psi_S}) \cdot 3^{\gamma(m-l)} 3^{-\nu(k-l)})) \overline{\mathbf{A}}(\square_l) \\ &\leq (1 + \mathcal{O}_\Psi(256\Theta 3^{\gamma(m-l)} 3^{-\nu(k-l)})) \overline{\mathbf{A}}(\square_l), \end{aligned} \quad (2.91)$$

where in the last line we used that  $3^{-l}K_{\Psi_S} \leq 1$  since  $l \geq n - l_0 \geq \log K_{\Psi_S}/\log 3$  by assumption. By a union bound, we deduce that, for every  $k \in \mathbb{N} \cap ((m - h) \vee l, m]$  and  $T \geq 1$ ,

$$\begin{aligned} & \mathbb{P} \left[ \sup_{z \in 3^k \mathbb{Z}^d \cap \square_m} \mathbf{A}(z + \square_k) \mathbf{1}_{\{\mathcal{S} \leq 3^m\}} \not\leq (1 + 256\Theta 3^{\gamma(m-l)} 3^{-\nu(k-l)} T) \overline{\mathbf{A}}(\square_l) \right] \\ & \leq \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \mathbb{P} \left[ \mathbf{A}(z + \square_k) \mathbf{1}_{\{\mathcal{S} \leq 3^m\}} \not\leq (1 + 256\Theta 3^{\gamma(m-l)} 3^{-\nu(k-l)} T) \overline{\mathbf{A}}(\square_l) \right] \leq \frac{3^{d(m-k)}}{\Psi(T)}. \end{aligned}$$

We set  $l := n - l_0 + \lceil \beta(k - n + l_0) \rceil$ . Observe that this choice of  $l$  satisfies  $\max\{n - l_0, \beta k\} < l < k$  announced above.

$$m - l \geq m - k + (1 - \beta)(k - n + l_0) - 1.$$

We also define

$$T := 2^{-10} \delta \Theta^{-1} 3^{\mu(k-n+l_0)} 3^{(\rho-\gamma)(m-k)},$$

and we observe that,

$$T \geq 2^{-10} \delta \Theta^{-1} 3^{\mu(l_0-h)} 3^{\mu(m-n)} \geq 1,$$

provided that

$$l_0 \geq h + \frac{7 + \log(\delta^{-1}\Theta)}{\mu}. \quad (2.92)$$

Note that this also implies that  $l < m - h$ . Substituting this choice of  $T$  into the inequalities above therefore yields that

$$\mathbb{P} \left[ \sup_{z \in 3^k \mathbb{Z}^d \cap \square_m} \mathbf{A}(z + \square_k) \mathbf{1}_{\{\mathcal{S} \leq 3^m\}} \not\leq (1 + \delta 3^{\rho(m-k)}) \overline{\mathbf{A}}(\square_l) \right] \leq \frac{3^{d(m-k)}}{\Psi(2^{-10} \delta \Theta^{-1} 3^{\mu(l_0-h)} 3^{\mu(m-n)})}.$$

Summing over  $k \in \{m - h + 1, \dots, m\}$  and using (1.29), (C.13) with  $p = 1$ , and a union bound, we obtain

$$\begin{aligned} & \mathbb{P} \left[ \sup_{k \in \mathbb{N} \cap [m-h+1, m]} \sup_{z \in 3^k \mathbb{Z}^d \cap \square_m} \mathbf{A}(z + \square_k) \mathbf{1}_{\{\mathcal{S} \leq 3^m\}} \not\leq (1 + \delta 3^{\rho(m-k)}) \overline{\mathbf{A}}(\square_{n-l_0}) \right] \\ & \leq \frac{3^{dh}}{\Psi(2^{-10} \delta \Theta^{-1} 3^{\mu(l_0-h)} 3^{\mu(m-n)})} \leq \frac{1}{\Psi(K_{\Psi}^{-6} 3^{-dh} 2^{-10} \delta \Theta^{-1} 3^{\mu(l_0-h)} 3^{\mu(m-n)})}. \end{aligned}$$

If we impose the another restriction on  $l_0$ , namely that

$$l_0 \geq \left(1 + \frac{d}{\mu}\right)h + \frac{6}{\mu}(1 + \log K_{\Psi}) + \frac{1}{\mu} \log(\delta^{-1}\Theta),$$

then  $K_{\Psi}^{-6} 3^{-dh} 2^{-10} \delta \Theta^{-1} 3^{\mu(l_0-h)} \geq 1$  and we can therefore upper bound the right side of the previous display by

$$\frac{1}{\Psi(3^{\mu(m-n)})}.$$

For the small scales, we proceed more crudely: by (2.72), we have that, for every  $k \leq m - h$ ,

$$3^m \geq \mathcal{S} \implies \sup_{k \in \mathbb{N} \cap [-\infty, m-h]} \sup_{z \in 3^k \mathbb{Z}^d \cap \square_m} \mathbf{A}(z + \square_k) \mathbf{1}_{\{\mathcal{S} \leq 3^m\}} \leq 3^{\gamma(m-k)} \mathbf{E}_0 \leq \delta 3^{\rho(m-k)} \overline{\mathbf{A}}(\square_{n-l_0}),$$

provided that we choose  $h > 0$  large enough so that

$$32\Theta(1 + 3^{-(n-l_0)}K_{\Psi_S}^2) \leq \delta 3^{(\rho-\gamma)h}.$$

Since  $(1 + 3^{-(n-l_0)}K_{\Psi_S}^2) \leq 2$  and  $\log 64 < 4 \log 3$ , it suffices to take any  $h \in \mathbb{N}$  satisfying

$$h \geq \frac{1}{\rho - \gamma} \left( 4 + \log(\delta^{-1}\Theta) \right).$$

Therefore, we may define a new minimal scale  $\mathcal{S}'$  by taking the maximum of  $\mathcal{S}$  and the following random scale:

$$\sup \left\{ 3^m : \sup_{k \in \mathbb{N} \cap [m-h, m]} \sup_{z \in 3^k \mathbb{Z}^d \cap \square_m} \mathbf{A}(z + \square_k) \mathbf{1}_{\{\mathcal{S} \leq 3^m\}} \not\leq (1 + \delta 3^{\rho(m-k)}) \overline{\mathbf{A}}(\square_{n-l_0}) \right\}.$$

We have shown that (2.90) holds, and

$$\mathbb{P}[\mathcal{S}' > 3^m] \leq \frac{1}{\Psi_{\mathcal{S}}(3^m)} + \frac{1}{\Psi(3^{\mu(m-n)})} \leq \frac{1}{\Psi_{\mathcal{S}'}(3^{m-n})},$$

where we define the new  $\Psi_{\mathcal{S}'}$  by

$$\Psi_{\mathcal{S}'}(t) := \frac{1}{2} \min \left\{ \Psi_{\mathcal{S}}(3^n t), \Psi(t^\mu) \right\}.$$

This completes the proof of the lemma.  $\square$

**2.7. Intrinsic geometry, adapted cubes, and consequences of subadditivity.** The next step is to change the geometry. We next introduce the (metric) geometric mean of the matrices  $\mathbf{b}_0$  and  $\mathbf{s}_{*,0}$ , which we denote by

$$\mathbf{m}_0 := \mathbf{E}_{11} \# \mathbf{E}_{22}^{-1} = \mathbf{b}_0 \# \mathbf{s}_{*,0} = (\mathbf{s}_0 + \mathbf{h}_0 \mathbf{s}_{*,0}^{-1} \mathbf{h}_0) \# \mathbf{s}_{*,0} \quad \text{and} \quad \mathbf{M}_0 := \begin{pmatrix} \mathbf{m}_0 & 0 \\ 0 & \mathbf{m}_0^{-1} \end{pmatrix}. \quad (2.93)$$

This notion of geometric mean of two positive matrices is given in Appendix B. We will work in cubes which are adapted to the matrix  $\mathbf{m}_0$ . We denote the smallest eigenvalue of  $\mathbf{m}_0$  by

$$\lambda_0 := |\mathbf{m}_0^{-1}|^{-1}.$$

We define another symmetric matrix  $\mathbf{q}_0$  by taking a large integer  $k_0 \in \mathbb{N}$  (to be selected below in terms of  $d$ ) and

$$(\mathbf{q}_0)_{ij} := 3^{-k_0} \left[ 3^{k_0} \lambda_0^{-1/2} (\mathbf{m}_0^{1/2})_{ij} \right]. \quad (2.94)$$

In other words, we take the matrix  $\lambda_0^{-1/2} \mathbf{m}_0^{1/2}$  and slightly alter each entry so that it belongs to the lattice  $3^{-k_0} \mathbb{Z}^d$ . Note that  $\mathbf{q}_0$  is symmetric, and it satisfies

$$|\mathbf{q}_0 - \lambda_0^{-1/2} \mathbf{m}_0^{1/2}| \leq C 3^{-k_0},$$

where  $C$  depends only on  $d$ . In view of the fact that

$$\mathbf{I}_d \leq \lambda_0^{-1/2} \mathbf{m}_0^{1/2} \leq \Pi^{1/2} \mathbf{I}_d,$$

we also have that

$$(1 - C 3^{-k_0}) \lambda_0^{-1/2} \mathbf{m}_0^{1/2} \leq \mathbf{q}_0 \leq (1 + C 3^{-k_0}) \lambda_0^{-1/2} \mathbf{m}_0^{1/2},$$

By making  $k_0$  sufficiently large, depending only on  $d$ , we obtain that

$$\frac{99}{100}\lambda_0^{-1/2}\mathbf{m}_0^{1/2} \leq \mathbf{q}_0 \leq \frac{101}{100}\lambda_0^{-1/2}\mathbf{m}_0^{1/2}. \quad (2.95)$$

We next introduce the cubes adapted to the geometry of  $\mathbf{s}_0$  by

$$\diamond_k := \mathbf{q}_0(\square_k) = \left\{x \in \mathbb{R}^d : \mathbf{q}_0^{-1}x \in \square_k\right\}. \quad (2.96)$$

These are hyperrectangles, but they may of course have edges which are not parallel to the coordinate axes. Note that

$$\mathbf{m}_0 \text{ is a scalar matrix} \implies \diamond_k = \square_k. \quad (2.97)$$

In view of the bounds

$$\lambda I_d \leq \mathbf{m}_0 \leq \Lambda I_d, \quad (2.98)$$

the eccentricity of these cubes (the ratio of largest to smallest side), is at most  $\frac{101}{99}\Pi^{1/2}$ , and in fact they satisfy

$$\frac{99}{100}\square_k \subseteq \diamond_k \subseteq \frac{101}{100}\Pi^{1/2}\square_k. \quad (2.99)$$

For convenience, we let  $\mathbb{L}_0$  denote the lattice

$$\mathbb{L}_0 := \mathbf{q}_0(\mathbb{Z}^d) = \{\mathbf{q}_0 z : z \in \mathbb{Z}^d\}.$$

Note that  $\{z + \diamond_n : z \in 3^n \mathbb{L}_0 \cap \diamond_m\}$  is a partition (up to a set of measure zero) of  $\diamond_m$ . By the construction of  $\mathbf{q}_0$ , it is clear that

$$\mathbb{L}_0 \subseteq 3^{-k_0} \mathbb{Z}^d. \quad (2.100)$$

This is very convenient, since it means that  $3^n \mathbb{L}_0 \subseteq \mathbb{Z}^d$  for all  $n$  larger than a universal constant depending only on  $d$ .

We need to work with the adapted rectangles  $\diamond_m$  to avoid artificial factors of  $\Pi$  from creeping into our estimates. To see why this is so, consider a constant-coefficient equation like  $-\nabla \cdot \mathbf{s}_0 \nabla u = 0$ . This equation can be considered to have an ellipticity ratio of one, because the coefficients are constant. We could perform a simple affine change of coordinates and transform this equation to the Laplace equation. However, if we do not perform this change of variables, and we start performing standard elliptic estimates (such as for instance the Caccioppoli inequality) in standard Euclidean balls or cubes, we will see powers of the ratio  $\Pi$  of the largest to the smallest eigenvalue of  $\mathbf{s}_0$  appear in our estimates. These factors would not appear if we were clever enough to have changed variables beforehand.

To avoid these extra factors of  $\Pi$ , we can either perform the affine change of variables and then work in Euclidean balls or cubes, or else work in the original coordinate system but use the  $\mathbf{s}_0$ -adapted balls or cubes. Neither choice is particularly pleasant, but we have taken the latter approach because the change of variables alters the  $\mathbb{Z}^d$ -stationarity assumption to stationarity with respect to the lattice  $\lambda_0^{1/2} \mathbf{s}_0^{-1/2} \mathbb{Z}^d$ , which has its own notational problems. This choice makes our arguments more straightforward to adapt to the discrete setting, for instance, in which the equation is not posed on the continuum  $\mathbb{R}^d$  but on the lattice  $\mathbb{Z}^d$ .

Using subadditivity and a Whitney-type decomposition of the adapted cubes into (normal) triadic cubes, we can reduce upper bounds on the coarse-grained matrices in adapted cubes to those of (2.67) and (2.68).



**Lemma 2.8** (Upper bounds for  $\mathbf{A}$  in adapted cubes). *Let  $\delta \in (0, 1]$ . There exists  $C(d) < \infty$  such that, if we define  $h' \in \mathbb{N}$  to be the smallest integer satisfying*

$$3^{h'} \geq \frac{C\Pi}{\delta(1-\gamma)}, \quad (2.101)$$

*then, with  $\mathcal{S}_{h+h'}$  being the random scale in the statement of Lemma 2.3, we have, for every  $m \in \mathbb{N}$ ,*

$$3^m \geq \mathcal{S}_{h+h'} \implies \mathbf{A}(y + \diamond_n) \leq (1+\delta)3^{\gamma(m-n-h)_+} \mathbf{E}_0, \quad \forall n \leq m, \quad y \in \mathbb{R}^d, \quad y + \diamond_n \subseteq \diamond_m. \quad (2.102)$$

*Proof.* Fix  $h' \in 2\mathbb{N}$  to satisfy (2.101), where we will make the constant  $C$  sufficiently large where needed, but depending only on  $d$ . Let  $m \in \mathbb{N}$  be such that  $3^m \geq \mathcal{S}_{h+h'}$ , where  $\mathcal{S}_{h+h'}$  is the minimal scale given by Lemma 2.3. Let  $l := \lceil \log_3(C(d)\Pi^{1/2}) \rceil$  with sufficiently large  $C(d)$  such that  $\diamond_m \subseteq \square_{m+l}$ . By taking larger constant  $C(d)$  in (2.101), we may assume that  $h' \geq 2l$ . Suppose that  $y \in \mathbb{R}^d$  and  $n \in \mathbb{Z}$  satisfy  $y + \diamond_n \subseteq \square_{m+l}$ . Then  $y + \diamond_n$  can be written as the disjoint union, up to a null set, of a family  $\{V_j(y) : -\infty < j \leq n\}$  of sets such that each  $V_j(y)$  is the disjoint union of cubes of the form  $z + \square_j$  with  $z \in 3^j\mathbb{Z}^d$ , and

$$|V_j(y)| \leq C\Pi^{1/2}3^{j-n}|\diamond_n| \quad \text{and} \quad \sum_{j=-\infty}^n \frac{|V_j(y)|}{|\diamond_n|} = 1. \quad (2.103)$$

We can obtain such a partition recursively as follows. Define first

$$V_n(y) := \bigcup \{z + \square_n : z + \square_n \subseteq y + \diamond_n\}$$

and then, having defined  $V_n(y), \dots, V_j(y)$ , we define  $V_{j-1}(y)$  by

$$V_{j-1}(y) := \bigcup \{z + \square_{j-1} : z \in 3^{j-1}\mathbb{Z}^d, z + \square_{j-1} \subseteq (y + \diamond_n) \setminus (V_n(y) \cup \dots \cup V_j(y))\}.$$

By subadditivity, the assumption  $3^{m+l} \geq \mathcal{S}_{h+h'}$  and (2.66), we obtain that

$$\begin{aligned} \mathbf{A}(y + \diamond_n) &\leq \sum_{j=-\infty}^n \frac{|V_j(y)|}{|\diamond_n|} \mathbf{A}(V_j(y)) \\ &\leq \sum_{j=-\infty}^n \frac{|V_j(y)|}{|\diamond_n|} 3^{\gamma(m+l-h'-h-j)_+} \mathbf{E}_0 \\ &\leq \left(1 + \sum_{j=-\infty}^{n+l-h'} (C\Pi^{1/2}3^{j-n}) 3^{\gamma(n+l-h'-j)_+}\right) 3^{\gamma(m-h-n)_+} \mathbf{E}_0 \\ &\leq \left(1 + \frac{C\Pi}{1-\gamma} 3^{-h'}\right) 3^{\gamma(m-h-n)_+} \mathbf{E}_0. \end{aligned}$$

The second last inequality is a consequence (2.103) and the triangle inequality:

$$(m+l-h'-h-j)_+ \leq (m-h-n)_+ + (n+l-h'-j)_+,$$

and the last inequality follows from the definition of  $l$ . If we now choose the constant  $C(d)$  in (2.101) large enough, (2.102) follows. This completes the proof.  $\square$

We next formalize a version of Lemma 2.5 in the adapted cubes.

**Lemma 2.9** (Concentration for adapted cubes). *There exists a constant  $C(d) < \infty$  such that, for every  $m, n \in \mathbb{N}$  with  $\beta m < n < m$ ,*

$$\left| \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathbf{E}_0^{-1/2} (\mathbf{A}(z + \diamond_n) - \overline{\mathbf{A}}(z + \diamond_n)) \mathbf{E}_0^{-1/2} \right| \leq \frac{CK_{\Psi_S}^2 \Pi}{1 - \gamma} 3^{\gamma(m-n)-m} + \mathcal{O}_{\Psi_S} \left( \frac{C\Pi}{1 - \gamma} 3^{\gamma(m-n)-m} \right) + \mathcal{O}_{\Psi} \left( C\Pi^{1/2} 3^{-(\nu-\gamma)(m-n)} \right). \quad (2.104)$$

*Proof.* Denote  $T := 3^{\gamma(m-n)}$  and let  $\mathcal{S}_{h'}$  be as in Lemma 2.3 where  $h' \in \mathbb{N}$  is chosen as small as possible such that (2.101). The proof is now similar to the one of Lemma 2.5. There is a slightly annoying complication caused by the fact that the assumption (P3) is formulated in terms of (normal) triadic cubes, whereas now we need to apply it to sums of the adapted cubes. It turns out that this difficulty can be handled quite crudely, while only dropping a factor of  $\Pi^{1/2}$ , by putting the adapted cubes into groups based on membership in slightly larger Euclidean cubes. Throughout, we fix  $m, n \in \mathbb{N}$  with  $\beta m < n \leq m$ .

We let  $k_0$  be the smallest positive integer such that  $\diamond_0 \subseteq \square_{n_0}$ ; in view of (2.99), we have that  $3^{n_0} \leq 3\Pi^{1/2}$ . For each  $x \in \mathbb{R}^d$ , let  $[x]$  denote the nearest point of the lattice  $3^{n+n_0}\mathbb{Z}^d$  to  $x$ , with the lexicographical ordering used as a tiebreaker if this point is not unique. We have then that

$$x + \diamond_n \subseteq [x] + \square_{n+n_0+1}, \quad \forall x \in \mathbb{R}^d.$$

Meanwhile, each  $z \in 3^{n+n_0}\mathbb{Z}^d$  satisfies  $z = [x]$  for at most  $3^{n_0+1}$  many distinct elements  $x$  belonging to the lattice  $3^n \mathbb{L}_0$ .

Select a smooth cutoff function  $\varphi : \mathbb{R}_+ \rightarrow [0, 1]$  satisfying

$$\mathbf{1}_{[0, T]} \leq \varphi \leq \mathbf{1}_{[0, 2T]}, \quad |\varphi'| \leq 2T^{-1}. \quad (2.105)$$

We define, for each  $z \in 3^{n+n_0}\mathbb{Z}^d \cap \square_{m+n_0+1}$ ,

$$\mathbf{X}_z := \sum_{x \in z + 3^n \mathbb{L}_0 \cap (z + \diamond_m)} \varphi(|\mathbf{E}_0^{-1/2} \mathbf{A}(x + \diamond_n) \mathbf{E}_0^{-1/2}|) \mathbf{E}_0^{-1/2} \mathbf{A}(x + \diamond_n) \mathbf{E}_0^{-1/2}.$$

Since there are at most  $3^{n_0+1} \leq 9\Pi^{1/2}$  many distinct elements in the sum, we have that

$$|\mathbf{X}_z| \leq 18\Pi^{1/2}T.$$

It is clear that  $\mathbf{X}_z$  is  $\mathcal{F}(z + \square_{n+n_0+1})$ -measurable and, similar to (2.79), we have that

$$|D_{z+\square_n} \mathbf{X}_z| \leq 36\Pi^{1/2}T. \quad (2.106)$$

We want to sum  $\mathbf{X}_z$  over  $z \in 3^{n+n_0}\mathbb{Z}^d \cap \square_{m+n_0+1}$ , but there is some overlap in the cubes  $z + \square_{n+n_0+1}$ . So we break the sum into  $3^d$  many different sums, each with  $z$ 's corresponding to disjoint cubes, and apply (P3) to each of these. The result is

$$\left| \sum_{z \in 3^{n+n_0}\mathbb{Z}^d \cap \square_{m+n_0+1}} (\mathbf{X}_z - \mathbb{E}[\mathbf{X}_z]) \right| \leq \mathcal{O}_{\Psi}(C_d \Pi^{1/2} T \cdot 3^{-\nu(m-n)}).$$

Since  $\mathbf{1}_{\{\mathcal{S}_{h'} \leq 3^m\}} \varphi(|\mathbf{E}_0^{-1/2} \mathbf{A}(z + \diamond_n) \mathbf{E}_0^{-1/2}|) = \mathbf{1}_{\{\mathcal{S}_{h'} \leq 3^m\}}$  for every  $z \in 3^n \mathbb{L}_0 \cap \diamond_m$ , we deduce that

$$\begin{aligned} & \mathbf{1}_{\{\mathcal{S}_{h'} \leq 3^m\}} \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathbf{E}_0^{-1/2} (\mathbf{A}(z + \diamond_n) - \overline{\mathbf{A}}(z + \diamond_n)) \mathbf{E}_0^{-1/2} \\ &= \mathbf{1}_{\{\mathcal{S}_{h'} \leq 3^m\}} \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} (\mathbf{X}_z - \mathbb{E}[\mathbf{X}_z]) + \mathbf{1}_{\{\mathcal{S}_{h'} \leq 3^m\}} \mathbb{E}[\mathbf{E}_0^{-1/2} \mathbf{A}(\diamond_n) \mathbf{E}_0^{-1/2} \mathbf{1}_{\{\mathcal{S}_{h'} > 3^m\}}]. \end{aligned}$$

It follows that

$$\begin{aligned} & \left| \mathbf{1}_{\{\mathcal{S}_{h'} \leq 3^m\}} \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathbf{E}_0^{-1/2} (\mathbf{A}(z + \diamond_n) - \overline{\mathbf{A}}(z + \diamond_n)) \mathbf{E}_0^{-1/2} \right| \\ & \leq |\mathbb{E}[\mathbf{E}_0^{-1/2} \mathbf{A}(\diamond_n) \mathbf{E}_0^{-1/2} \mathbf{1}_{\{\mathcal{S}_{h'} > 3^m\}}]| + \mathcal{O}_\Psi(C\Pi^{1/2} T 3^{-\nu(m-n)}). \end{aligned}$$

In view of (C.5), we apply (2.102) with  $\delta = 1$  and  $h = 0$  to get that

$$|\mathbb{E}[\mathbf{E}_0^{-1/2} \mathbf{A}(\diamond_n) \mathbf{E}_0^{-1/2} \mathbf{1}_{\{\mathcal{S}_{h'} > 3^m\}}]| \leq 6 \cdot 3^{-n\gamma} \mathbb{E}[\mathcal{S}_{h'}^\gamma \mathbf{1}_{\{\mathcal{S}_{h'} > 3^m\}}] \leq \frac{CK_\Psi^2 \Pi}{1 - \gamma} 3^{\gamma(m-n)-m}.$$

Similarly,

$$\sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} |\mathbf{E}_0^{-1/2} \mathbf{A}(z + \diamond_n) \mathbf{E}_0^{-1/2}| \mathbf{1}_{\{\mathcal{S}_{h'} > 3^m\}} \leq 6 \cdot 3^{-n\gamma} \mathcal{S}_{h'}^\gamma \mathbf{1}_{\{\mathcal{S}_{h'} > 3^m\}} \leq \mathcal{O}_{\Psi_S} \left( \frac{C\Pi}{1 - \gamma} 3^{\gamma(m-n)-m} \right).$$

Combining the last three displays implies (2.104) and completes the proof.  $\square$

In the next lemma, we use subadditivity arguments similar to the one Lemma 2.8 to compare the means of the coarse-grained matrices in Euclidean triadic cubes to those in the adapted cubes.

**Lemma 2.10.** *There exists  $C(d) < \infty$  such that, for every  $y \in \mathbb{R}^d$  and  $k, n, m \in \mathbb{N}$  with  $k < n < m$ ,*

$$\overline{\mathbf{A}}(y + \diamond_n) \leq \overline{\mathbf{A}}(\square_k) + C\Pi 3^{k-n} \mathbf{E}_0 \quad (2.107)$$

and

$$\overline{\mathbf{A}}(\square_m) \leq \overline{\mathbf{A}}(\diamond_n) + C\Pi 3^{n-m} \mathbf{E}_0. \quad (2.108)$$

*Proof.* Fix  $n, k \in \mathbb{N}$  with  $k \leq n$ . We partition the adapted cube  $\diamond_n$  as

$$\diamond_n = V \cup (\diamond_n \setminus V)$$

where

$$V := \bigcup \{z + \square_k : z \in 3^k \mathbb{Z}^d, z + \square_{k+1} \subseteq \diamond_n\}.$$

The boundary layer  $\diamond_n \setminus V$  has thickness at most  $\sqrt{d} 3^{k+1}$ , and so its volume is at most the perimeter of  $\diamond_n$  times  $C 3^{k+1}$ . Therefore,

$$\frac{|\diamond_n \setminus V|}{|\diamond_n|} \leq C\Pi^{1/2} 3^{k-n}.$$

Subadditivity yields

$$\begin{aligned} \mathbf{A}(\diamond_n) & \leq \frac{|V|}{|\diamond_n|} \mathbf{A}(V) + \frac{|\diamond_n \setminus V|}{|\diamond_n|} \mathbf{A}((y + \diamond_n) \setminus V) \\ & \leq \sum_{z \in 3^k \mathbb{Z}^d, z + \square_{k+1} \subseteq \diamond_n} \mathbf{A}(z + \square_k) + C\Pi^{1/2} 3^{k-n} \mathbf{A}(\diamond_n \setminus V). \end{aligned}$$

Following the the proof of Lemma 2.8 above, we use a Whitney-type partition of the boundary layer  $\diamond_n \setminus V$  into triadic subcubes and then apply subadditivity and (2.67) to obtain

$$|\mathbf{E}_0^{-1/2} \mathbf{A}(\diamond_n \setminus V) \mathbf{E}_0^{-1/2}| \leq \mathcal{O}_{\Psi_S}(C\Pi^{1/2}).$$

Combining the above and taking expectations, we obtain (2.107).

To get a bound in the opposite direction, we need to partition the cube  $\square_m$  into cubes of the form  $y' + \diamond_n$  with  $y' \in 3^n \mathbb{L}_0$ , plus a small boundary layer. We write

$$W := \bigcup \{y' + \diamond_n : y' \in 3^n \mathbb{L}_0, y' + \diamond_{n+1} \subseteq \square_m\}.$$

We have the following upper bound for the volume fraction of the boundary layer  $\square_m \setminus W$ :

$$\frac{|\square_m \setminus W|}{|\square_m|} \leq C\Pi^{1/2} 3^{n-m}.$$

Subadditivity yields

$$\begin{aligned} \mathbf{A}(\square_m) &\leq \mathbf{A}(W) + \frac{|\square_m \setminus W|}{|\square_m|} (\mathbf{A}(\square_m \setminus W) - \mathbf{A}(W)) \\ &\leq \sum_{y' \in 3^n \mathbb{L}_0, y' + \diamond_n \subseteq \square_m} \mathbf{A}(y' + \diamond_n) + C\Pi^{1/2} 3^{n-m} \mathbf{A}(\square_m \setminus W). \end{aligned}$$

Using once again a Whitney-type partition of the boundary layer  $\square_m \setminus W$ , subadditivity and the estimate (2.67), we obtain

$$\mathbf{A}(\square_m \setminus W) \leq \mathcal{O}_{\Psi_S}(C\Pi^{1/2}).$$

Combining the above and taking expectations, we get

$$\overline{\mathbf{A}}(\square_m) \leq \sum_{y' \in 3^n \mathbb{L}_0, y' + \diamond_n \subseteq \square_m} \overline{\mathbf{A}}(y' + \diamond_n) + C\Pi 3^{n-m} = \overline{\mathbf{A}}(\diamond_n) + C\Pi 3^{n-m}.$$

This completes the proof of (2.108) and thus of the lemma.  $\square$

**2.8. Embeddings into fractional Sobolev and Besov spaces.** In this subsection we show that the space  $\mathcal{A}(U)$  of solution embeds into certain fractional Besov spaces, provided that certain bounds on the coarse-grained matrices are satisfied.

For each  $s \in (0, 1)$ ,  $p \in [1, \infty)$ ,  $q \in [1, \infty)$  and  $n \in \mathbb{N}$ , we define a (volume-normalized) Besov seminorm in the cube  $\square_n$  by

$$[g]_{\underline{B}_{p,q}^s(\square_n)} := \left( \sum_{k=-\infty}^n \left( 3^{-spk} \sum_{z \in 3^{k-1} \mathbb{Z}^d, z + \square_k \subseteq \square_n} \|g - (g)_{z + \square_k}\|_{\underline{L}^p(z + \square_k)}^p \right)^{q/p} \right)^{1/q}. \quad (2.109)$$

In the case  $q = \infty$ , we define the Besov seminorms for every  $s \in [0, 1]$  and  $p \in [1, \infty)$  by

$$[g]_{\underline{B}_{p,\infty}^s(\square_n)} := \sup_{k \in (-\infty, n] \cap \mathbb{Z}} 3^{-sk} \left( \sum_{z \in 3^{k-1} \mathbb{Z}^d, z + \square_k \subseteq \square_n} \|g - (g)_{z + \square_k}\|_{\underline{L}^p(z + \square_k)}^p \right)^{1/p}. \quad (2.110)$$

The corresponding (volume-normalized) Besov norms are defined by

$$\|g\|_{\underline{B}_{p,q}^s(\square_n)} := 3^{-sn} |(g)_{\square_n}| + [g]_{\underline{B}_{p,q}^s(\square_n)}. \quad (2.111)$$

The Banach space  $B_{p,q}^s(\square_n)$  is defined to be the closure of  $C^\infty(\overline{\square_n})$  with respect to  $\|\cdot\|_{B_{p,q}^s(\square_n)}$ .

Note that  $s \in \{0, 1\}$  is allowed if  $q = \infty$ , and we actually obtain the more familiar Sobolev spaces. Indeed,  $[\cdot]_{\underline{B}_{p,\infty}^1(\square_n)}$  is equivalent to (volume-normalized)  $W^{1,p}(\square_n)$  seminorm and, similarly,  $[\cdot]_{\underline{B}_{p,\infty}^0(\square_n)}$  is equivalent to the volume-normalized  $L^p(\square_n)$  norm modulo constants:

$$\|g - (g)_{\square_n}\|_{\underline{L}^p(\square_n)} \leq [g]_{\underline{B}_{p,\infty}^0(\square_n)} \leq C(d)\|g - (g)_{\square_n}\|_{\underline{L}^p(\square_n)}.$$

We also work with weak Besov norms with negative regularity exponents which are defined as the dual spaces of  $B_{p,q}^s$ . For every  $s \in (0, 1]$ ,  $p \in [1, \infty]$  and  $q \in [1, \infty]$ , we let  $p'$  and  $q'$  denote the Hölder conjugate exponents of  $p$  and  $q$ , respectively, and we define

$$[f]_{\underline{B}_{p,q}^{-s}(\square_n)} := \sup \left\{ \int_{\square_n} fg : g \in B_{p',q'}^s(\square_n), \|g\|_{B_{p',q'}^s(\square_n)} \leq 1 \right\}. \quad (2.112)$$

The dual space of the subspace of  $B_{p',q'}^s(\square_n)$  with zero boundary values is defined by

$$[f]_{\underline{B}_{p,q}^{-s}(\square_n)} := \sup \left\{ \int_{\square_n} fg : g \in C_c^\infty(\square_n), [g]_{\underline{B}_{p',q'}^s(\square_n)} \leq 1 \right\}. \quad (2.113)$$

Finally, we introduce another variant of these negative spaces by defining

$$[f]_{\underline{\hat{B}}_{p,q}^{-s}(\square_n)} := 3^{d+s} \left( \sum_{k=-\infty}^n \left( 3^{spk} \sum_{z \in 3^k \mathbb{Z}^d \cap \square_n} |(f)_{z+\square_k}|^p \right)^{q/p} \right)^{1/q}. \quad (2.114)$$

The latter definition will sometimes be useful when estimating the negative seminorms from above, since we have that, for every  $f$ ,

$$[f]_{\underline{B}_{p,q}^{-s}(\square_n)} \leq [f]_{\underline{\hat{B}}_{p,q}^{-s}(\square_n)} \leq [f]_{\underline{B}_{p,q}^{-s}(\square_n)}. \quad (2.115)$$

The first inequality in (2.115) is immediate from the definitions and the second inequality is a consequence of Lemma A.2 in the appendix.

In the next lemma, we use the coarse-graining inequalities (2.38) and (2.39) to obtain embeddings of the solution space  $\mathcal{A}(U)$  into Besov spaces. In fact, we show that bounds on the coarse-grained matrices imply bounds on the weak Besov norms of the gradient and flux of a solution. In other words, we obtain weak, spatially-averaged information about an arbitrary solution in terms of the total energy of the solution and the coarse-grained matrices. Note that the estimates in (2.116) and (2.117) are obvious if the we replace the coarse grained matrices in each cube by the supremum of  $\mathbf{s}^{-1}$  and  $\mathbf{b}$ , respectively, in that cube. That we are able to prove the estimate *without* using pointwise information about the coefficient field, and instead relying on the coarse-grained matrices, is extremely important to our approach.

**Lemma 2.11.** *For every  $s \in [0, 1)$ ,  $n \in \mathbb{N}$  and  $u \in \mathcal{A}(\square_n)$ ,*

$$[\nabla u]_{\underline{\hat{B}}_{2,1}^{-s}(\square_n)} \leq 3^{d+s} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\square_n)} \sum_{k=-\infty}^n 3^{sk} \max_{z \in 3^k \mathbb{Z}^d \cap \square_n} |\mathbf{s}_*^{-1}(z + \square_k)|^{1/2} \quad (2.116)$$

and

$$[\mathbf{a} \nabla u]_{\underline{\hat{B}}_{2,1}^{-s}(\square_n)} \leq 3^{d+s} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\square_n)} \sum_{k=-\infty}^n 3^{sk} \max_{z \in 3^k \mathbb{Z}^d \cap \square_n} |\mathbf{b}(z + \square_k)|^{1/2}. \quad (2.117)$$

*Proof.* For the gradient we have by (2.38) that

$$\begin{aligned}
& \sum_{k=-\infty}^n 3^{sk} \left( \sum_{z \in 3^k \mathbb{Z}^d \cap \square_n} |(\nabla u)_{z+\square_k}|^2 \right)^{1/2} \\
& \leq \sum_{k=-\infty}^n 3^{sk} \left( \sum_{z \in 3^k \mathbb{Z}^d \cap \square_n} |\mathbf{s}_*^{-1}(z+\square_k)| |\mathbf{s}_*^{1/2}(z+\square_k)(\nabla u)_{z+\square_k}|^2 \right)^{1/2} \\
& \leq \sum_{k=-\infty}^n 3^{sk} \max_{z \in 3^k \mathbb{Z}^d \cap \square_n} |\mathbf{s}_*^{-1}(z+\square_k)|^{1/2} \left( \sum_{z \in 3^k \mathbb{Z}^d \cap \square_n} \int_{z+\square_k} \nabla u \cdot \mathbf{s} \nabla u \right)^{1/2} \\
& = \left( \int_{\square_n} \nabla u \cdot \mathbf{s} \nabla u \right)^{1/2} \sum_{k=-\infty}^n 3^{sk} \max_{z \in 3^k \mathbb{Z}^d \cap \square_n} |\mathbf{s}_*^{-1}(z+\square_k)|^{1/2}.
\end{aligned}$$

This is (2.116). The bound (2.117) for the flux follows similarly, using (2.39).  $\square$

The previous lemma implies a *coarse-grained Poincaré inequality* for solutions, because the left side of (2.116) actually control a *positive* Besov norms of  $u - (u)_{\square_n}$ . The latter assertion is a purely functional analytic fact which is given in Lemma A.3 in Appendix A.

**Lemma 2.12** (Coarse-grained Poincaré inequality). *There exists a constant  $C(d) < \infty$  such that, for every  $s \in [0, 1)$ ,  $n \in \mathbb{N}$  and  $u \in \mathcal{A}(\diamond_n)$ ,*

$$\|u - (u)_{\square_n}\|_{\underline{B}_{2,\infty}^s(\square_n)} \leq C \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\square_n)} \sum_{k=-\infty}^n 3^{(1-s)k} \max_{z \in 3^k \mathbb{Z}^d \cap \square_n} |\mathbf{s}_*^{-1}(z+\square_k)|^{1/2}. \quad (2.118)$$

In particular, under assumption (P2), we have that, for every  $s \in [0, 1 - \gamma/2)$ ,  $n \in \mathbb{N}$  with  $3^n \geq \mathcal{S}$  and  $u \in \mathcal{A}(\square_n)$ ,

$$\|u - (u)_{\square_n}\|_{\underline{B}_{2,\infty}^s(\square_n)} \leq \frac{C \lambda^{-1/2} 3^{(1-s)n}}{2 - 2s - \gamma} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\square_n)}.$$

*Proof.* We combine (A.5) in Lemma A.3 with (2.116) to obtain

$$\|u - (u)_{\square_n}\|_{\underline{B}_{2,\infty}^s(\square_n)} \leq C \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\square_n)} \sum_{k=-\infty}^n 3^{(1-s)k} \max_{z \in 3^k \mathbb{Z}^d \cap \square_n} |\mathbf{s}_*^{-1}(z+\square_k)|^{1/2}.$$

Thus, if  $3^n \geq \mathcal{S}$ , then we use (P2) and recall the definition of  $\lambda$  in (1.26) to get

$$\begin{aligned}
\|u - (u)_{\square_n}\|_{\underline{B}_{2,\infty}^s(\square_n)} & \leq C \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\square_n)} \sum_{k=-\infty}^n 3^{(1-s)k} \max_{z \in 3^k \mathbb{Z}^d \cap \square_n} |\mathbf{s}_*^{-1}(z+\square_k)|^{1/2} \\
& \leq C \lambda^{-1/2} 3^{(1-s)n} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\square_n)} \sum_{k=-\infty}^n 3^{(1-s-\gamma/2)(k-n)} \\
& \leq \frac{C \lambda^{-1/2} 3^{(1-s)n}}{2 - 2s - \gamma} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\square_n)}.
\end{aligned}$$

This completes the proof of (2.118).  $\square$

The next lemma says that, if a solution  $u \in \mathcal{A}(\square_n)$  has the property that its gradient and flux belong to compatible Besov spaces, then we are able to test the equation with  $\varphi u$  and integrate by parts. In view of the above lemmas, this allows us to justify basic energy estimates in our more general (non-uniformly elliptic) framework—such as the Caccioppoli inequality—opening up the way for basic elliptic theory.

**Lemma 2.13.** *Let  $n \in \mathbb{N}$ ,  $s \in (0, 1)$  and  $\varepsilon \in (0, 1 - s)$ . If  $u \in \mathcal{A}(\square_n)$  is such that*

$$[u]_{\underline{B}_{2,\infty}^{s+\varepsilon}(\square_n)} + [\mathbf{a}\nabla u]_{\underline{B}_{2,1}^{-s}(\square_n)} < \infty, \quad (2.119)$$

*then, for every  $\varphi \in C_c^\infty(\square_n)$ ,*

$$\oint_{\square_n} \varphi \mathbf{s} \nabla u \cdot \nabla u = - \oint_{\square_n} u \mathbf{a} \nabla u \cdot \nabla \varphi. \quad (2.120)$$

*Proof.* Without loss of generality, we may assume that  $(u)_{\square_n} = 0$ . For  $k \in \mathbb{N}$  with  $k \geq 10$ , we set  $u_k := (u \wedge k) \vee (-k)$  and  $v_k := u - u_k$ . Observe that  $v_k \rightarrow 0$  in  $L^2(\square_n)$  as  $k \rightarrow \infty$ . Let  $\varphi \in C_c^\infty(\square_n)$ . Then  $u_k \varphi \in H_{s,0}^1(\square_n)$ , because  $u_k$  is bounded, and since  $u \in \mathcal{A}(\square_n)$ ,

$$\begin{aligned} 0 &= \int_{\square_n} \mathbf{a} \nabla u \cdot \nabla (u_k \varphi) = \int_{\square_n} \varphi \mathbf{a} \nabla u \cdot \nabla u_k + \int_{\square_n} u_k \mathbf{a} \nabla u \cdot \nabla \varphi \\ &= \int_{\square_n} \varphi \mathbf{s} \nabla u_k \cdot \nabla u_k + \int_{\square_n} u_k \mathbf{a} \nabla u \cdot \nabla \varphi. \end{aligned} \quad (2.121)$$

We will argue that (2.119) allows us to pass to the limit  $k \rightarrow \infty$  in (2.121) and thereby obtain (2.120). We first observe that the first term on the right side of (2.121) converges to the energy of  $u$  by the monotone convergence theorem. To show that we can pass to the limit in the second term on the right, we use duality to get

$$\left| \int_{\square_n} v_k \mathbf{a} \nabla u \cdot \nabla \varphi \right| \leq [\mathbf{a} \nabla u]_{\underline{B}_{2,1}^{-s}(\square_n)} \|v_k \nabla \phi\|_{\underline{B}_{2,\infty}^s(\square_n)}.$$

The first factor on the right is finite by assumption (2.119). We will show that the second factor on the right vanishes in the limit  $k \rightarrow \infty$ . To that end, we apply (A.8), which gives

$$\begin{aligned} &\|v_k \nabla \phi\|_{\underline{B}_{2,\infty}^s(\square_n)} \\ &\leq C 3^{-sn} (3^n \|\nabla^2 \varphi\|_{L^\infty(\square_n)} + \|\nabla \varphi\|_{L^\infty(\square_n)}) \|v_k\|_{\underline{L}^2(\square_n)} + C \|\nabla \varphi\|_{L^\infty(\square_n)} [v_k]_{\underline{B}_{2,\infty}^s(\square_n)}. \end{aligned} \quad (2.122)$$

The first term on the right side of (2.122) converges to zero as  $k \rightarrow \infty$  since, as mentioned above,  $v_k \rightarrow 0$  in  $L^2(\square_n)$ . For the second term, we use that

$$\begin{aligned} \|u_k - (u_k)_{z+\square_j}\|_{\underline{L}^2(z+\square_j)}^2 &\leq \oint_{z+\square_j} \oint_{z+\square_j} |u_k(x) - u_k(y)|^2 dx dy \\ &\leq \oint_{z+\square_j} \oint_{z+\square_j} |u(x) - u(y)|^2 dx dy \leq 4 \|u - (u)_{z+\square_j}\|_{\underline{L}^2(z+\square_j)}^2. \end{aligned}$$

Select  $\tilde{\varepsilon} \in (0, 1)$  so that  $s(1 - \tilde{\varepsilon})^{-1} = s + \varepsilon$ , and then use the Hölder inequality to obtain that

$$\begin{aligned} [v_k]_{\underline{B}_{2,\infty}^s(\square_n)} &= \sup_{j \in (-\infty, n] \cap \mathbb{Z}} 3^{sj} \left( \sum_{z \in 3^j \mathbb{Z}^d \cap \square_n} \|v_k - (v_k)_{z+\square_j}\|_{\underline{L}^2(z+\square_j)}^2 \right)^{1/2} \\ &\leq \sup_{j \in (-\infty, n] \cap \mathbb{Z}} 3^{sj} \left( \sum_{z \in 3^j \mathbb{Z}^d \cap \square_n} \|v_k - (v_k)_{z+\square_j}\|_{\underline{L}^2(z+\square_j)}^{2(1-\tilde{\varepsilon})} \|v_k\|_{\underline{L}^2(z+\square_j)}^{2\tilde{\varepsilon}} \right)^{1/2} \\ &\leq \|v_k\|_{\underline{L}^2(\square_n)}^{\tilde{\varepsilon}} \sup_{j \in (-\infty, n] \cap \mathbb{Z}} 3^{sj} \left( \sum_{z \in 3^j \mathbb{Z}^d \cap \square_n} \|v_k - (v_k)_{z+\square_j}\|_{\underline{L}^2(z+\square_j)}^2 \right)^{(1-\tilde{\varepsilon})/2} \\ &\leq \|v_k\|_{\underline{L}^2(\square_n)}^{\tilde{\varepsilon}} \sup_{j \in (-\infty, n] \cap \mathbb{Z}} 3^{sj} \left( 4 \sum_{z \in 3^j \mathbb{Z}^d \cap \square_n} \|u - (u)_{z+\square_j}\|_{\underline{L}^2(z+\square_j)}^2 \right)^{(1-\tilde{\varepsilon})/2} \\ &\leq 2 \|v_k\|_{\underline{L}^2(\square_n)}^{\tilde{\varepsilon}} [u]_{\underline{B}_{2,\infty}^{s+\varepsilon}(\square_n)}^{1-\tilde{\varepsilon}}. \end{aligned}$$



In view of the assumption (2.119) and the fact that  $v_k \rightarrow 0$  in  $L^2(\square_n)$ , we deduce that  $[v_k]_{\underline{B}_{2,\infty}^s(\square_n)} \rightarrow 0$  as  $k \rightarrow \infty$  and thus the second term on the right side of (2.122) vanishes in the limit  $k \rightarrow \infty$ . Thus, we may pass to the limit  $k \rightarrow \infty$  in (2.121) to obtain (2.120). The proof is complete.  $\square$

We record here some analogues of the above estimates in the  $\mathbf{m}_0$ -adapted geometry, since these will be needed in what follows. We will not give the proofs, since they can be obtained by repeating the arguments above, or by applying the statements above after performing an affine change of coordinates.

- We first extend the Besov norms with positive regularity, defined in (2.123), to  $\diamond_n$  by defining, for every  $s \in (0, 1)$ ,  $p \in [1, \infty)$ ,  $q \in [1, \infty)$  and  $n \in \mathbb{N}$ ,

$$[g]_{\underline{B}_{p,q}^s(\diamond_n)} := \left( \sum_{k=-\infty}^n \left( 3^{-spk} \sum_{z \in 3^{k-1}\mathbb{L}_0, z+\diamond_k \subseteq \diamond_n} \|g - (g)_{z+\diamond_k}\|_{\underline{L}^p(z+\diamond_k)}^p \right)^{q/p} \right)^{1/q}. \quad (2.123)$$

We also define  $[g]_{\underline{B}_{p,\infty}^s(\diamond_n)}$  similarly, in analogy to (2.110).

- The negative Besov norms are defined following (2.112), (2.113) and (2.114):

$$[f]_{\underline{B}_{p,q}^{-s}(\diamond_n)} := \sup \left\{ \int_{\diamond_n} fg : g \in B_{p',q'}^s(\diamond_n), \|g\|_{\underline{B}_{p',q'}^s(\diamond_n)} \leq 1 \right\}, \quad (2.124)$$

$$[f]_{\underline{B}_{p,q}^{-s}(\diamond_n)} := \sup \left\{ \int_{\diamond_n} fg : g \in C_c^\infty(\square_n), [g]_{\underline{B}_{p',q'}^s(\diamond_n)} \leq 1 \right\}, \quad (2.125)$$

and

$$[f]_{\dot{\underline{B}}_{p,q}^{-s}(\diamond_n)} := 3^{d+s} \left( \sum_{k=-\infty}^n \left( 3^{spk} \sum_{z \in 3^k\mathbb{L}_0 \cap \diamond_n} |(f)_{z+\diamond_k}|^p \right)^{q/p} \right)^{1/q}. \quad (2.126)$$

For every  $s \in (0, 1]$ ,  $p \in [1, \infty)$  and  $q \in [1, \infty)$ , we have that

$$[f]_{\underline{B}_{p,q}^{-s}(\diamond_n)} \leq [f]_{\dot{\underline{B}}_{p,q}^{-s}(\diamond_n)} \leq [f]_{\dot{\underline{B}}_{p,q}^{-s}(\diamond_n)}. \quad (2.127)$$

- The statement of Lemma A.3 is modified as follows: there exists a constant  $C(d) < \infty$  such that, for every  $n \in \mathbb{N}$ ,  $s \in [0, 1)$  and  $u \in H_s^1(\diamond_n)$ , we have that

$$\|u - (u)_{\diamond_n}\|_{\underline{B}_{2,\infty}^s(\diamond_n)} \leq C[\mathbf{q}_0 \nabla u]_{\dot{\underline{B}}_{2,1}^{s-1}(\diamond_n)}, \quad (2.128)$$

and, if  $\phi \in C_c^\infty(\diamond_n)$  satisfies  $3^n \|\mathbf{q}_0 \nabla \phi\|_{L^\infty(\diamond_n)} + 3^{2n} \|(\mathbf{q}_0 \nabla)^2 \phi\|_{L^\infty(\diamond_n)} \leq 1$ , then

$$\|(u - (u)_{\diamond_n}) \mathbf{m}_0^{1/2} \nabla \phi\|_{\underline{B}_{2,\infty}^s(\diamond_n)} \leq C 3^{-n} [\mathbf{m}_0^{1/2} \nabla u]_{\dot{\underline{B}}_{2,1}^{s-1}(\diamond_n)}. \quad (2.129)$$

- The statement of Lemma 2.11 can be modified as follows. For every  $s \in (0, 1)$ ,  $n \in \mathbb{N}$  and  $u \in \mathcal{A}(\diamond_n)$ , we have that

$$[\mathbf{m}_0^{1/2} \nabla u]_{\dot{\underline{B}}_{2,1}^{-s}(\diamond_n)} \leq \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_n)} \sum_{k=-\infty}^n 3^{sk} \max_{z \in 3^k\mathbb{L}_0 \cap \diamond_n} |\mathbf{m}_0 \mathbf{s}_*^{-1}(z + \diamond_k)|^{1/2} \quad (2.130)$$

and

$$[\mathbf{m}_0^{-1/2} \mathbf{a} \nabla u]_{\dot{\underline{B}}_{2,1}^{-s}(\diamond_n)} \leq \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_n)} \sum_{k=-\infty}^n 3^{sk} \max_{z \in 3^k\mathbb{L}_0 \cap \diamond_n} |\mathbf{m}_0^{-1} \mathbf{b}(z + \diamond_k)|^{1/2}. \quad (2.131)$$

**2.9. Gradient and flux estimates in weak norms.** We conclude this subsection by a lemma, which ties the weak norms and coarsened coefficients in a very precise way. The statement is a bit ugly at first glance, but the explicit form of the estimate will prove useful.

**Lemma 2.14.** *Let  $\rho \in (0, 2)$ ,  $s \in (\rho/2, 1]$  and  $h, n \in \mathbb{N}$  with  $h < n$ . Also let  $\mathbf{m} \in \mathbb{R}^{d \times d}$  be positive and symmetric and  $\mathbf{h} \in \mathbb{R}^{d \times d}$  be antisymmetric, and denote*

$$\mathbf{M} := \begin{pmatrix} \mathbf{m} + \mathbf{h}^t \mathbf{m}^{-1} \mathbf{h} & -\mathbf{h}^t \mathbf{m} \\ -\mathbf{m} \mathbf{h} & \mathbf{m}^{-1} \end{pmatrix}.$$

Given a symmetric and positive matrix  $\mathbf{E} \in \mathbb{R}^{2d \times 2d}$ , define the random variable

$$\mathcal{M}_{n,\rho} := \sup_{k \in \mathbb{Z} \cap (-\infty, n]} 3^{-\rho(n-k)} \max_{z \in 3^k \mathbb{L}_0 \cap \Diamond_n} |(\mathbf{E}^{-1/2}(\mathbf{A}(z + \Diamond_k) - \mathbf{E})\mathbf{E}^{-1/2})_+|. \quad (2.132)$$

Then there exists a universal constant  $C < \infty$  such that, for every  $\delta \in (0, 1]$  and  $p, q \in \mathbb{R}^d$ , by writing  $v_n := v(\cdot, \Diamond_n, p, q)$ ,

$$\begin{aligned} & 3^{-sn} \left[ \mathbf{M}^{1/2} \begin{pmatrix} \nabla v_n - (\nabla v_n)_{\Diamond_n} \\ \mathbf{a} \nabla v_n - (\mathbf{a} \nabla v_n)_{\Diamond_n} \end{pmatrix} \right]_{\underline{B}_{2,1}^{-s}(\Diamond_n)} \\ & \leq C 3^d |\mathbf{M}^{-1/2} \mathbf{E} \mathbf{M}^{-1/2}|^{1/2} \left| \mathbf{E}^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right| \sum_{k=n-h}^n 3^{s(k-n)} \left( \sum_{z \in 3^k \mathbb{L}_0 \cap \Diamond_n} |\mathbf{E}^{-1/2}(\mathbf{A}(z + \Diamond_k) - \mathbf{A}(\Diamond_n))\mathbf{E}^{-1/2}|^2 \right)^{1/2} \\ & \quad + C 3^d |\mathbf{M}^{-1/2} \mathbf{E} \mathbf{M}^{-1/2}|^{1/2} \left| \mathbf{E}^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right| \sum_{k=n-h}^n 3^{s(k-n)} \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \Diamond_n} \mathbf{E}^{-1/2}(\mathbf{A}(z + \Diamond_k) - \mathbf{A}(\Diamond_n))\mathbf{E}^{-1/2} \right|^{1/2} \\ & \quad + \frac{C 3^d \delta^{-1/2}}{s - \rho/2} \left( \mathbf{1}_{\{\mathcal{M}_{n,\rho} > \delta\}} \mathcal{M}_{n,\rho}^{1/2} + 3^{-(s-\rho/2)h} \mathbf{1}_{\{\mathcal{M}_{n,\rho} \leq \delta\}} \right) |\mathbf{M}^{-1/2} \mathbf{E} \mathbf{M}^{-1/2}|^{1/2} \|\mathbf{s}^{1/2} \nabla v_n\|_{\underline{L}^2(\Diamond_n)}. \quad (2.133) \end{aligned}$$

*Proof.* As in the statement, we write  $v_n := v(\cdot, \Diamond_n, p, q)$  to shorten the notation. Fix  $\delta \in (0, \infty)$ ,  $\rho \in (0, 2)$ ,  $s \in (\rho/2, 1]$  and  $n, h \in \mathbb{N}$  with  $h < n$ . We suppress  $\rho, \delta$  from the notation with  $\mathcal{M}_{\rho, \delta}$ . Fix also  $p, q \in \mathbb{R}^d$ . We denote, for any Lipschitz domain  $U$ ,

$$X(\cdot, U, p, q) := \begin{pmatrix} \nabla v(\cdot, U, p, q) \\ \mathbf{a} \nabla v(\cdot, U, p, q) \end{pmatrix}. \quad (2.134)$$

Moreover, for every  $k \in \mathbb{Z}$  and  $z \in \mathbb{R}^d$ , we denote  $X_{z,k} := X(\cdot, z + \Diamond_k, p, q)$  and, for  $z = 0$ , we suppress  $z$  from the notation and write  $X_k = X_{0,k}$ . We also denote  $v_{z,k} := v(\cdot, z + \Diamond_k, p, q)$ .

*Step 1.* We first show that, for any Lipschitz domains  $U, V$ ,

$$\begin{aligned} & \left| \mathbf{M}^{1/2} \left( (X(\cdot, U, p, q))_U - (X(\cdot, V, p, q))_V \right) \right|^2 \\ & \leq |\mathbf{E}^{1/2} \mathbf{M}^{-1} \mathbf{E}^{1/2}| |\mathbf{E}^{-1/2}(\mathbf{A}(U) - \mathbf{A}(V))\mathbf{E}^{-1/2}|^2 \left| \mathbf{E}^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2. \quad (2.135) \end{aligned}$$

We have the following identity by (2.33):

$$(X(\cdot, U, p, q))_U = (\mathbf{R} \mathbf{A}(U) + \mathbf{I}_{2d}) \begin{pmatrix} -p \\ q \end{pmatrix} \quad \text{with } \mathbf{R} := \begin{pmatrix} 0 & \mathbf{I}_d \\ \mathbf{I}_d & 0 \end{pmatrix}. \quad (2.136)$$

$$\begin{aligned} & \left| \mathbf{M}^{1/2} \left( (X(\cdot, U, p, q))_U - (X(\cdot, V, p, q))_V \right) \right|^2 \\ &= \left| \begin{pmatrix} -p \\ q \end{pmatrix} \cdot (\mathbf{A}(U) - \mathbf{A}(V)) \mathbf{RMR} (\mathbf{A}(U) - \mathbf{A}(V)) \begin{pmatrix} -p \\ q \end{pmatrix} \right|, \end{aligned}$$

from which (2.135) follows since  $\mathbf{RMR} = \mathbf{M}^{-1}$ .

*Step 2.* We show that, for every  $X \in \mathcal{S}(\diamond_n)$ ,

$$\begin{aligned} & 3^{-sn} [\mathbf{M}^{1/2} X]_{\underline{B}_{2,1}^{-s}(\diamond_n)} \\ & \leq 3^d |\mathbf{E}^{1/2} \mathbf{M}^{-1} \mathbf{E}^{1/2}|^{1/2} \|\mathbf{A}^{1/2} X\|_{\underline{L}^2(\diamond_n)} \sum_{k=-\infty}^n 3^{s(k-n)} \max_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{E}^{-1/2} \mathbf{A}(z + \diamond_k) \mathbf{E}^{-1/2}|^{1/2}. \end{aligned} \quad (2.137)$$

To see this, we use  $\mathbf{R}^2 = \mathbf{I}_{2d}$ ,  $\mathbf{A}(U) = \mathbf{R} \mathbf{A}_*^{-1}(U) \mathbf{R}$ ,  $\mathbf{M}^{-1} = \mathbf{RMR}$ , noting that also implies  $\mathbf{M}^{1/2} = \mathbf{RM}^{-1/2} \mathbf{R}$ , and the fact that  $\mathbf{R}$  is unitary, to get

$$\begin{aligned} |\mathbf{M}^{1/2} \mathbf{A}_*^{-1}(z + \diamond_k) \mathbf{M}^{1/2}| &= |\mathbf{RM}^{-1/2} \mathbf{R} \mathbf{A}_*^{-1}(z + \diamond_k) \mathbf{RM}^{-1/2} \mathbf{R}| = |\mathbf{RM}^{-1/2} \mathbf{A}(z + \diamond_k) \mathbf{M}^{-1/2} \mathbf{R}| \\ &= |\mathbf{M}^{-1/2} \mathbf{A}(z + \diamond_k) \mathbf{M}^{-1/2}| \end{aligned}$$

Thus, we obtain by (2.40) that

$$\begin{aligned} \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{M}^{1/2}(X)_{z + \diamond_k}|^2 &\leq \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{A}_*^{-1/2}(z + \diamond_k) \mathbf{M} \mathbf{A}_*^{-1/2}(z + \diamond_k)| |\mathbf{A}_*^{1/2}(z + \diamond_k)(X)_{z + \diamond_k}|^2 \\ &= \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{M}^{1/2} \mathbf{A}_*^{-1}(z + \diamond_k) \mathbf{M}^{1/2}| |\mathbf{A}_*^{1/2}(z + \diamond_k)(X)_{z + \diamond_k}|^2 \\ &\leq |\mathbf{E}^{1/2} \mathbf{M}^{-1} \mathbf{E}^{1/2}| \|\mathbf{A}^{1/2} X\|_{\underline{L}^2(\diamond_n)}^2 \max_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{E}^{-1/2} \mathbf{A}(z + \diamond_k) \mathbf{E}^{-1/2}|, \end{aligned} \quad (2.138)$$

which gives us (2.137).

*Step 3.* We next show that

$$\begin{aligned} & \mathbf{1}_{\{\mathcal{M} > \delta\}} 3^{-sn} [\mathbf{M}^{1/2} (X_n - (X_n)_{\diamond_n})]_{\underline{B}_{2,1}^{-s}(\diamond_n)} \\ & \leq \frac{16 \cdot 3^d}{s - \rho/2} \frac{1 + \delta^{1/2}}{\delta^{1/2}} \mathbf{1}_{\{\mathcal{M} > \delta\}} \mathcal{M}^{1/2} |\mathbf{E}^{1/2} \mathbf{M}^{-1} \mathbf{E}^{1/2}|^{1/2} \|\mathbf{s}^{1/2} \nabla v_n\|_{\underline{L}^2(\diamond_n)}. \end{aligned} \quad (2.139)$$

In view of (2.137) and (2.138) we obtain that

$$\begin{aligned} & \mathbf{1}_{\{\mathcal{M} > \delta\}} 3^{-sn} [\mathbf{M}^{1/2} X_n]_{\underline{B}_{2,1}^{-s}(\diamond_n)} \\ & \leq 2 \cdot 3^d |\mathbf{E}^{1/2} \mathbf{M}^{-1} \mathbf{E}^{1/2}|^{1/2} \|\mathbf{A}^{1/2} X_n\|_{\underline{L}^2(\diamond_n)} \mathbf{1}_{\{\mathcal{M} > \delta\}} \sum_{k=-\infty}^n 3^{s(k-n)} (1 + \mathcal{M}^{1/2} 3^{\frac{\rho}{2}(n-k)}) \\ & \leq \frac{16 \cdot 3^d}{s - \rho/2} \frac{1 + \delta^{1/2}}{\delta^{1/2}} \mathcal{M}^{1/2} \mathbf{1}_{\{\mathcal{M} > \delta\}} |\mathbf{E}^{1/2} \mathbf{M}^{-1} \mathbf{E}^{1/2}|^{1/2} \|\mathbf{s}^{1/2} \nabla v_n\|_{\underline{L}^2(\diamond_n)}, \end{aligned} \quad (2.140)$$

proving (2.139).

*Step 4.* We conclude the proof by proving (2.133) under the event  $\{\mathcal{M} \leq \delta\}$ . First, by (2.138), we see that

$$\begin{aligned} & \mathbf{1}_{\{\mathcal{M} \leq \delta\}} \sum_{k=-\infty}^{n-h} 3^{s(k-n)} \left( \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{M}^{1/2}((X_n)_{z+\diamond_k} - (X_n)_{\diamond_n})|^2 \right)^{1/2} \\ & \leq 2 |\mathbf{E}^{1/2} \mathbf{M}^{-1} \mathbf{E}^{1/2}|^{1/2} \|\mathbf{s}^{1/2} \nabla v_n\|_{\underline{L}^2(\diamond_n)} \sum_{k=-\infty}^{n-h} 3^{s(k-n)} (1 + \delta^{1/2} 3^{\frac{\rho}{2}(n-k)}) \\ & \leq 4 \left( \frac{3^{-hs}}{s} + \frac{\delta^{1/2} 3^{-(s-\rho/2)h}}{s-\rho/2} \right) |\mathbf{E}^{1/2} \mathbf{M}^{-1} \mathbf{E}^{1/2}|^{1/2} \|\mathbf{s}^{1/2} \nabla v_n\|_{\underline{L}^2(\diamond_n)}. \end{aligned}$$

Second, using the triangle inequality, for each  $k \in \mathbb{Z}$  with  $n-h \leq k \leq n$ , we get that

$$\begin{aligned} & \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{M}^{1/2}((X_n)_{z+\diamond_k} - (X_n)_{\diamond_n})|^2 \\ & \leq 2 \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \left( |\mathbf{M}^{1/2}((X_{z,k})_{z+\diamond_k} - (X_n)_{\diamond_n})|^2 + |\mathbf{M}^{1/2}(X_n - X_{z,k})_{z+\diamond_k}|^2 \right). \end{aligned} \quad (2.141)$$

The contribution of the first term on the right side of (2.141) can be estimated using (2.135):

$$\begin{aligned} & |\mathbf{M}^{1/2}((X_{z,k})_{z+\diamond_k} - (X_n)_{\diamond_n})|^2 \\ & \leq |\mathbf{E}^{1/2} \mathbf{M}^{-1} \mathbf{E}^{1/2}| |\mathbf{E}^{-1/2} (\mathbf{A}(z+\diamond_k) - \mathbf{A}(\diamond_n)) \mathbf{E}^{-1/2}|^2 \left| \mathbf{E}^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2. \end{aligned} \quad (2.142)$$

We estimate the second term on the right side of (2.141) using the same computation as in (2.138), but now for  $X_n - X_{z,k}$  instead of  $X_n$ :

$$\begin{aligned} & \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{M}^{1/2}(X_n - X_{z,k})_{z+\diamond_k}|^2 \\ & \leq |\mathbf{E}^{1/2} \mathbf{M}^{-1} \mathbf{E}^{1/2}| \max_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{E}^{-1/2} \mathbf{A}(z+\diamond_k) \mathbf{E}^{-1/2}| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \|\mathbf{A}^{1/2}(X_n - X_{z,k})\|_{\underline{L}^2(z+\diamond_k)}^2. \end{aligned}$$

By the quadratic response (2.29), (2.30), the first variation (2.28) and (2.13), we have that

$$\begin{aligned} & \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \|\mathbf{A}^{1/2}(X_n - X_{z,k})\|_{\underline{L}^2(z+\diamond_k)}^2 = 2 \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (\|\mathbf{s}^{1/2} \nabla v_{z,k}\|_{\underline{L}^2(z+\diamond_k)}^2 - \|\mathbf{s}^{1/2} \nabla v_n\|_{\underline{L}^2(\diamond_n)}^2) \\ & = 4 \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \begin{pmatrix} -p \\ q \end{pmatrix} \cdot (\mathbf{A}(z+\diamond_k) - \mathbf{A}(\diamond_n)) \begin{pmatrix} -p \\ q \end{pmatrix} \\ & \leq 4 \left| \mathbf{E}^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2 \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{E}^{-1/2} (\mathbf{A}(z+\diamond_k) - \mathbf{A}(\diamond_n)) \mathbf{E}^{-1/2}|. \end{aligned}$$

Putting the above estimates together gives us (2.133) under the event  $\{\mathcal{M} \leq \delta\}$ . The proof is complete.  $\square$

### 3. Renormalization in high contrast

The purpose of this section is to give an estimate of the length scale at which a general elliptic coefficient field  $\mathbf{a}(x)$ , with (possibly very large) ellipticity ratio  $\Theta \in [1, \infty)$ , has homogenized to within a specified finite error. The precise statement is given in the following theorem.

**Theorem 3.1** (Homogenization in high contrast). *Let  $\alpha := (\min\{\nu, 1\} - \gamma)(1 - \beta)$ . There exists a constant  $C(d) < \infty$  such that, for every  $\sigma \in (0, \frac{1}{2}\Theta]$  and  $m \in \mathbb{N}$  satisfying*

$$m \geq \frac{C}{\sigma^2} \left( \log K_{\Psi_S} + \frac{1}{\alpha^2} \log \left( \frac{\Pi K_{\Psi}}{\alpha \sigma} \right) \right) \log^2(1 + \Theta), \quad (3.1)$$

*the renormalized ellipticity ratio  $\Theta_m$  defined in (2.84) satisfies*

$$\Theta_m - 1 \leq \sigma. \quad (3.2)$$

The reader is encouraged to ignore the details of the rather explicit form of (3.1), on first reading, and notice only that the theorem asserts that, for a constant  $C$  depending on the parameters  $(d, \gamma, \beta)$ , but not on  $\Theta$  or  $\Pi$ ,

$$m \geq C \log(1 + \max\{\Pi, K_{\Psi}, K_{\Psi_S}\}) \log^2(1 + \Theta) \implies \Theta_m - 1 \leq 10^{-6}.$$

If we wish, we can ignore also the parameters  $(K_{\Psi}, K_{\Psi_S})$  and use the trivial bound  $\Theta \leq \Pi$  to obtain that, for a constant  $C(d, \gamma, \beta, \nu, K_{\Psi}, K_{\Psi_S}) < \infty$ ,

$$m \geq C \log^3(1 + \Pi) \implies \Theta_m - 1 \leq 10^{-6}.$$

This matches the length scale appearing in the statement of Theorem A in the introduction.

Since the parameter  $\sigma$  in Theorem 3.1 can be taken arbitrarily small, the theorem statement provides an explicit convergence rate for  $\Theta_m - 1$ . However, this rate is not very useful when  $\sigma$  is small. The main role of the theorem is therefore to reduce  $\Theta_m - 1$  from a possibly very large number to a somewhat small number, say,  $10^{-6}$ . In the next section we use this estimate as a starting point for the derivation of a much better estimate on the rate of  $\Theta_m - 1$  to zero.

The main step in the proof of Theorem 3.1 lies in the following proposition, which formalizes one step of the renormalization procedure. It says that we can reduce the renormalized diffusivity by a constant factor by zooming out on the order of  $\log^2 \Pi$  many triadic scales.

**Proposition 3.2** (One renormalization step). *There exists a constant  $C(d) < \infty$  such that, for every  $\sigma \in (0, \frac{1}{2}]$  and  $m \in \mathbb{N}$  satisfying*

$$m \geq \frac{C}{\sigma^2} \left( \log K_{\Psi_S} + \frac{1}{\alpha^2} \log \left( \frac{\Pi K_{\Psi}}{\alpha \sigma} \right) \right) \log(1 + \Theta), \quad (3.3)$$

*where  $\alpha := (\min\{\nu, 1\} - \gamma)(1 - \beta)$ , we have*

$$\Theta_m - 1 \leq \sigma \Theta. \quad (3.4)$$

We have written the dependence of the lower bound on  $m$  in (3.3) explicitly in all parameters except for  $d$ . If we wish, we can write it in a nicer-looking (but less informative) way as

$$m \geq \frac{C |\log \sigma|}{\sigma^2} \log^2(1 + \Pi),$$

but now the constant  $C$  depends on  $(d, K_{\Psi_S}, K_{\Psi}, \gamma, \beta, \nu)$ , but not on  $\sigma$  nor on the ellipticity ratios  $\Pi$  and  $\Theta$ . Therefore Proposition 3.2 says, informally, that the renormalized ellipticity ratio is reduced by a constant factor if we zoom out on the order of  $\log^2(1 + \Pi)$  many geometric scales.

The proof of Proposition 3.2 is the main focus of this section. Once its proof is complete, we will iterate the statement on the order of  $\log(1 + \Theta)$  many times, renormalizing at each step with the help of Proposition 2.6, to obtain Theorem 3.1.

**3.1. A reduction: finding a good range of scales.** The first step in the proof of Proposition 3.2 is to make a reduction to the following statement.

**Proposition 3.3.** *Suppose that  $\delta, \sigma \in (0, 1/2]$  and  $l \in \mathbb{N}$  satisfy*

$$\max \left\{ 3^{-\frac{1}{4}(1-\beta)l} \Pi, K_{\Psi}^8 \Pi 3^{-\frac{1}{2}(\nu-\gamma)(1-\beta)l}, \frac{K_{\Psi}^{8d} \Pi^2}{(1-\gamma)^2} 3^{-l}, \frac{3^{-(1-\gamma)l}}{1-\gamma} \right\} \leq \delta \sigma^2. \quad (3.5)$$

*Suppose that  $m \in \mathbb{N}$  with  $m \geq 100l$  and that the matrix  $\mathbf{E}_0$  in (P2) satisfies*

$$\overline{\mathbf{A}}(\square_0) \leq \mathbf{E}_0 \quad \text{and} \quad |\mathbf{E}_0^{1/2} \overline{\mathbf{A}}^{-1}(\square_m) \mathbf{E}_0^{1/2} - \mathbf{I}_{2d}| \leq \delta \sigma^2. \quad (3.6)$$

*Then there exists a constant  $\delta_0(d) > 0$  such that  $\delta \leq \delta_0$  implies*

$$\Theta_m - 1 \leq \sigma \Theta. \quad (3.7)$$

The proof of Proposition 3.3 is given in Section 3.2, below. We first demonstrate that it implies Proposition 3.2, which relies on the following lemma. The idea is very simple: since the deterministic matrix  $\overline{\mathbf{A}}(\square_n)$  is monotone in  $n$  and its determinant is bounded between 1 and  $\Theta^d$ , we can find a sequence of consecutive  $n$  for which it does not change much.

**Lemma 3.4** (Pigeonhole lemma). *For every  $\delta_1 \in (0, 1/2]$  and  $m_1, l \in \mathbb{N}$  satisfying*

$$m_1 \geq \frac{4d \log(4\Theta_0)}{\delta_1} l, \quad (3.8)$$

*there exists  $m \in \mathbb{N} \cap [m_1, 2m_1]$  satisfying*

$$|\overline{\mathbf{A}}^{-1/2}(\square_m) \overline{\mathbf{A}}(\square_{m-l}) \overline{\mathbf{A}}^{-1/2}(\square_m) - \mathbf{I}_{2d}| \leq \delta_1. \quad (3.9)$$

*Proof.* For short, we denote  $\overline{\mathbf{A}}_j := \overline{\mathbf{A}}(\square_j)$  for every  $j \in \mathbb{N}$ . Observe that, for each  $m, n \in \mathbb{N}$  with  $n \leq m$ ,  $\det(\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_n \overline{\mathbf{A}}_m^{-1/2}) \geq 1$  by subadditivity. Fix  $K \in \mathbb{N}$  with  $K \geq 2$  and compute

$$\sum_{k=K}^{2K-1} \log \det(\overline{\mathbf{A}}_{(k+1)l}^{-1/2} \overline{\mathbf{A}}_{kl} \overline{\mathbf{A}}_{(k+1)l}^{-1/2}) = \log \det(\overline{\mathbf{A}}_{2Kl}^{-1/2} \overline{\mathbf{A}}_{Kl} \overline{\mathbf{A}}_{2Kl}^{-1/2}) \leq 2d \log(4\Theta_0).$$

Indeed, by letting  $\mathbf{h}_0$  being the minimizing matrix in (2.84) for  $n = Kl$ , using (2.23),  $\overline{\mathbf{s}}_{*,2Kl} \leq \overline{\mathbf{s}}_{2Kl}$  and  $\overline{\mathbf{A}}_{\mathbf{h}_0,2Kl} \leq \overline{\mathbf{A}}_{\mathbf{h}_0,Kl}$ , we obtain, by the fact that  $n \mapsto \Theta_n$  is monotone decreasing, that

$$|\overline{\mathbf{A}}_{2Kl}^{-1/2} \overline{\mathbf{A}}_{Kl} \overline{\mathbf{A}}_{2Kl}^{-1/2}| \leq 4(|\overline{\mathbf{s}}_{2Kl}^{-1/2} \mathbf{b}_{\mathbf{h}_0,Kl} \overline{\mathbf{s}}_{2Kl}^{-1/2}| \vee |\mathbf{b}_{\mathbf{h}_0,2Kl}^{1/2} \mathbf{s}_{*,Kl}^{-1} \mathbf{b}_{\mathbf{h}_0,2Kl}^{1/2}|) \leq 4\Theta_{Kl} \leq 4\Theta_0.$$

By the pigeonhole principle, we can find at least one element  $k \in \{K, \dots, 2K-1\}$  such that

$$\log |\overline{\mathbf{A}}_{(k+1)l}^{-1/2} \overline{\mathbf{A}}_{kl} \overline{\mathbf{A}}_{(k+1)l}^{-1/2}| \leq \log \det(\overline{\mathbf{A}}_{(k+1)l}^{-1/2} \overline{\mathbf{A}}_{kl} \overline{\mathbf{A}}_{(k+1)l}^{-1/2}) \leq \frac{d \log(4\Theta_0)}{K-1}.$$

If we impose the restriction that  $K \geq 4d \log(4\Theta_0)$ , then we may deduce from this that

$$|\overline{\mathbf{A}}_{(k+1)l}^{-1/2} \overline{\mathbf{A}}_{kl} \overline{\mathbf{A}}_{(k+1)l}^{-1/2} - \mathbf{I}_{2d}| \leq \frac{2d \log(4\Theta_0)}{K-1}.$$

Taking  $K := \lceil l^{-1} m_1 \rceil$  and, setting  $m := (k-1)l$ , we obtain, under the condition (3.8), the existence of  $m \in \{m_1, \dots, 2m_1\}$  satisfying

$$|\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_{m-l} \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}| \leq \frac{4dl \log(4\Theta_0)}{m_1} \leq \delta_1.$$

This completes the proof.  $\square$

*Reduction of Proposition 3.2 to Proposition 3.3.* Fix  $\delta, \sigma \in (0, 1/2]$  and let  $L(\delta, \sigma)$  be defined by

$$L(\delta, \sigma) := \left\lceil 8d \log K_{\Psi_S} + \frac{100d}{\alpha^2} \log \left( \frac{2^{10} \Pi K_{\Psi}}{\alpha \delta \sigma^2} \right) \right\rceil, \quad (3.10)$$

where we define  $\alpha := (\min\{\nu, 1\} - \gamma)(1 - \beta)$ . We apply Lemma 3.4 with the following choices of parameters:

$$\delta_1 := \frac{1}{2} \delta \sigma^2, \quad m_1 := \frac{8d \log(4\Theta_0)}{\delta_1} n \quad \text{and} \quad n := 100L(\delta, \sigma). \quad (3.11)$$

The lemma gives us an  $m \in \mathbb{N}$  satisfying

$$m_1 \leq m \leq 2m_1 = \frac{1600d \log(4\Theta_0)}{\delta_1} L(\delta, \sigma) \quad (3.12)$$

and

$$|\overline{\mathbf{A}}^{-1/2}(\square_m) \overline{\mathbf{A}}(\square_{m-2n}) \overline{\mathbf{A}}^{-1/2}(\square_m) - \mathbf{I}_{2d}| \leq \frac{1}{2} \delta \sigma^2. \quad (3.13)$$

We intend to apply Proposition 2.6 with parameters  $n_0 = m - n$ ,  $l_0 = n$ ,  $\rho = \frac{1}{2}(\min\{\nu, 1\} + \gamma)$  and  $\frac{1}{4} \delta \sigma^2$  instead of  $\delta$ . In order to apply the proposition, we need to check that the condition (2.86) is valid. This is however immediate from the choice  $n = 100L(\delta, \sigma)$  and the definition of  $L(\delta, \sigma)$  above. The application of the proposition yields that the probability measure  $\mathbb{P}_{m-n}$ , defined in (2.83) as the pushforward of  $\mathbb{P}$  under the dilation map  $\mathbf{a} \mapsto \mathbf{a}(3^{m-n} \cdot)$ , satisfies assumptions (P1), (P2) and (P3) with the new parameters

$$\begin{cases} \mathbf{E}_{\text{new}} := (1 + \frac{1}{4} \delta \sigma^2) \overline{\mathbf{A}}(\square_{m-2n}), \\ \gamma_{\text{new}} := \frac{1}{2}(\min\{1, \nu\} + \gamma) \\ K_{\Psi, \text{new}} := K_{\Psi} \\ K_{\Psi_S, \text{new}} := \max\{K_{\Psi_S}, K_{\Psi}^{\lceil 1/\mu \rceil}\}, \\ \Theta_{\text{new}} := (1 + \frac{1}{4} \delta \sigma^2)^2 \Theta_{m-n} \leq (1 + \delta \sigma^2) \Theta, \\ \Pi_{\text{new}} := 2^{10} \Pi. \end{cases} \quad (3.14)$$

We will now apply Proposition 3.3 with  $(\mathbb{P}_{m-n}, n)$  in place of  $(\mathbb{P}, m)$ . This requires that  $n \geq 100l$  for some  $l$  satisfying (3.5) with the new parameters in (3.14). For this we take  $l = L(\delta, \sigma)$ , we note that  $n = 100l$  and that (3.5) with the new parameters is valid by the definition of  $L(\delta, \sigma)$  in (3.10). To verify the final hypothesis (3.6), we observe that have that  $\overline{\mathbf{A}}(\square_{m-n}) \leq \mathbf{E}_{\text{new}}$  and

$$\begin{aligned} |\overline{\mathbf{A}}^{-1/2}(\square_m) \mathbf{E}_{\text{new}} \overline{\mathbf{A}}^{-1/2}(\square_m) - \mathbf{I}_{2d}| &\leq (1 + \frac{1}{4} \delta \sigma^2) |\overline{\mathbf{A}}^{-1/2}(\square_m) \overline{\mathbf{A}}(\square_{m-2n}) \overline{\mathbf{A}}^{-1/2}(\square_m) - \mathbf{I}_{2d}| + \frac{1}{4} \delta \sigma^2 \\ &\leq \delta \sigma^2. \end{aligned} \quad (3.15)$$

We note that  $\overline{\mathbf{A}}(\square_m)$  for  $\mathbb{P}$  is the same as  $\overline{\mathbf{A}}(\square_n)$  for  $\mathbb{P}_{m-n}$ . The application of Proposition 3.3 therefore yields that  $\Theta_m$  (which is  $\Theta_n$  for  $\mathbb{P}_{m-n}$ ) satisfies

$$\Theta_m - 1 \leq \sigma \Theta_{\text{new}} \leq 4\sigma \Theta. \quad (3.16)$$

This yields (3.4) with  $4\sigma$  in place of  $\sigma$ ; the factor of 4 may be removed by shrinking  $\delta$  by a factor of two. By the upper bound on  $m$  in (3.12) and the fact that  $k \mapsto \Theta_k$  is monotone nonincreasing, the proof is now complete.  $\square$



**3.2. One renormalization step.** We present the proof of Proposition 3.3 in this subsection. Throughout, we work with the following fixed parameters:

- $\delta_0 = \delta_0(d) \in (0, 1/2]$  is a small constant which will be selected at the end of the proof;
- $\sigma \in (0, 1/2]$  and  $\delta \in (0, \delta_0]$  are a given constants in the statement of Proposition 3.3.
- We fix an integer  $l \in \mathbb{N}$  representing a mesoscopic scale; we take it to be the smallest positive integer satisfying the condition (3.5), which we repeat here for convenience:

$$\max \left\{ 3^{-\frac{1}{4}(1-\beta)l} \Pi, K_{\Psi}^8 \Pi 3^{-\frac{1}{2}(\nu-\gamma)(1-\beta)l}, \frac{K_{\Psi_S}^{8d} \Pi^2}{(1-\gamma)^2} 3^{-l}, \frac{3^{-(1-\gamma)l}}{1-\gamma} \right\} \leq \delta \sigma^2. \quad (3.17)$$

We require also that  $l$  is large enough that  $3^l \mathbb{L}_0 \subseteq \mathbb{Z}^d$ ; in view of (2.100), this can be ensured by taking  $\delta_0$  sufficiently small (depending only on  $d$ ). We emphasize that the parameter  $l$  above depends on  $\delta$  and  $\sigma$ , in addition to the other parameters  $(K_{\Psi}, K_{\Psi_S}, \gamma, \nu, \beta, \Pi)$ , but this dependence is quite explicit.

- We assume that  $m \in \mathbb{N}$  with  $m \geq 100l$  is such that (3.6) holds, that is,

$$\overline{\mathbf{A}}(\square_0) \leq \mathbf{E}_0 \quad \text{and} \quad |\overline{\mathbf{A}}^{-1/2}(\square_m) \mathbf{E}_0 \overline{\mathbf{A}}^{-1/2}(\square_m) - \mathbf{I}_{2d}| \leq \delta \sigma^2. \quad (3.18)$$

To simplify the presentation, throughout this subsection we work with the following notational conventions and assumptions:

- To keep the notation short, for every  $j \in \mathbb{N}$  we define  $\overline{\mathbf{A}}_j := \overline{\mathbf{A}}(\square_j)$ ,  $\overline{\mathbf{s}}_j := \overline{\mathbf{s}}(\square_j)$ ,  $\overline{\mathbf{s}}_{*,j} := \overline{\mathbf{s}}_*(\square_j)$  and  $\overline{\mathbf{k}}_j := \overline{\mathbf{k}}(\square_j)$ . Similarly we define  $\overline{\mathbf{b}}_j := \overline{\mathbf{s}}_j + \overline{\mathbf{k}}_j^t \overline{\mathbf{s}}_{*,j}^{-1} \overline{\mathbf{k}}_j$  and, given a constant matrix  $\mathbf{h}$ , we set  $\overline{\mathbf{b}}_{\mathbf{h},j} := \overline{\mathbf{s}}_j + (\overline{\mathbf{k}}_j - \mathbf{h})^t \overline{\mathbf{s}}_{*,j}^{-1} (\overline{\mathbf{k}}_j - \mathbf{h})$ .
- The coefficient field is “centered” so that the anti-symmetric part of a certain annealed coarse-grained matrix vanishes. By subtracting the matrix  $\frac{1}{2}(\overline{\mathbf{k}}_m - \overline{\mathbf{k}}_m^t)$  from the coefficient field, and recentering both  $\mathbf{E}_0$  and  $\mathbf{A}(U)$  accordingly (as in Section 2.3), we may assume that

$$\overline{\mathbf{k}}_m = \overline{\mathbf{k}}_m^t. \quad (3.19)$$

This centering does not alter  $\Theta$  and it enlarges  $\Pi$  by at most a factor of 100, as shown below. This inflation of  $\Pi$  can be absorbed by making the  $\delta_0$  in the statement of Proposition 3.3 smaller by a factor  $10^{-4}$ .

We stress that (3.19) is assumed to be in force for the rest of the proof of Proposition 3.3.

We next write down some inequalities involving the coarse-grained matrices which are needed throughout this subsection. We first claim that, in view of the above centering condition in (3.19), there exists a constant  $C(d) < \infty$  such that

$$|\mathbf{m}_0^{-1/2} \mathbf{b}_0 \mathbf{m}_0^{-1/2}| + |\mathbf{s}_{*,0}^{-1/2} \mathbf{m}_0 \mathbf{s}_{*,0}^{-1/2}| \leq 13d\Theta^{1/2}. \quad (3.20)$$

Recall that  $\mathbf{m}_0$  is defined to be  $\mathbf{m}_0 := (\mathbf{s}_0 + (\mathbf{k}_0 - \mathbf{h}_0) \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{h}_0)) \# \mathbf{s}_{0,*}$ , where  $\#$  denotes the metric geometric mean and  $\mathbf{h}_0 \in \mathbb{R}_{\text{skew}}^{d \times d}$  minimizes the quantity  $|\mathbf{s}_0 + (\mathbf{k}_0 - \mathbf{h})^t \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{h})|$  over  $\mathbf{h} \in \mathbb{R}_{\text{skew}}^{d \times d}$ . It follows from the definitions of  $\mathbf{m}_0$ ,  $\Theta$  and the geometric mean  $\#$  that

$$|\mathbf{s}_{*,0}^{-1/2} \mathbf{m}_0 \mathbf{s}_{*,0}^{-1/2}| = |\mathbf{s}_{*,0}^{-1/2} (\mathbf{s}_0 + (\mathbf{k}_0 - \mathbf{h}_0) \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{h}_0)) \mathbf{s}_{*,0}^{-1/2}|^{1/2} = \Theta^{1/2}.$$

To prove (3.20), we first observe that (3.18) gives us the bound

$$\mathbf{E}_0 \leq |\overline{\mathbf{A}}_m^{-1/2} \mathbf{E}_0 \overline{\mathbf{A}}_m^{-1/2}| \overline{\mathbf{A}}_m \leq (1 + \delta\sigma^2) \overline{\mathbf{A}}_m \leq (1 + \delta\sigma^2) \overline{\mathbf{A}}_0 \leq (1 + \delta\sigma^2) \mathbf{E}_0 \leq 2\mathbf{E}_0. \quad (3.21)$$

Thus, by (2.51), we have that  $\overline{\mathbf{s}}_{*,m} \leq 2\mathbf{s}_{*,0}$  and  $(\mathbf{k}_0 - \overline{\mathbf{k}}_m)^t \mathbf{s}_{*,m}^{-1} (\mathbf{k}_0 - \overline{\mathbf{k}}_m) \leq 2\mathbf{s}_0$ . It follows that

$$\begin{aligned} |\mathbf{m}_0^{-1/2} (\mathbf{k}_0 - \overline{\mathbf{k}}_m)^t \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \overline{\mathbf{k}}_m) \mathbf{m}_0^{-1/2}| &\leq 2 |\mathbf{m}_0^{-1/2} (\mathbf{k}_0 - \overline{\mathbf{k}}_m)^t \mathbf{s}_{*,m}^{-1} (\mathbf{k}_0 - \overline{\mathbf{k}}_m) \mathbf{m}_0^{-1/2}| \\ &\leq 4 |\mathbf{m}_0^{-1/2} \mathbf{s}_0 \mathbf{m}_0^{-1/2}| \leq 4\Theta^{1/2}. \end{aligned} \quad (3.22)$$

Since  $\overline{\mathbf{k}}_m$  is symmetric by (3.19), we obtain that

$$|\mathbf{m}_0^{-1/2} (\mathbf{k}_0 - \mathbf{k}_0^t) \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{k}_0^t) \mathbf{m}_0^{-1/2}| \leq 4 |\mathbf{m}_0^{-1/2} (\mathbf{k}_0 - \overline{\mathbf{k}}_m)^t \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \overline{\mathbf{k}}_m) \mathbf{m}_0^{-1/2}| \leq 16\Theta^{1/2}.$$

Set  $\tilde{\mathbf{h}}_0 := \frac{1}{2}(\mathbf{k}_0 - \mathbf{k}_0^t)$  and notice that this is the anti-symmetric  $\mathbf{h}$  which minimizes the quantity

$$\min_{\mathbf{h} \in \mathbb{R}_{\text{skew}}^{d \times d}} \text{trace} \left( \mathbf{m}_0^{-1/2} (\mathbf{s}_0 + (\mathbf{k}_0 - \mathbf{h})^t \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{h})) \mathbf{m}_0^{-1/2} \right).$$

Using the previous displays, we therefore obtain

$$\begin{aligned} |\mathbf{m}_0^{-1/2} \mathbf{b}_0 \mathbf{m}_0^{-1/2}| &\leq 4 |\mathbf{m}_0^{-1/2} (\mathbf{s}_0 + (\mathbf{k}_0 - \tilde{\mathbf{h}}_0)^t \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \tilde{\mathbf{h}}_0)^t) \mathbf{m}_0^{-1/2}| + 4 |\mathbf{m}_0^{-1/2} \tilde{\mathbf{h}}_0 \mathbf{s}_{*,0}^{-1} \tilde{\mathbf{h}}_0 \mathbf{m}_0^{-1/2}| \\ &\leq 4 \text{trace} \left( \mathbf{m}_0^{-1/2} (\mathbf{s}_0 + (\mathbf{k}_0 - \tilde{\mathbf{h}}_0)^t \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \tilde{\mathbf{h}}_0)) \mathbf{m}_0^{-1/2} \right) + 16\Theta^{1/2} \\ &= 4 \inf_{\mathbf{h} \in \mathbb{R}_{\text{skew}}^{d \times d}} \text{trace} \left( \mathbf{m}_0^{-1/2} (\mathbf{s}_0 + (\mathbf{k}_0 - \mathbf{h})^t \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{h})) \mathbf{m}_0^{-1/2} \right) + 16\Theta^{1/2} \\ &\leq 4d \inf_{\mathbf{h} \in \mathbb{R}_{\text{skew}}^{d \times d}} |\mathbf{m}_0^{-1/2} (\mathbf{s}_0 + (\mathbf{k}_0 - \mathbf{h})^t \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{h})) \mathbf{m}_0^{-1/2}| + 16\Theta^{1/2} \\ &= (4d + 16)\Theta^{1/2}. \end{aligned}$$

This proves the claim (3.20). By a similar argument, putting  $\mathbf{m}_0 = \mathbf{I}_d$  in the above computations, yields that the centering enlarges the aspect ratio  $\Pi$  by at most a factor of 100. Indeed, we obtain

$$|(\mathbf{k}_0 - \mathbf{k}_0^t) \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{k}_0^t)| \leq 24|\mathbf{s}_0|$$

and thus, using also (3.21),

$$\Pi = |\mathbf{E}_{11}| |\mathbf{E}_{22}^{-1}| \leq 4 |\mathbf{s}_0 + (\mathbf{k}_0 - \mathbf{k}_0^t) \mathbf{s}_{*,0}^{-1} (\mathbf{k}_0 - \mathbf{k}_0^t)| |\mathbf{s}_{*,0}^{-1}| \leq 100 |\mathbf{s}_0| |\mathbf{s}_{*,0}^{-1}| \leq 100 \Pi_{\text{old}}, \quad (3.23)$$

where  $\Pi_{\text{old}}$  is the constant  $\Pi$  before we centered according to (3.19).

We continue by transferring the assumed bounds (3.18) from Euclidean cubes to the adapted cubes and, using the mixing condition, obtain an estimate on the variance of  $\mathbf{A}(\diamond_n)$  across roughly the same range of scales. We note that  $x \mapsto \mathbf{A}(x + \diamond_n)$  is  $\mathbb{Z}^d$ -stationary for every  $n \in \mathbb{N}$  with  $n \geq l$ , since  $3^l \mathbb{L}_0 \subseteq \mathbb{Z}^d$ .

**Lemma 3.5.** *There exists a constant  $C(d) < \infty$  such that*

$$\max_{n \in \mathbb{N} \cap [l, m-l]} |\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}(\diamond_n) \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}| \leq C\delta\sigma^2 \quad (3.24)$$

and

$$\max_{n \in \mathbb{N} \cap [l, m-l]} \mathbb{E} \left[ |\overline{\mathbf{A}}_m^{-1/2} \mathbf{A}(\diamond_n) \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}|^2 \right] \leq C\delta\sigma^2. \quad (3.25)$$

*Proof.* Define  $h := \lceil \frac{1}{4}(1 - \beta)l \rceil$ . By (3.17) we have that each of the following conditions are valid:

$$\frac{K_{\Psi}^8 \Pi^2}{(1 - \gamma)^2} 3^{-l} \leq \delta \sigma^2, \quad K_{\Psi}^8 \Pi 3^{-\frac{1}{2}(\nu - \gamma)(1 - \beta)l} \leq \delta \sigma^2 \quad \text{and} \quad 3^{-\frac{1}{4}(1 - \beta)l} \Pi \leq \delta \sigma^2. \quad (3.26)$$

*Step 1.* The proof of (3.24). We will show that

$$\max_{n \in \mathbb{N} \cap [h, m - h]} |\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}(\diamond_n) \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}| \leq C \delta \sigma^2. \quad (3.27)$$

This inequality is a consequence of (3.18) and Lemma 2.10. The latter implies that there exists a constant  $C(d) < \infty$  such that, for every  $h, n, m \in \mathbb{N}$  with  $h \leq n \leq m - h$ ,

$$\begin{aligned} \mathbf{I}_{2d} &= \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_m \overline{\mathbf{A}}_m^{-1/2} \leq \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}(\diamond_n) \overline{\mathbf{A}}_m^{-1/2} + C 3^{n-m} \overline{\mathbf{A}}_m^{-1/2} \mathbf{E}_0 \overline{\mathbf{A}}_m^{-1/2} \\ &\leq \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_0 \overline{\mathbf{A}}_m^{-1/2} + C \Pi (3^{n-m} + 3^{-n}) \overline{\mathbf{A}}_m^{-1/2} \mathbf{E}_0 \overline{\mathbf{A}}_m^{-1/2} \\ &\leq (1 + C \Pi 3^{-h}) (1 + \delta \sigma^2) \mathbf{I}_{2d} \end{aligned}$$

This string of inequalities implies (3.24) provided that  $3^h \geq 2C(\delta \sigma^2)^{-1} \Pi$ , which is guaranteed by the choice  $h := \lceil \frac{1}{4}(1 - \beta)l \rceil$  and (3.17).

*Step 2.* We next show that,<sup>6</sup> for every  $m, n, k \in \mathbb{N}$  with  $k \leq n$ , we have that

$$\begin{aligned} &\mathbb{E} \left[ \left| \overline{\mathbf{A}}^{-1/2}(\square_m) \mathbf{A}(\diamond_n) \overline{\mathbf{A}}^{-1/2}(\square_m) - \mathbf{I}_{2d} \right|^2 \right] \\ &\leq 40d \max_{j \in \{k, n\}} \left| \overline{\mathbf{A}}^{-1/2}(\square_m) \overline{\mathbf{A}}^{-1/2}(\square_m) - \mathbf{I}_{2d} \right| + 27 \left| \overline{\mathbf{A}}^{-1/2}(\square_m) \overline{\mathbf{A}}(\diamond_k) \overline{\mathbf{A}}^{-1/2}(\square_m) - \mathbf{I}_{2d} \right|^2 \\ &\quad + 27 \mathbb{E} \left[ \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \overline{\mathbf{A}}^{-1/2}(\square_m) (\mathbf{A}(z + \diamond_k) - \overline{\mathbf{A}}(z + \diamond_k)) \overline{\mathbf{A}}^{-1/2}(\square_m) \right|^2 \right]. \end{aligned} \quad (3.28)$$

By the triangle inequality, we get that

$$\begin{aligned} &|\overline{\mathbf{A}}_m^{-1/2} \mathbf{A}(\diamond_n) \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}|^2 \\ &\leq 3 \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \overline{\mathbf{A}}_m^{-1/2} (\mathbf{A}(z + \diamond_k) - \overline{\mathbf{A}}(\diamond_k)) \overline{\mathbf{A}}_m^{-1/2} \right|^2 + 3 \left| \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}(\diamond_k) \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d} \right|^2 \\ &\quad + 3 \left| \overline{\mathbf{A}}_m^{-1/2} (\mathbf{A}(\diamond_n) - \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbf{A}(z + \diamond_k)) \overline{\mathbf{A}}_m^{-1/2} \right|^2. \end{aligned} \quad (3.29)$$

The first term and the second term appear on the right side of (3.28). The last term on the right side of (3.29) is the square of the additivity defect, written in terms of  $\mathbf{A}$ . By subadditivity, we also have that

$$\overline{\mathbf{A}}_m^{-1/2} \left( \mathbf{A}(\diamond_n) - \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbf{A}(z + \diamond_k) \right) \overline{\mathbf{A}}_m^{-1/2} \leq 0. \quad (3.30)$$

---

<sup>6</sup>The statement in Step 2 is valid without the assumptions of Proposition 3.3, as the proof does not use them. It also does not use the recentering assumption (3.19). We will need these observations in Section 5 when we need to reuse the arguments here.

We therefore obtain, using also (3.27),

$$\begin{aligned}
& \mathbb{E} \left[ \left| \overline{\mathbf{A}}_m^{-1/2} \left( \mathbf{A}(\diamond_n) - \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbf{A}(z + \diamond_k) \right) \overline{\mathbf{A}}_m^{-1/2} \right| \right] \\
& \leq 2d \left| \mathbb{E} \left[ \overline{\mathbf{A}}_m^{-1/2} \left( \mathbf{A}(\diamond_n) - \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbf{A}(z + \diamond_k) \right) \overline{\mathbf{A}}_m^{-1/2} \right] \right| \\
& \leq 2d \left| \overline{\mathbf{A}}_m^{-1/2} (\overline{\mathbf{A}}(\diamond_k) - \overline{\mathbf{A}}(\diamond_n)) \overline{\mathbf{A}}_m^{-1/2} \right| \\
& \leq 4d \max_{j \in \{k, n\}} \left| \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}(\diamond_j) \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d} \right|.
\end{aligned}$$

We use (3.30) to (crudely) estimate one factor from above: we have

$$\begin{aligned}
& \left| \overline{\mathbf{A}}_m^{-1/2} \left( \mathbf{A}(\diamond_n) - \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbf{A}(z + \diamond_k) \right) \overline{\mathbf{A}}_m^{-1/2} \right| \\
& \leq \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \overline{\mathbf{A}}_m^{-1/2} \mathbf{A}(z + \diamond_k) \overline{\mathbf{A}}_m^{-1/2} \right| \\
& \leq 1 + \left| \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}(\diamond_k) \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d} \right| + \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \overline{\mathbf{A}}_m^{-1/2} (\mathbf{A}(z + \diamond_k) - \overline{\mathbf{A}}(z + \diamond_k)) \overline{\mathbf{A}}_m^{-1/2} \right|.
\end{aligned}$$

For any square integrable, nonnegative random variable  $X$  we have  $\mathbb{E}[X^2] \leq 2\mathbb{E}[X] + 4\mathbb{E}[(X-1)_+^2]$ . Therefore, by the previous two displays,

$$\begin{aligned}
& \mathbb{E} \left[ \left| \overline{\mathbf{A}}_m^{-1/2} \left( \mathbf{A}(\diamond_n) - \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbf{A}(z + \diamond_k) \right) \overline{\mathbf{A}}_m^{-1/2} \right|^2 \right] \\
& \leq 8d \max_{j \in \{k, n\}} \left| \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}(\diamond_j) \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d} \right| + 8 \left| \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}(\diamond_k) \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d} \right|^2 \\
& \quad + 8 \mathbb{E} \left[ \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \overline{\mathbf{A}}_m^{-1/2} (\mathbf{A}(z + \diamond_k) - \overline{\mathbf{A}}(z + \diamond_k)) \overline{\mathbf{A}}_m^{-1/2} \right|^2 \right].
\end{aligned}$$

Combining this with (3.29) yields (3.28).

*Step 3.* The proof of (3.25). Let  $n \in \mathbb{N}$  with  $l \leq n \leq m - l$  and let  $k := n - h$ . We then have that  $k - \beta n = (1 - \beta)n - h \geq (1 - \beta)l - h > 0$  and  $k \geq h$ . Appealing to Lemma 2.9, we obtain that,

$$\begin{aligned}
& \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbf{E}_0^{-1/2} (\mathbf{A}(z + \diamond_k) - \overline{\mathbf{A}}(z + \diamond_k)) \mathbf{E}_0^{-1/2} \right| \\
& \leq \frac{CK_{\Psi_S}^2 \Pi}{1 - \gamma} 3^{\gamma h - l} + \mathcal{O}_{\Psi_S} \left( \frac{C\Pi}{1 - \gamma} 3^{\gamma h - l} \right) + \mathcal{O}_{\Psi} \left( C\Pi^{1/2} 3^{-(\nu - \gamma)h} \right). \quad (3.31)
\end{aligned}$$

Since  $h := \lceil \frac{1}{4}(1 - \beta)l \rceil$ , it follows by (3.17) that

$$\begin{aligned}
& \mathbb{E} \left[ \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \overline{\mathbf{A}}_m^{-1/2} (\mathbf{A}(z + \diamond_k) - \overline{\mathbf{A}}(\diamond_k)) \overline{\mathbf{A}}_m^{-1/2} \right|^2 \right] \\
& \leq \frac{CK_{\Psi_S}^8 \Pi^2 3^{-l}}{(1 - \gamma)^2} + CK_{\Psi}^8 \Pi 3^{-\frac{1}{2}(\nu - \gamma)(1 - \beta)l} \leq C\delta\sigma^2. \quad (3.32)
\end{aligned}$$

Since  $k \geq h$ , the expectation of the third term on the right side of (3.28) is controlled by (3.27):

$$\max_{j \in \{k, n\}} |\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}(\diamond_j) \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}| \leq C \delta^2 \sigma^8 \leq C \delta \sigma^2. \quad (3.33)$$

Therefore, putting together (3.32), (3.33) and (3.28) completes the proof of the lemma.  $\square$

Let us recall the definition of the renormalized ellipticity ratio at scale  $3^m$ :

$$\Theta_m := \inf_{\mathbf{h} \in \mathbb{R}_{\text{skew}}^{d \times d}} |\overline{\mathbf{s}}_{*,m}^{-1/2} (\overline{\mathbf{s}}_m + (\overline{\mathbf{k}}_m - \mathbf{h}) \overline{\mathbf{s}}_{*,m}^{-1} (\overline{\mathbf{k}}_m - \mathbf{h})) \overline{\mathbf{s}}_{*,m}^{-1/2}|. \quad (3.34)$$

Since  $\overline{\mathbf{k}}_m$  is symmetric, we have that

$$\begin{aligned} \Theta_m - 1 &\leq |\overline{\mathbf{s}}_{*,m}^{-1/2} \overline{\mathbf{b}}_m \overline{\mathbf{s}}_{*,m}^{-1/2} - \mathbf{I}_d| \\ &\leq \text{trace}(\overline{\mathbf{s}}_{*,m}^{-1/2} \overline{\mathbf{b}}_m \overline{\mathbf{s}}_{*,m}^{-1/2} - \mathbf{I}_d) \\ &= \inf_{\mathbf{h} \in \mathbb{R}_{\text{skew}}^{d \times d}} \text{trace}(\overline{\mathbf{s}}_{*,m}^{-1/2} (\overline{\mathbf{s}}_m + (\overline{\mathbf{k}}_m - \mathbf{h}) \overline{\mathbf{s}}_{*,m}^{-1} (\overline{\mathbf{k}}_m - \mathbf{h})) \overline{\mathbf{s}}_{*,m}^{-1/2} - \mathbf{I}_d) \leq d(\Theta_m - 1). \end{aligned} \quad (3.35)$$

Our next goal is to relate  $\Theta_m - 1$  to the variational quantities  $J$  and  $J^*$  and their maximizers. It is convenient to introduce the following variant of  $J$ :

$$\tilde{J}(U, p, q) := J(U, p, q) - \frac{1}{2} \mathbb{E} \left[ \int_U \nabla v(\cdot, U, p, q) \right] \cdot \mathbb{E} \left[ \int_U \mathbf{a} \nabla v(\cdot, U, p, q) \right]. \quad (3.36)$$

This “centers” the quantity  $J$  by removing the part of the energy due to the “bias” in the spatial averages of the gradient and flux of its maximizer  $v(\cdot, U, p, q)$ .<sup>7</sup> We let  $\tilde{J}^*(U, p, q)$  denote the analogous quantity defined for the adjoint coefficient field  $\mathbf{a}^t$ .

We also introduce the following matrices: first, the anti-symmetric part of  $\overline{\mathbf{k}}(U)$  is denoted by

$$\overline{\mathbf{h}}(U) := \frac{1}{2} (\overline{\mathbf{k}} - \overline{\mathbf{k}}^t)(U), \quad (3.37)$$

and the geometric mean of  $\overline{\mathbf{b}}_{\overline{\mathbf{h}}(U)}(U)$  and  $\overline{\mathbf{s}}_*(U)$  is given by

$$\overline{\mathbf{t}}(U) := (\overline{\mathbf{s}} + (\overline{\mathbf{k}} - \overline{\mathbf{h}})^t \overline{\mathbf{s}}_*^{-1} (\overline{\mathbf{k}} - \overline{\mathbf{h}}))(U) \# \overline{\mathbf{s}}_*(U) = (\overline{\mathbf{b}}_{\overline{\mathbf{h}}(U)} \# \overline{\mathbf{s}}_*)(U). \quad (3.38)$$

Note that our centering assumption (3.19) implies that  $\overline{\mathbf{h}}(\square_m) = 0$ , so that

$$\overline{\mathbf{t}}_m := \overline{\mathbf{b}}_m \# \overline{\mathbf{s}}_{*,m} = (\overline{\mathbf{s}}(\square_m) + \overline{\mathbf{k}}(\square_m) \overline{\mathbf{s}}_*^{-1}(\square_m) \overline{\mathbf{k}}(\square_m)) \# \overline{\mathbf{s}}_*(\square_m). \quad (3.39)$$

In the next lemma we control the ratio of  $\overline{\mathbf{t}}(U)$  to  $\overline{\mathbf{s}}_*(U)$  by an expression involving the expectations of  $\tilde{J}(U, p, q)$  and  $\tilde{J}^*(p, q')$  for particular choices of  $p, q, q'$ ; note that, in view of (3.35) and (3.39), this will allow us to control  $\Theta_m - 1$  in terms of the latter for  $U = \square_m$ .

**Lemma 3.6.** *For every bounded Lipschitz domain  $U \subseteq \mathbb{R}^d$ ,*

$$\begin{aligned} |(\overline{\mathbf{s}}_*^{-1/2} \overline{\mathbf{b}}_{\overline{\mathbf{h}}(U)} \overline{\mathbf{s}}_*^{-1/2})(U) - \mathbf{I}_d| &\leq 2d \sup_{|e|=1} \left( \mathbb{E} \left[ \tilde{J}(U, \overline{\mathbf{t}}^{-1/2}(U)e, \overline{\mathbf{t}}^{1/2}(U)e - \overline{\mathbf{h}}(U) \overline{\mathbf{t}}^{-1/2}(U)e) \right] \right. \\ &\quad \left. + \mathbb{E} \left[ \tilde{J}^*(U, \overline{\mathbf{t}}^{-1/2}(U)e, \overline{\mathbf{t}}^{1/2}(U)e + \overline{\mathbf{h}}(U) \overline{\mathbf{t}}^{-1/2}(U)e) \right] \right). \end{aligned} \quad (3.40)$$

<sup>7</sup>This bias does not usually appear in the theory, because a choice of the parameters  $p, q$  is typically made so that one of the factors vanishes. In the high contrast setting, this would create additional error terms which are too large. Here we must be more careful in our choice of these parameters  $p, q$  so as to balance various error terms.

*Proof.* The lemma is a consequence of the following claim: For every  $p, q, h \in \mathbb{R}^d$ ,

$$\begin{aligned} & \mathbb{E}[\tilde{J}(U, p, q - h) + \tilde{J}^*(U, p, q + h)] \\ &= q \cdot (\bar{\mathbf{s}}(U) \bar{\mathbf{s}}_*^{-1}(U) - \mathbf{I}_d) p + \bar{\mathbf{s}}_*^{-1}(U) q \cdot (\bar{\mathbf{k}}(U) + \bar{\mathbf{k}}^t(U)) \bar{\mathbf{s}}_*^{-1}(U) (\bar{\mathbf{k}}(U) p - h). \end{aligned} \quad (3.41)$$

Indeed, we obtain (3.40) from (3.41) by using the definitions of  $\bar{\mathbf{t}}(U)$  and  $\bar{\mathbf{h}}(U)$  in (3.38) and (3.37), respectively, and the following computation (all of the matrices are evaluated at  $U$ ):

$$\begin{aligned} & \sum_{j=1}^d \mathbb{E} \left[ \tilde{J}(U, \bar{\mathbf{t}}^{-1/2} e_j, \bar{\mathbf{t}}^{1/2} e_j - \bar{\mathbf{h}} \bar{\mathbf{t}}^{-1/2} e_j) + \tilde{J}^*(U, \bar{\mathbf{t}}^{-1/2} e_j, \bar{\mathbf{t}}^{1/2} e_j + \bar{\mathbf{h}} \bar{\mathbf{t}}^{-1/2} e_j) \right] \\ &= \sum_{j=1}^d \left( e_j \cdot \bar{\mathbf{t}}^{1/2} (\bar{\mathbf{s}} \bar{\mathbf{s}}_*^{-1} - \mathbf{I}_d) \bar{\mathbf{t}}^{-1/2} e_j + \frac{1}{2} e_j \cdot \bar{\mathbf{t}}^{1/2} \bar{\mathbf{s}}_*^{-1} (\bar{\mathbf{k}} + \bar{\mathbf{k}}^t) \bar{\mathbf{s}}_*^{-1} (\bar{\mathbf{k}} + \bar{\mathbf{k}}^t) \bar{\mathbf{t}}^{-1/2} e_j \right) \\ &= \text{trace} \left( \bar{\mathbf{t}}^{1/2} (\bar{\mathbf{s}} \bar{\mathbf{s}}_*^{-1} - \mathbf{I}_d) \bar{\mathbf{t}}^{-1/2} \right) + \frac{1}{2} \text{trace} \left( \bar{\mathbf{t}}^{1/2} \bar{\mathbf{s}}_*^{-1} (\bar{\mathbf{k}} + \bar{\mathbf{k}}^t) \bar{\mathbf{s}}_*^{-1} (\bar{\mathbf{k}} + \bar{\mathbf{k}}^t) \bar{\mathbf{t}}^{-1/2} \right) \\ &= \text{trace} \left( \bar{\mathbf{s}}_*^{-1/2} \bar{\mathbf{s}} \bar{\mathbf{s}}_*^{-1/2} - \mathbf{I}_d \right) + \frac{1}{2} \text{trace} \left( \bar{\mathbf{s}}_*^{-1/2} (\bar{\mathbf{k}} - \bar{\mathbf{h}})^t \bar{\mathbf{s}}_*^{-1} (\bar{\mathbf{k}} - \bar{\mathbf{h}}) \bar{\mathbf{s}}_*^{-1/2} \right) \\ &\geq \frac{1}{2} \text{trace} \left( \bar{\mathbf{s}}_*^{-1/2} (\bar{\mathbf{s}} + (\bar{\mathbf{k}} - \bar{\mathbf{h}})^t \bar{\mathbf{s}}_*^{-1} (\bar{\mathbf{k}} - \bar{\mathbf{h}})) \bar{\mathbf{s}}_*^{-1/2} - \mathbf{I}_d \right) \\ &\geq \frac{1}{2} \left| \bar{\mathbf{s}}_*^{-1/2} (\bar{\mathbf{s}} + (\bar{\mathbf{k}} - \bar{\mathbf{h}})^t \bar{\mathbf{s}}_*^{-1} (\bar{\mathbf{k}} - \bar{\mathbf{h}})) \bar{\mathbf{s}}_*^{-1/2} - \mathbf{I}_d \right|, \end{aligned} \quad (3.42)$$

where the cyclic property of traces was applied to obtain the last equality and the non-negativity of the matrices inside of the traces to obtain the last inequalities.

We turn to the proof of (3.41). We suppress  $U$  from the notation, since it plays no role in the argument. Recall the definition of  $\mathbf{R}$  in (2.32) and the formula for  $\bar{\mathbf{A}}_*^{-1}$  in (2.17). Notice in particular that  $\mathbf{R} \bar{\mathbf{A}} \mathbf{R} = \bar{\mathbf{A}}_*^{-1}$  and  $\mathbf{R}^2 = \mathbf{I}_{2d}$ . The formula in (2.33) reads as

$$\mathbb{E} \left[ \oint \left( \begin{array}{c} \nabla v(\cdot, p, q - h) \\ \mathbf{a} \nabla v(\cdot, p, q - h) \end{array} \right) \right] = (\mathbf{R} \bar{\mathbf{A}} + \mathbf{I}_{2d}) \begin{pmatrix} -p \\ q - h \end{pmatrix}.$$

For the adjoint maximizers, the corresponding formula is

$$\mathbb{E} \left[ \oint \left( \begin{array}{c} \nabla v^*(\cdot, p, q + h) \\ -\mathbf{a}^t \nabla v^*(\cdot, p, q + h) \end{array} \right) \right] = (\mathbf{R} \bar{\mathbf{A}} - \mathbf{I}_{2d}) \begin{pmatrix} p \\ q + h \end{pmatrix}.$$

Using  $\mathbf{R}^2 = \mathbf{I}_{2d}$  and (2.13), we find that

$$\begin{aligned} \frac{1}{2} \mathbb{E} \left[ \oint \nabla v(\cdot, p, q - h) \right] \cdot \mathbb{E} \left[ \oint \mathbf{a} \nabla v(\cdot, p, q - h) \right] &= \frac{1}{4} \begin{pmatrix} -p \\ q - h \end{pmatrix} \cdot (\bar{\mathbf{A}} + \mathbf{R}) \mathbf{R} (\bar{\mathbf{A}} + \mathbf{R}) \begin{pmatrix} -p \\ q - h \end{pmatrix} \\ &= \mathbb{E}[J(p, q - h)] + \frac{1}{4} \begin{pmatrix} -p \\ q - h \end{pmatrix} \cdot (\bar{\mathbf{A}} \mathbf{R} \bar{\mathbf{A}} - \mathbf{R}) \begin{pmatrix} -p \\ q - h \end{pmatrix} \end{aligned}$$

and, similarly,

$$\begin{aligned} \frac{1}{2} \mathbb{E} \left[ \oint \nabla v^*(\cdot, p, q + h) \right] \cdot \mathbb{E} \left[ \oint \mathbf{a}^t \nabla v^*(\cdot, p, q + h) \right] &= -\frac{1}{4} \begin{pmatrix} p \\ q + h \end{pmatrix} \cdot (\bar{\mathbf{A}} \mathbf{R} - \mathbf{I}_{2d}) (\bar{\mathbf{A}} - \mathbf{R}) \begin{pmatrix} p \\ q + h \end{pmatrix} \\ &= \mathbb{E}[J^*(p, q + h)] - \frac{1}{4} \begin{pmatrix} p \\ q + h \end{pmatrix} \cdot (\bar{\mathbf{A}} \mathbf{R} \bar{\mathbf{A}} - \mathbf{R}) \begin{pmatrix} p \\ q + h \end{pmatrix}. \end{aligned}$$

Summing these up yields that

$$\begin{aligned}\mathbb{E}[\tilde{J}(U, p, q-h) + \tilde{J}^*(U, p, q+h)] &= \frac{1}{2} \binom{p}{h} \cdot (\overline{\mathbf{A}}\mathbf{R}\overline{\mathbf{A}} - \mathbf{R}) \binom{0}{q} + \frac{1}{2} \binom{0}{q} \cdot (\overline{\mathbf{A}}\mathbf{R}\overline{\mathbf{A}} - \mathbf{R}) \binom{p}{h} \\ &= \binom{p}{h} \cdot (\overline{\mathbf{A}}\mathbf{R}\overline{\mathbf{A}} - \mathbf{R}) \binom{0}{q}.\end{aligned}$$

We next compute

$$\overline{\mathbf{A}} \binom{p}{h} = \begin{pmatrix} \bar{\mathbf{b}}p - \bar{\mathbf{k}}^t \bar{\mathbf{s}}_*^{-1} h \\ -\bar{\mathbf{s}}_*^{-1} \bar{\mathbf{k}}p + \bar{\mathbf{s}}_*^{-1} h \end{pmatrix} \quad \text{and} \quad \mathbf{R}\overline{\mathbf{A}} \binom{0}{q} = \mathbf{R} \begin{pmatrix} -\bar{\mathbf{k}}^t \bar{\mathbf{s}}_*^{-1} q \\ \bar{\mathbf{s}}_*^{-1} q \end{pmatrix} = \begin{pmatrix} \bar{\mathbf{s}}_*^{-1} q \\ -\bar{\mathbf{k}}^t \bar{\mathbf{s}}_*^{-1} q \end{pmatrix}.$$

It follows that

$$\begin{aligned}\binom{p}{h} \cdot \overline{\mathbf{A}}\mathbf{R}\overline{\mathbf{A}} \binom{0}{q} &= \begin{pmatrix} \bar{\mathbf{b}}p - \bar{\mathbf{k}}^t \bar{\mathbf{s}}_*^{-1} h \\ -\bar{\mathbf{s}}_*^{-1} \bar{\mathbf{k}}p + \bar{\mathbf{s}}_*^{-1} h \end{pmatrix} \cdot \begin{pmatrix} \bar{\mathbf{s}}_*^{-1} q \\ -\bar{\mathbf{k}}^t \bar{\mathbf{s}}_*^{-1} q \end{pmatrix} \\ &= p \cdot \bar{\mathbf{b}} \bar{\mathbf{s}}_*^{-1} q - h \cdot \bar{\mathbf{s}}_*^{-1} \bar{\mathbf{k}} \bar{\mathbf{s}}_*^{-1} q + p \cdot \bar{\mathbf{k}}^t \bar{\mathbf{s}}_*^{-1} \bar{\mathbf{k}}^t \bar{\mathbf{s}}_*^{-1} q - h \cdot \bar{\mathbf{s}}_*^{-1} \bar{\mathbf{k}}^t \bar{\mathbf{s}}_*^{-1} q \\ &= p \cdot \bar{\mathbf{s}} \bar{\mathbf{s}}_*^{-1} q - h \cdot \bar{\mathbf{s}}_*^{-1} (\bar{\mathbf{k}} + \bar{\mathbf{k}}^t) \bar{\mathbf{s}}_*^{-1} q + p \cdot \bar{\mathbf{k}}^t \bar{\mathbf{s}}_*^{-1} (\bar{\mathbf{k}} + \bar{\mathbf{k}}^t) \bar{\mathbf{s}}_*^{-1} q.\end{aligned}$$

Combining the above displays yields

$$\mathbb{E}[\tilde{J}(U, p, q-h) + \tilde{J}^*(U, p, q+h)] = p \cdot (\bar{\mathbf{s}} \bar{\mathbf{s}}_*^{-1} - \mathbf{I}_d) q + (\bar{\mathbf{k}}p - h) \cdot \bar{\mathbf{s}}_*^{-1} (\bar{\mathbf{k}} + \bar{\mathbf{k}}^t) \bar{\mathbf{s}}_*^{-1} q.$$

This completes the proof of (3.41) and thus of the lemma.  $\square$

Motivated by (3.35) and Lemma 3.6 (recall that  $\bar{\mathbf{h}}(\square_m) = 0$  by (3.19) and (3.37)), our goal is now to get upper bound estimates on

$$\mathbb{E}[\tilde{J}(\square_m, p, q) + \tilde{J}^*(\square_m, p, q)], \quad (3.43)$$

with the choices

$$p := \bar{\mathbf{t}}_m^{-1/2} e \quad \text{and} \quad q := \bar{\mathbf{t}}_m^{1/2} e, \quad (3.44)$$

where  $\bar{\mathbf{t}}_m$  is defined by (3.39) and  $e \in \mathbb{R}^d$  with  $|e| = 1$  is the unit vector attaining the supremum on the right side (3.40) with  $U = \square_m$ : that is,

$$\mathbb{E}[\tilde{J}(\square_m, p, q) + \tilde{J}^*(\square_m, p, q)] = \sup_{|e'|=1} \mathbb{E}[\tilde{J}(\square_m, \bar{\mathbf{t}}_m^{-1/2} e', \bar{\mathbf{t}}_m^{1/2} e') + \tilde{J}^*(\square_m, \bar{\mathbf{t}}_m^{-1/2} e', \bar{\mathbf{t}}_m^{1/2} e')]. \quad (3.45)$$

Note that, with these choices of  $(p, q)$  in (3.44), the combination of (3.35) and (3.40) implies

$$\Theta_m - 1 \leq 2d \mathbb{E}[\tilde{J}(\square_m, p, q) + \tilde{J}^*(\square_m, p, q)]. \quad (3.46)$$

To get a bound on the right side of (3.46), we first switch to the adapted cubes at the cost of giving up a few scales: using (2.108), we have that, for  $n = m - l$ ,

$$\begin{aligned}\mathbb{E}[\tilde{J}(\square_m, p, q) + \tilde{J}^*(\square_m, p, q)] &= \mathbb{E}[J(\square_m, p, q) + J^*(\square_m, p, q)] - \frac{1}{2} P \cdot Q - \frac{1}{2} P^* \cdot Q^* \\ &\leq \mathbb{E}[J(\diamond_n, p, q) + J^*(\diamond_n, p, q)] - \frac{1}{2} P \cdot Q - \frac{1}{2} P^* \cdot Q^* + C \Pi^2 3^{-l},\end{aligned} \quad (3.47)$$

where we let  $P, Q, P^*, Q^* \in \mathbb{R}^d$  denote the vectors

$$\begin{pmatrix} P \\ Q \end{pmatrix} := \mathbb{E} \left[ \int_{\square_m} \begin{pmatrix} \nabla v(\cdot, \square_m, p, q) \\ \mathbf{a} \nabla v(\cdot, \square_m, p, q) \end{pmatrix} \right] \quad \text{and} \quad \begin{pmatrix} P^* \\ Q^* \end{pmatrix} := \mathbb{E} \left[ \int_{\square_m} \begin{pmatrix} \nabla v^*(\cdot, \square_m, p, q) \\ \mathbf{a}^t \nabla v^*(\cdot, \square_m, p, q) \end{pmatrix} \right]. \quad (3.48)$$

The last equality is valid by (2.33). Therefore what we want to bound is the quantity

$$\mathbb{E}[J(\diamond_n, p, q)] - \frac{1}{2} P \cdot Q, \quad (3.49)$$

where  $p$  and  $q$  are as in (3.44),  $e$  is chosen so that (3.45) holds,  $P$  and  $Q$  are defined in (3.48), and  $n \in \mathbb{N}$  is some suitably chosen parameter with  $n < m$ . The analogous bound for  $J^*$  is a consequence of the one for  $J$ , if we apply it with the random field  $\mathbf{a}^t$  in place of  $\mathbf{a}$ .

The idea now is to write the quantity in (3.49) as the integral of the product of the centered gradient and centered flux of their maximizers:

$$\begin{aligned} \mathbb{E}[J(\diamond_n, p, q)] - \frac{1}{2} P \cdot Q &= \frac{1}{2} \begin{pmatrix} Q \\ P \end{pmatrix} \cdot \mathbb{E} \left[ \begin{pmatrix} \nabla v(\cdot, \diamond_n, p, q) - P \\ \mathbf{a} \nabla v(\cdot, \diamond_n, p, q) - Q \end{pmatrix} \right] \\ &\quad + \frac{1}{2} \mathbb{E} \left[ \int_{y+\diamond_n} (\nabla v(\cdot, \diamond_n, p, q) - P) \cdot (\mathbf{a} \nabla v(\cdot, \diamond_n, p, q) - Q) \right]. \end{aligned} \quad (3.50)$$

Note the formula (3.50) is valid for any  $P$  and  $Q$  and does not use the particular choice in (3.48).

The way one would normally proceed (in the uniformly elliptic case) with estimating the right side of (3.50) is to use the additivity defect to estimate the expectation on the first line, since the spatial averages of gradients and fluxes can be expressed in terms of the coarse-grained matrices themselves. The second line is then typically estimated using the Caccioppoli inequality, which reduces the energy term to an  $L^2$  type oscillation of the maximizer itself, which can then be reduced once again to the spatial averages of the gradient and thus the additivity defect.

The problem with this strategy in our context is that the Caccioppoli inequality produces estimates with factors of the (pointwise) ellipticity constants of the microscopic matrix  $\mathbf{a}(x)$ . Since we do not assume that  $\mathbf{a}(\cdot)$  is uniformly elliptic, this strategy is unavailable; even if we did make a uniform ellipticity assumption, this estimate would produce factors of the ellipticity ratio that would result in a much worse estimate than the one we will prove.

We instead proceed in a more *coarse-grained* fashion by directly relating the energy of the maximizer to the weak Sobolev norms of the gradient and flux—while paying only the price required by the *coarse-grained* ellipticity ratio.

This is the content of the next lemma, in which we control the left side of (3.50) by using a similar identity to (3.50), but with a cutoff function smuggled in, and then a quantitative “div-curl” type argument<sup>8</sup> to control the energy term.

**Lemma 3.7.** *There exists a constant  $C(d) < \infty$  such that, for  $n = m - l$ , we have the estimate*

$$\begin{aligned} &\left| \mathbb{E}[J(\diamond_n, p, q)] - \frac{1}{2} P \cdot Q \right| \\ &\leq C 3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \begin{pmatrix} \nabla v(\cdot, \diamond_n, p, q) - P \\ \mathbf{a} \nabla v(\cdot, \diamond_n, p, q) - Q \end{pmatrix} \right]_{\dot{B}_{2,1}^{-1/2}(\diamond_n)}^2 \right] + C \delta^{1/2} \sigma \Theta^{1/2} \Theta_m^{1/2}. \end{aligned} \quad (3.51)$$

---

<sup>8</sup>Recall that the “div-curl lemma” is a classical statement in homogenization that says that products of weakly converging gradients and divergence-free fields also weakly converge.



*Proof.* We first prove a preliminary statement which is valid for general  $P, Q \in \mathbb{R}^d$  (not only for the specific choices made in (3.48)), and also for a general  $\mathbf{q}_0$  and general  $\mathbf{M}_0$  with a defining property that  $3^{k_0}\mathbb{L}_0 \subseteq \mathbb{Z}^d$  and  $\mathbf{q}_0 = \text{const} \cdot \mathbf{m}_0^{1/2}$  (not only for the specific choices made in (2.93) and (2.94)). We then conclude in Step 6 below using the particular definitions (3.48), (2.93) and (2.94).

To make the notation simple, we drop  $p, q$  by denoting, for every  $z \in \mathbb{Z}^d$  and  $k, n \in \mathbb{N}$  with  $n > k + 3 > k \geq k_0$ ,

$$v_{z,k} := v(\cdot, z + \diamond_k, p, q), \quad J(z + \diamond_k) = J(z + \diamond_k, p, q), \quad \bar{\tau}_{n,k} := \mathbb{E}[J(\diamond_k) - J(\diamond_n)].$$

Denote also  $v_n := v(\cdot, \diamond_n, p, q)$ . We fix a nonnegative smooth test function  $\varphi \in C_c^\infty(\diamond_n)$  such that

$$(\varphi)_{\diamond_n} = 1, \quad 0 \leq \varphi \leq 2 \quad \text{and} \quad \|\mathbf{q}_0^j \nabla^j \varphi\|_{L^\infty(\diamond_n)} \leq 3^{-j(n-2)}, \quad j \in \{1, 2\}.$$

Observe that

$$\begin{aligned} \mathbb{E}[J(\diamond_n)] - \frac{1}{2}P \cdot Q &= \mathbb{E}\left[\int_{\diamond_n} \frac{1}{2}\varphi(\nabla v_n - P) \cdot (\mathbf{a}\nabla v_n - Q)\right] \\ &\quad + \mathbb{E}\left[J(\diamond_n) - \int_{\diamond_n} \frac{1}{2}\varphi \nabla v_n \cdot \mathbf{s} \nabla v_n\right] \\ &\quad + \frac{1}{2}Q \cdot \mathbb{E}[(\varphi - 1)\nabla v_n]_{\diamond_n} + \frac{1}{2}P \cdot \mathbb{E}[(\varphi - 1)\mathbf{a}\nabla v_n]_{\diamond_n} \\ &\quad + \frac{1}{2}Q \cdot (\mathbb{E}[(\nabla v_n)_{\diamond_n}] - P) + \frac{1}{2}P \cdot (\mathbb{E}[(\mathbf{a}\nabla v_n)_{\diamond_n}] - Q). \end{aligned} \quad (3.52)$$

We proceed by estimating each of the four lines on the right side of (3.52) separately.

*Step 1.* We show that there exists a constant  $C(d) < \infty$  such that

$$\left|\int_{\diamond_n} \varphi(\nabla v_n - P) \cdot (\mathbf{a}\nabla v_n - Q)\right| \leq C3^{-n}[\mathbf{m}_0^{1/2}(\nabla v_n - P)]_{\underline{B}_{2,1}^{-1/2}(\diamond_n)}[\mathbf{m}_0^{-1/2}(\mathbf{a}\nabla v_n - Q)]_{\underline{B}_{2,1}^{-1/2}(\diamond_n)}. \quad (3.53)$$

By Lemma 2.13 we may test the equation with  $(u - \ell_P)\varphi$  with  $\ell_P(x) = (v_n)_{\diamond_n} + P \cdot x$  and integrate by parts. Using (2.95) and the duality pairing between  $\underline{B}_{2,\infty}^{1/2}$  and  $\underline{B}_{2,1}^{-1/2}$ , we get that

$$\begin{aligned} \left|\int_{\diamond_n} \varphi(\nabla v_n - P) \cdot (\mathbf{a}\nabla v_n - Q)\right| &= \left|\int_{\diamond_n} (v_n - \ell_P)\mathbf{m}_0^{1/2}\nabla\varphi \cdot \mathbf{m}_0^{-1/2}(\mathbf{a}\nabla v_n - Q)\right| \\ &\leq C\|(v_n - \ell_P)\mathbf{m}_0^{1/2}\nabla\varphi\|_{\underline{B}_{2,\infty}^{1/2}(\diamond_n)}[\mathbf{m}_0^{-1/2}(\mathbf{a}\nabla v_n - Q)]_{\underline{B}_{2,1}^{-1/2}(\diamond_n)}. \end{aligned}$$

By applying (2.129), we find that

$$\|(v_n - \ell_P)\mathbf{m}_0^{1/2}\nabla\varphi\|_{\underline{B}_{2,\infty}^{1/2}(\diamond_n)} \leq C3^{-n}[\mathbf{m}_0^{1/2}(\nabla v_n - P)]_{\underline{B}_{2,1}^{-1/2}(\diamond_n)},$$

and (3.53) follows.

*Step 2.* We show that, for every  $k \in \mathbb{N}$  with  $k_0 \leq k \leq n$ ,

$$\left|\mathbb{E}\left[\int_{\diamond_n} \frac{1}{2}\varphi \nabla v_n \cdot \mathbf{s} \nabla v_n\right] - \mathbb{E}[J(\diamond_n)]\right| \leq 7\bar{\tau}_{n,k} + 5\bar{\tau}_{n,k}^{1/2}\mathbb{E}[J(\diamond_n)]^{1/2} + 3^{k+2-n}\mathbb{E}[J(\diamond_n)]. \quad (3.54)$$

We split the energy term, writing it as

$$\begin{aligned} \mathbb{E} \left[ \oint_{\diamond_n} \varphi \nabla v_n \cdot \mathbf{s} \nabla v_n \right] &= \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (\varphi)_{z+\diamond_k} \mathbb{E} \left[ \oint_{z+\diamond_k} \nabla v_n \cdot \mathbf{s} \nabla v_n \right] \\ &\quad + \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbb{E} \left[ \oint_{z+\diamond_k} (\varphi - (\varphi)_{z+\diamond_k}) \nabla v_n \cdot \mathbf{s} \nabla v_n \right]. \end{aligned} \quad (3.55)$$

The second term is very small: we have that

$$\begin{aligned} &\left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbb{E} \left[ \oint_{z+\diamond_k} (\varphi - (\varphi)_{z+\diamond_k}) \nabla v_n \cdot \mathbf{s} \nabla v_n \right] \right| \\ &\leq \max_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \|\varphi - (\varphi)_{z+\diamond_k}\|_{L^\infty(z+\diamond_k)} \mathbb{E} \left[ \oint_{\diamond_n} \nabla v_n \cdot \mathbf{s} \nabla v_n \right] \leq 3^{-(n-k)+2} \mathbb{E}[J(\diamond_n)]. \end{aligned} \quad (3.56)$$

Write next, for short, the energy density of  $J(z+\diamond_k)$  (similarly for  $J(\diamond_n)$ ) as

$$E_{k,z} = -\frac{1}{2} \nabla v_{k,z} \cdot \mathbf{s} \nabla v_{k,z} - \mathbf{a} \nabla v_{k,z} \cdot p + \nabla v_{k,z} \cdot q.$$

By (2.28) and (2.30), we have  $(E_{k,z})_{z+\diamond_k} = J(z+\diamond_k)$ ,  $(E_n)_{\diamond_n} = J(\diamond_n)$  and

$$\mathbb{E}[J(\diamond_n)] = \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbb{E} \left[ \oint_{z+\diamond_k} E_n \right] = \frac{1}{2} \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbb{E} \left[ \oint_{z+\diamond_k} (-\mathbf{a} \nabla v_n \cdot p + \nabla v_n \cdot q) \right].$$

Thus have the identity

$$\begin{aligned} &\mathbb{E}[J(\diamond_n)] - \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (\varphi)_{z+\diamond_k} \mathbb{E} \left[ \oint_{z+\diamond_k} \frac{1}{2} \nabla v_n \cdot \mathbf{s} \nabla v_n \right] \\ &= \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (1 - (\varphi)_{z+\diamond_k}) \mathbb{E}[J(z+\diamond_k)] \\ &\quad + \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (1 + (\varphi)_{z+\diamond_k}) \mathbb{E} \left[ \oint_{z+\diamond_k} (E_n - E_{k,z}) \right] \\ &\quad + \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (\varphi)_{z+\diamond_k} \mathbb{E} \left[ \oint_{z+\diamond_k} (\mathbf{a} \nabla(v_n - v_{k,z})p - \nabla(v_n - v_{k,z})q) \right]. \end{aligned} \quad (3.57)$$

The first term on the right side vanishes since, by  $\mathbb{Z}^d$ -stationarity and  $(\varphi)_{\diamond_n} = 1$ ,

$$\sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (1 - (\varphi)_{z+\diamond_k}) \mathbb{E}[J(z+\diamond_k)] = \mathbb{E}[J(\diamond_k)] \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (1 - (\varphi)_{z+\diamond_k}) = 0.$$

To bound the second term on the right of (3.57), we use (2.29) to get

$$\begin{aligned} &\left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbb{E} \left[ \oint_{z+\diamond_k} (E_{k,z} - E_n) \right] \right| \\ &= \frac{1}{2} \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbb{E} \left[ \oint_{z+\diamond_k} (\nabla v_{k,z} - \nabla v_n) \cdot \mathbf{s} (\nabla v_{k,z} - \nabla v_n) \right] = \bar{\tau}_{n,k}. \end{aligned} \quad (3.58)$$

To bound the third term on the right of (3.57), we apply (2.34) and Hölder's inequality to obtain

$$\begin{aligned}
& \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbb{E} \left[ \oint_{z + \diamond_k} (-\mathbf{a} \nabla(v_{k,z} - v_n) \cdot p + \nabla(v_{k,z} - v_n) \cdot q) \right] \right| \\
& \leq 4 \mathbb{E} \left[ \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (J(z + \diamond_k))^{1/2} \left( \oint_{z + \diamond_k} \frac{1}{2} (\nabla v_{k,z} - \nabla v_n) \cdot \mathbf{s} (\nabla v_{k,z} - \nabla v_n) \right)^{1/2} \right] \\
& \leq 4 \bar{\tau}_{n,k}^{1/2} \left( \mathbb{E}[J(\diamond_n)] + \bar{\tau}_{n,k} \right)^{1/2} \leq 4 \bar{\tau}_{n,k} + 4 \bar{\tau}_{n,k}^{1/2} \mathbb{E}[J(\diamond_n)]^{1/2}.
\end{aligned}$$

By combining the previous four displays with (3.56), we obtain (3.54).

*Step 3.* We next show that there exists  $C(d) < \infty$  such that, for every  $k \in \mathbb{N}$  with  $k_0 \leq k \leq n$ ,

$$\begin{aligned}
& |\mathbb{E}[(\varphi - 1) \nabla v_n]_{\diamond_n} \cdot Q| \leq 4 \bar{\tau}_{n,k}^{1/2} |\mathbf{s}_{*,0}^{-1/2} Q| \\
& + C 3^{-(n-k)} \mathbb{E}[J(\diamond_n)]^{1/2} |\mathbf{s}_{*,0}^{-1/2} Q| \left( \sum_{j=-\infty}^k 3^{\frac{3}{2}(j-k)} \sum_{z' \in z + 3^j \mathbb{L}_0 \cap \diamond_k} |\mathbf{E}_0^{-1/2} \overline{\mathbf{A}}(z' + \diamond_j) \mathbf{E}_0^{-1/2}| \right)^{1/2}. \quad (3.59)
\end{aligned}$$

Using the fact that  $(\varphi)_{\diamond_n} = 1$ , we have

$$\begin{aligned}
\mathbb{E}[(1 - \varphi) \nabla v_n]_{\diamond_n} &= \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (1 - (\varphi)_{z + \diamond_k}) \mathbb{E}[(\nabla v_n - \nabla v_{z,k})_{z + \diamond_k}] \\
&+ \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (1 - (\varphi)_{z + \diamond_k}) \mathbb{E}[(\nabla v_{z,k})_{z + \diamond_k}] \\
&+ \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbb{E} \left[ \oint_{z + \diamond_k} ((\varphi)_{z + \diamond_k} - \varphi) \nabla v_n \right]. \quad (3.60)
\end{aligned}$$

We next use a bound for the expectation of spatial averages of  $\nabla(v_n - v_{z,k})$  which is analogous to (2.38). We have

$$\begin{aligned}
& \frac{1}{2} \mathbb{E} \left[ (\nabla(v_n - v_{z,k}))_{z + \diamond_k} \cdot \overline{\mathbf{s}}_*(z + \diamond_k) \mathbb{E}[(\nabla(v_n - v_{z,k}))_{z + \diamond_k}] \right] \\
&= \sup_{q' \in \mathbb{R}^d} \mathbb{E} \left[ q' \cdot (\nabla(v_n - v_{z,k}))_{z + \diamond_k} - \frac{1}{2} q' \cdot \overline{\mathbf{s}}_*^{-1}(z + \diamond_k) q' \right] \\
&= \sup_{q' \in \mathbb{R}^d} \mathbb{E} \left[ q' \cdot (\nabla(v_n - v_{z,k}))_{z + \diamond_k} - J(z + \diamond_k, 0, q') \right] \\
&= \sup_{q' \in \mathbb{R}^d} \mathbb{E} \left[ \inf_{u \in \mathcal{A}(z + \diamond_k)} \oint_{z + \diamond_k} \left( \frac{1}{2} \nabla u \cdot \mathbf{s} \nabla u + q' \cdot \nabla(v_n - v_{z,k} - u) \right) \right] \\
&\leq \frac{1}{2} \mathbb{E} \left[ \oint_{z + \diamond_k} \nabla(v_n - v_{z,k}) \cdot \mathbf{s} \nabla(v_n - v_{z,k}) \right]. \quad (3.61)
\end{aligned}$$

The contribution of the first term on the right in (3.60) can then be estimated using the above display and  $\mathbb{Z}^d$ -stationarity:

$$\begin{aligned}
& \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \left| (1 - (\varphi)_{z + \diamond_k}) \mathbb{E}[(\nabla v_n - \nabla v_{z,k})_{z + \diamond_k}] \cdot Q \right| \\
& \leq 4 \bar{\tau}_{n,k}^{1/2} |\overline{\mathbf{s}}_*^{-1}(\diamond_k) Q| \leq 4 \bar{\tau}_{n,k}^{1/2} |\mathbf{E}_0^{-1/2} \overline{\mathbf{A}}(\diamond_k) \mathbf{E}_0^{-1/2}| |\mathbf{s}_{*,0}^{-1} Q|. \quad (3.62)
\end{aligned}$$

The second term on the right side of (3.60), by  $(\varphi)_{\diamond_n} = 1$  and  $\mathbb{Z}^d$ -stationarity, is zero:

$$\sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (1 - (\varphi)_{z+\diamond_k}) \mathbb{E}[(\nabla v_{z,k})_{z+\diamond_k}] = \mathbb{E}[(\nabla v_{0,k})_{\diamond_k}] \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} (1 - (\varphi)_{z+\diamond_k}) = 0.$$

The last term on the right side of (3.60) is small, and we start estimating it, by using (2.38):

$$\begin{aligned} \left| \oint_{z+\diamond_k} ((\varphi)_{z+\diamond_k} - \varphi) \nabla v_n \cdot Q \right| &\leq [\varphi]_{B_{1,\infty}^1(z+\diamond_k)} [\nabla v_n \cdot Q]_{B_{1,1}^{-1}(z+\diamond_k)} \\ &\leq C 3^{-n} \sum_{j=-\infty}^k 3^j \sum_{z' \in 3^j \mathbb{L}_0 \cap (z+\diamond_k)} |(\nabla v_n \cdot Q)_{z'+\diamond_j}| \\ &\leq C 3^{-n} \sum_{j=-\infty}^k 3^j \sum_{z' \in z+3^j \mathbb{L}_0 \cap \diamond_k} \|\mathbf{s}^{1/2} \nabla v_n\|_{\underline{L}^2(z'+\diamond_j)} |\mathbf{s}_*^{-1/2}(z'+\diamond_j) Q|. \end{aligned}$$

By Hölder's inequality, we then obtain

$$\begin{aligned} \mathbb{E} \left[ \left| \oint_{z+\diamond_k} ((\varphi)_{z+\diamond_k} - \varphi) \nabla v_n \cdot Q \right| \right] &\leq C 3^{-(n-k)} \mathbb{E}[J(\diamond_n)]^{1/2} \sum_{j=-\infty}^k 3^{\frac{3}{2}(j-k)} \left( \mathbb{E} \left[ \sum_{z' \in z+3^j \mathbb{L}_0 \cap \diamond_k} Q \cdot \mathbf{s}_*^{-1}(z'+\diamond_j) Q \right] \right)^{1/2} \\ &\leq C 3^{-(n-k)} \mathbb{E}[J(\diamond_n)]^{1/2} |\mathbf{s}_{*,0}^{-1/2} Q| \sum_{j=-\infty}^k 3^{\frac{3}{2}(j-k)} \left( \sum_{z' \in z+3^j \mathbb{L}_0 \cap \diamond_k} |\mathbf{E}_0^{-1/2} \overline{\mathbf{A}}(z'+\diamond_j) \mathbf{E}_0^{-1/2}| \right)^{1/2}. \quad (3.63) \end{aligned}$$

Combining the above displays yields (3.59).

*Step 4.* Similar to the previous step, we argue next that there exists  $C(d) < \infty$  such that, for every  $k \in \mathbb{N}$  with  $k_0 \leq k \leq n$ ,

$$\begin{aligned} |\mathbb{E}[(\varphi - 1) \mathbf{a} \nabla v_n]_{\diamond_n} \cdot P| &\leq 4 \overline{\tau}_{n,k}^{1/2} |\mathbf{b}_0^{1/2} P| \\ &+ C 3^{-(n-k)} \mathbb{E}[J(\diamond_n)]^{1/2} |\mathbf{b}_0^{1/2} P| \left( \sum_{j=-\infty}^k 3^{\frac{3}{2}(j-k)} \sum_{z' \in z+3^j \mathbb{L}_0 \cap \diamond_k} |\mathbf{E}_0^{-1/2} \overline{\mathbf{A}}(z'+\diamond_j) \mathbf{E}_0^{-1/2}| \right)^{1/2}. \quad (3.64) \end{aligned}$$

The proof is almost a verbatim repetition of Step 3 above using identities for fluxes in place of gradients, in particular, an appropriate version of (2.39) similar to (3.61). We omit the details.

*Step 5.* We next show that<sup>9</sup> there exists a constant  $C(d) < \infty$  such that, for every  $k, n \in \mathbb{N}$

---

<sup>9</sup>The statement asserted in Step 5 is valid without the assumptions of Proposition 3.3 and without the centering assumption (3.19), and for general  $p, q, P, Q \in \mathbb{R}^d$ . Indeed neither the assumptions of the proposition, the centering assumption, nor the choices of these parameters are used in the argument of Lemma 3.7 before Step 6. This more general statement will be used later, both in Section 4 and Section 5.

with  $k_0 \leq k \leq n-4$  and  $\varepsilon \in (0, 1]$ , we have that

$$\begin{aligned}
\left| \mathbb{E}[J(\diamond_n)] - \frac{1}{2}P \cdot Q \right| &\leq 50(1 + \varepsilon^{-1})\bar{\tau}_{n,k} + C(\varepsilon + 3^{-(n-k)})(|\mathbf{b}_0^{1/2}P| + |\mathbf{s}_{*,0}^{-1/2}Q|)^2 \\
&\quad + C3^{-(n-k)}(|\mathbf{b}_0^{1/2}P| + |\mathbf{s}_{*,0}^{-1/2}Q|)^2 \sum_{j=-\infty}^k 3^{\frac{3}{2}(j-k)} \sum_{z \in 3^j \mathbb{L}_0 \cap \diamond_k} |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}(z' + \diamond_j) \mathbf{E}_0^{-1/2}| \\
&\quad + \frac{1}{2} \left| Q \cdot (\mathbb{E}[(\nabla v_n)_{\diamond_n}] - P) + P \cdot (\mathbb{E}[(\mathbf{a} \nabla v_n)_{\diamond_n}] - Q) \right| \\
&\quad + C3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \left( \nabla v_n - P \right) \right]_{\underline{\dot{B}}_{2,1}^{-1/2}(\diamond_n)}^2 \right]. \tag{3.65}
\end{aligned}$$

Fix  $\varepsilon \in (0, 1]$ . To prove (3.65), we combine (3.52), (3.53), (3.54), (3.59) and (3.64) and obtain

$$\begin{aligned}
&\left| \mathbb{E}[J(\diamond_n)] - \frac{1}{2}P \cdot Q \right| \\
&\leq \bar{\tau}_{n,k}^{1/2} (7\bar{\tau}_{n,k}^{1/2} + 5\mathbb{E}[J(\diamond_n)]^{1/2} + 4|\mathbf{b}_0^{1/2}P| + 4|\mathbf{s}_{*,0}^{-1/2}Q|) + 3^{k+2-n} \mathbb{E}[J(\diamond_n)] \\
&\quad + C3^{-(n-k)} \mathbb{E}[J(\diamond_n)]^{1/2} (|\mathbf{b}_0^{1/2}P| + |\mathbf{s}_{*,0}^{-1/2}Q|) \left( \sum_{j=-\infty}^k 3^{\frac{3}{2}(j-k)} \sum_{z' \in z + 3^j \mathbb{L}_0 \cap \diamond_k} |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}(z' + \diamond_j) \mathbf{E}_0^{-1/2}| \right)^{1/2} \\
&\quad + \frac{1}{2} \left| Q \cdot (\mathbb{E}[(\nabla v_n)_{\diamond_n}] - P) + P \cdot (\mathbb{E}[(\mathbf{a} \nabla v_n)_{\diamond_n}] - Q) \right| \\
&\quad + C3^{-n} \mathbb{E} \left[ [\mathbf{m}_0^{1/2}(\nabla v_n - P)]_{\underline{\dot{B}}_{2,1}^{-1/2}(\diamond_n)} [\mathbf{m}_0^{-1/2}(\mathbf{a} \nabla v_n - Q)]_{\underline{B}_{2,1}^{-1/2}(\diamond_n)} \right].
\end{aligned}$$

By the triangle inequality and Young's inequality, we get that

$$\begin{aligned}
5\bar{\tau}_{n,k}^{1/2} \mathbb{E}[J(\diamond_n)]^{1/2} &\leq 5\bar{\tau}_{n,k}^{1/2} \left( \left| \mathbb{E}[J(\diamond_n)] - \frac{1}{2}P \cdot Q \right| + \frac{1}{2}|\mathbf{s}_{*,0}^{1/2}P \cdot \mathbf{s}_{*,0}^{-1/2}Q| \right)^{1/2} \\
&\leq 5\bar{\tau}_{n,k}^{1/2} \left( \left| \mathbb{E}[J(\diamond_n)] - \frac{1}{2}P \cdot Q \right|^{1/2} + \frac{1}{2}(|\mathbf{s}_{*,0}^{1/2}P| + |\mathbf{s}_{*,0}^{-1/2}Q|) \right) \\
&\leq \frac{1}{4} \left| \mathbb{E}[J(\diamond_n)] - \frac{1}{2}P \cdot Q \right| + (25 + \varepsilon^{-1})\bar{\tau}_{n,k} + \frac{25}{16} \varepsilon (|\mathbf{b}_0^{1/2}P| + |\mathbf{s}_{*,0}^{-1/2}Q|)^2.
\end{aligned}$$

Similarly,

$$5\bar{\tau}_{n,k}^{1/2} (|\mathbf{b}_0^{1/2}P| + |\mathbf{s}_{*,0}^{-1/2}Q|) \leq \frac{5}{2} \varepsilon^{-1} \bar{\tau}_{n,k} + \frac{5}{2} \varepsilon (|\mathbf{b}_0^{1/2}P| + |\mathbf{s}_{*,0}^{-1/2}Q|)^2.$$

Since  $n \geq k+4$ , we also deduce that

$$3^{k+2-n} \mathbb{E}[J(\diamond_n)] \leq \frac{1}{16} \left| \mathbb{E}[J(\diamond_n)] - \frac{1}{2}P \cdot Q \right| + C3^{-(n-k)} (|\mathbf{b}_0^{1/2}P| + |\mathbf{s}_{*,0}^{-1/2}Q|)^2$$

and, by Hölder's inequality,

$$\begin{aligned}
&3^{-(n-k)} \mathbb{E}[J(\diamond_n)]^{1/2} (|\mathbf{b}_0^{1/2}P| + |\mathbf{s}_{*,0}^{-1/2}Q|) \left( \sum_{j=-\infty}^k 3^{\frac{3}{2}(j-k)} \sum_{z' \in z + 3^j \mathbb{L}_0 \cap \diamond_k} |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}(z' + \diamond_j) \mathbf{E}_0^{-1/2}| \right)^{1/2} \\
&\leq \frac{1}{16} \left| \mathbb{E}[J(\diamond_n)] - \frac{1}{2}P \cdot Q \right| \\
&\quad + C3^{-(n-k)} (|\mathbf{b}_0^{1/2}P| + |\mathbf{s}_{*,0}^{-1/2}Q|)^2 \sum_{j=-\infty}^k 3^{\frac{3}{2}(j-k)} \sum_{z' \in z + 3^j \mathbb{L}_0 \cap \diamond_k} (1 + |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}(z' + \diamond_j) \mathbf{E}_0^{-1/2}|).
\end{aligned}$$

Consequently, we obtain (3.65) by Young's inequality.

*Step 6.* We finally prove (3.51). We fix  $n := m - l$  and  $k := n - l$ , and estimate the various quantities appearing in (3.65). First, we use (3.24) and (2.23) to control the additivity defect:

$$\begin{aligned} \bar{\tau}_{n,k} &= \mathbb{E}[J(\diamond_k) - J(\diamond_n)] = \frac{1}{2} \binom{-p}{q} \cdot (\bar{\mathbf{A}}(\diamond_k) - \bar{\mathbf{A}}(\diamond_n)) \binom{-p}{q} \\ &\leq \max_{j \in \{k,n\}} \left| \bar{\mathbf{A}}_m^{-1/2} \bar{\mathbf{A}}(\diamond_j) \bar{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d} \right| \left| \bar{\mathbf{A}}_m^{1/2} \binom{-p}{q} \right|^2 \leq C \delta \sigma^2 \left| \bar{\mathbf{A}}_m^{1/2} \binom{-p}{q} \right|^2. \end{aligned} \quad (3.66)$$

Second, by (3.17), we see that since  $k = m - 2l > 90l$

$$\frac{K_{\Psi_S}^{90} \Pi^{90}}{(1 - \gamma)^{90}} (3^{-(m-2l)} + 3^{-k}) \leq 1. \quad (3.67)$$

On the one hand, using (3.67) and (2.107) we see that

$$\sum_{j=k_0}^k 3^{\frac{3}{2}(j-k)} \sum_{z \in 3^j \mathbb{L}_0 \cap \diamond_k} |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}(z + \diamond_j) \mathbf{E}_0^{-1/2}| \leq \frac{3}{2} |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}_0 \mathbf{E}_0^{-1/2}| + C \Pi \sum_{j=k_0}^k 3^{-2k+j} \leq \frac{3}{2} + C \Pi 3^{-k}.$$

Here  $k_0(d)$  is as in (2.100) guaranteeing that  $\bar{\mathbf{A}}(z' + \diamond_j) = \bar{\mathbf{A}}(\diamond_j)$  for every  $j \in \mathbb{N}$  with  $j \geq k_0$ . On the other hand, letting  $\mathcal{S}_{h'}$  be as in Lemma 2.8 (applied with  $h = 0$ ,  $\delta = 1$  and the smallest integer  $h'$  satisfying (2.101)) and (3.67),

$$\begin{aligned} &\sum_{j=-\infty}^{k_0} 3^{\frac{3}{2}(j-k)} \sum_{z \in 3^j \mathbb{L}_0 \cap \diamond_k} |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}(z + \diamond_j) \mathbf{E}_0^{-1/2}| \\ &\leq C \sum_{j=-\infty}^{k_0} 3^{\frac{3}{2}(j-k)} (3^{\gamma(m-j)} + \mathbb{E}[S]^\gamma 3^{-\gamma j}) \leq C \left( \frac{K_{\Psi_S}^2 \Pi}{1 - \gamma} \right)^\gamma 3^{-k}. \end{aligned}$$

Therefore, (3.67) yields that

$$\sum_{j=-\infty}^k 3^{\frac{3}{2}(j-k)} \sum_{z' \in z + 3^j \mathbb{L}_0 \cap \diamond_k} |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}(z' + \diamond_j) \mathbf{E}_0^{-1/2}| \leq C \left( 1 + \frac{K_{\Psi_S}^2 \Pi}{1 - \gamma} 3^{-k} \right) \leq C. \quad (3.68)$$

Third, we give the estimate for  $|\mathbf{b}_0^{1/2} P| + |\mathbf{s}_{*,0}^{-1/2} Q|$ . For this, define

$$\widetilde{\mathbf{M}}_0 := \begin{pmatrix} \mathbf{b}_0 & 0 \\ 0 & \mathbf{s}_{*,0}^{-1} \end{pmatrix} \quad \text{and} \quad \mathbf{R} := \begin{pmatrix} 0 & \mathbf{I}_d \\ \mathbf{I}_d & 0 \end{pmatrix}.$$

In view of (3.44), (2.48) and (3.48), we have that

$$\begin{aligned} \left| \widetilde{\mathbf{M}}_0^{1/2} \begin{pmatrix} P \\ Q \end{pmatrix} \right|^2 &= \binom{-p}{q} \cdot ((\bar{\mathbf{A}}_m \mathbf{R} - \mathbf{I}_{2d}) \widetilde{\mathbf{M}}_0 (\mathbf{R} \bar{\mathbf{A}}_m - \mathbf{I}_{2d})) \binom{-p}{q} \\ &\leq 2 \left( |\bar{\mathbf{A}}_m^{1/2} \mathbf{R} \widetilde{\mathbf{M}}_0 \mathbf{R} \bar{\mathbf{A}}_m^{1/2}| + |\bar{\mathbf{A}}_m^{-1/2} \widetilde{\mathbf{M}}_0 \bar{\mathbf{A}}_m^{-1/2}| \right) \left| \bar{\mathbf{A}}_m^{1/2} \binom{-p}{q} \right|^2. \end{aligned}$$

Since  $\bar{\mathbf{A}}_m \leq \bar{\mathbf{A}}_0 \leq \mathbf{E}_0$ , we obtain by (2.23) that

$$|\bar{\mathbf{A}}_m^{1/2} \mathbf{R} \widetilde{\mathbf{M}}_0 \mathbf{R} \bar{\mathbf{A}}_m^{1/2}| \leq 2 (|\mathbf{s}_{*,0}^{-1/2} \mathbf{b}_m \mathbf{s}_{*,0}^{-1/2}| \vee |\mathbf{s}_{*,m}^{-1/2} \mathbf{b}_0 \mathbf{s}_{*,m}^{-1/2}|) \leq 2\Theta (|\mathbf{b}_0^{-1} \bar{\mathbf{b}}_m| \vee |\mathbf{s}_{*,m}^{-1/2} \mathbf{s}_{*,0} \mathbf{s}_{*,m}^{-1/2}|) \leq 2\Theta$$

and

$$|\overline{\mathbf{A}}_m^{-1/2} \widetilde{\mathbf{M}}_0 \overline{\mathbf{A}}_m^{-1/2}| = |\widetilde{\mathbf{M}}_0^{1/2} \overline{\mathbf{A}}_m^{-1} \widetilde{\mathbf{M}}_0^{1/2}| \leq 2(|\mathbf{b}_0^{1/2} \mathbf{s}_m^{-1} \mathbf{b}_0^{1/2}| \vee |\mathbf{s}_{*,0}^{-1/2} (\mathbf{s}_{*,m} + \mathbf{k}_m \mathbf{s}_m^{-1} \mathbf{k}_m^t) \mathbf{s}_{*,0}^{-1/2}|) \leq 2\Theta.$$

It follows that

$$|\mathbf{b}_0^{1/2} P| + |\mathbf{s}_{*,0}^{-1/2} Q| = \left| \widetilde{\mathbf{M}}_0^{1/2} \begin{pmatrix} P \\ Q \end{pmatrix} \right| \leq 4\Theta^{1/2} \left| \overline{\mathbf{A}}_m^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|. \quad (3.69)$$

Fourth, using the particular choice of  $P$  and  $Q$  in (3.48) and (2.33), we have, by (3.6) and (3.24),

$$\begin{aligned} \left| \begin{pmatrix} Q \\ P \end{pmatrix} \cdot \mathbb{E} \left[ \int_{\diamond_n} \begin{pmatrix} \nabla v_n \\ \mathbf{a} \nabla v_n \end{pmatrix} - \begin{pmatrix} P \\ Q \end{pmatrix} \right] \right| &= \left| \begin{pmatrix} P \\ Q \end{pmatrix} \cdot (\overline{\mathbf{A}}(\diamond_n) - \overline{\mathbf{A}}_m) \cdot \begin{pmatrix} -p \\ q \end{pmatrix} \right| \\ &\leq |\mathbf{E}_0^{-1/2} (\overline{\mathbf{A}}(\diamond_n) - \overline{\mathbf{A}}_m) \mathbf{E}_0^{-1/2}| \left| \mathbf{E}_0^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right| \left| \mathbf{E}_0^{1/2} \begin{pmatrix} P \\ Q \end{pmatrix} \right| \\ &\leq C\delta\sigma^2 \left| \overline{\mathbf{A}}_m^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right| (|\mathbf{b}_0^{1/2} P| + |\mathbf{s}_{*,0}^{-1/2} Q|). \end{aligned} \quad (3.70)$$

Inserting now (3.68), (3.70), (3.66) and (3.69) into (3.65) yields that

$$\begin{aligned} \left| \mathbb{E}[J(\diamond_n)] - \frac{1}{2} P \cdot Q \right| &\leq C3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \begin{pmatrix} \nabla v_n - P \\ \mathbf{a} \nabla v_n - Q \end{pmatrix} \right]_{\underline{B}_{2,1}^{-1/2}(\diamond_n)}^2 \right] \\ &\quad + C(\delta\sigma^2(\varepsilon^{-1} + \Theta^{1/2}) + (\varepsilon + 3^{-l})\Theta) \left| \overline{\mathbf{A}}_m^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2. \end{aligned}$$

By (3.17),  $3^{-l}\Theta \leq \delta\sigma^2$ . Thus, by taking  $\varepsilon := \delta^{1/2}\sigma\Theta^{-1/2}$ , we deduce that

$$\left| \mathbb{E}[J(\diamond_n)] - \frac{1}{2} P \cdot Q \right| \leq C\delta^{1/2}\sigma\Theta^{1/2} \left| \overline{\mathbf{A}}_m^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2.$$

To conclude, we hence need to estimate the last term on the right. We claim that

$$\left| \overline{\mathbf{A}}_m^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2 \leq 2|\overline{\mathbf{b}}_m^{1/2} p|^2 \vee |\overline{\mathbf{s}}_{*,m}^{-1/2} q|^2 \leq 2|\overline{\mathbf{s}}_{*,m}^{-1} \overline{\mathbf{b}}_m|^{1/2} \leq C\Theta_m^{1/2}, \quad (3.71)$$

which obviously then concludes the proof. To show (3.71), the first inequality is a consequence of (2.23). Next, since  $\mathbf{t}_m := \overline{\mathbf{b}}_m \# \overline{\mathbf{s}}_{*,m}^{-1}$ , by the properties of the metric geometric mean, namely the equation (B.1), we have that

$$\mathbf{t}_m^{-1/2} \overline{\mathbf{b}}_m \mathbf{t}_m^{-1/2} = \mathbf{t}_m^{1/2} \overline{\mathbf{s}}_{*,m}^{-1} \mathbf{t}_m^{1/2}.$$

Thus

$$\begin{aligned} |\mathbf{t}_m^{-1/2} \overline{\mathbf{b}}_m \mathbf{t}_m^{-1/2}| &= |\overline{\mathbf{s}}_{*,m}^{-1/2} \mathbf{t}_m \overline{\mathbf{s}}_{*,m}^{-1/2}| = |\overline{\mathbf{s}}_{*,m}^{-1/2} \mathbf{t}_m \overline{\mathbf{s}}_{*,m}^{-1} \mathbf{t}_m \overline{\mathbf{s}}_{*,m}^{-1/2}|^{1/2} = |\overline{\mathbf{s}}_{*,m}^{-1/2} \mathbf{t}_m^{1/2} (\mathbf{t}_m^{-1/2} \overline{\mathbf{b}}_m \mathbf{t}_m^{-1/2}) \mathbf{t}_m^{1/2} \overline{\mathbf{s}}_{*,m}^{-1/2}|^{1/2} \\ &= |\overline{\mathbf{s}}_{*,m}^{-1/2} \overline{\mathbf{b}}_m \overline{\mathbf{s}}_{*,m}^{-1/2}|^{1/2}. \end{aligned} \quad (3.72)$$

From the above display, we obtain

$$|\overline{\mathbf{b}}_m^{1/2} p| = |\overline{\mathbf{b}}_m^{1/2} \mathbf{t}_m^{-1/2} e| \leq |\mathbf{t}_m^{-1/2} \overline{\mathbf{b}}_m \mathbf{t}_m^{-1/2}|^{1/2} = |\overline{\mathbf{s}}_{*,m}^{-1/2} \overline{\mathbf{b}}_m \overline{\mathbf{s}}_{*,m}^{-1/2}|^{1/4} \quad (3.73)$$

and, similarly,

$$|\overline{\mathbf{s}}_{*,m}^{-1/2} q| = |\overline{\mathbf{s}}_{*,m}^{-1/2} \mathbf{t}_m^{1/2} e| \leq |\overline{\mathbf{s}}_{*,m}^{-1} \mathbf{t}_m|^{1/2} = |\overline{\mathbf{s}}_{*,m}^{-1/2} \overline{\mathbf{b}}_m \overline{\mathbf{s}}_{*,m}^{-1/2}|^{1/4}. \quad (3.74)$$

Thus we have proved (3.71) in view of (3.35). We may now use the definition of  $l$  in (3.17) to deduce that  $3^{-l}\Theta \leq \delta\sigma$ . The proof is complete.  $\square$

The previous lemma reduces our task to estimating the right side of (3.51). The is the content of the following lemma, which is based on an application of Lemma 2.14.

**Lemma 3.8.** *There exists a constant  $C(d) < \infty$  such that, for  $n = m - l$ , we have that*

$$3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \left( \nabla v(\cdot, \diamond_n, p, q) - P \right) \right]_{\underline{\dot{B}}_{2,1}^{-1/2}(\diamond_n)}^2 \right] \leq C \delta \sigma^2 \Theta_m^{1/2} \Theta^{1/2}. \quad (3.75)$$

*Proof.* Denote  $v_n := v(\cdot, \diamond_n, p, q)$ . Towards the application of Lemma 2.14, we take  $h = l$ . Notice that since  $n = m - l > 90l$ , we have, by (3.17), that

$$\frac{\Pi^4 K_{\Psi_S}^{4d+15}}{(1-\gamma)^5} 3^{-n} \leq \delta \sigma^2 \quad \text{and} \quad \frac{3^{-(1-\gamma)l}}{1-\gamma} \leq \delta \sigma^2. \quad (3.76)$$

Setting  $\mathbf{M} = \mathbf{M}_0$  and  $\mathbf{E} = \mathbf{E}_0$ , we observe that by (3.20) we have

$$|\mathbf{M}^{-1/2} \mathbf{E} \mathbf{M}^{-1/2}| \leq C \Theta^{1/2}. \quad (3.77)$$

Using (3.21), we see that  $\mathbf{E}_0 \leq 2\bar{\mathbf{A}}_m$  and, by (3.71) and (3.44), we have the estimate

$$\max \left\{ \left| \mathbf{E}_0^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|, |p \cdot q| \right\} \leq \max \left\{ 2 \left| \bar{\mathbf{A}}_m^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|, 1 \right\} \leq C \Theta_m^{1/4}. \quad (3.78)$$

Moreover, by the definitions of  $p$  and  $q$ ,

$$\|\mathbf{s}^{1/2} \nabla v_n\|_{\underline{\mathbb{L}}^2(\diamond_n)}^2 = \begin{pmatrix} -p \\ q \end{pmatrix} \cdot \mathbf{A}(\diamond_n) \begin{pmatrix} -p \\ q \end{pmatrix} - 2p \cdot q \leq \mathcal{M}_{n,\gamma} \left| \mathbf{E}_0^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2 \leq C \mathcal{M}_{n,\gamma} \Theta_m^{1/2}. \quad (3.79)$$

We next apply Lemma 2.14: we use the inequality (2.133) with  $h = n - l$ , square it, and then take the expectation of the result, and substitute for some factors on the right side with (3.77) and (3.78). Finally, we get rid of the squares on the right side by using Hölder's inequality for the sums in the following form

$$\begin{aligned} \left( \sum_{k=n-l}^n 3^{\frac{1}{2}(k-n)} X_k^{1/2} \right)^2 &= \left( \sum_{k=n-l}^n 3^{\frac{1}{2}\varepsilon(k-n)} 3^{\frac{1}{2}(1-\varepsilon)(k-n)} X_k^{1/2} \right)^2 \\ &\leq \left( \sum_{k=n-l}^n 3^{\varepsilon(k-n)} \right) \left( \sum_{k=n-l}^n 3^{(1-\varepsilon)(k-n)} X_k \right) \leq \frac{C}{\varepsilon} \sum_{k=n-l}^n 3^{(1-\varepsilon)(k-n)} X_k. \end{aligned}$$

As a result, we obtain, using also (3.79), for every  $\varepsilon \in (0, 1)$ ,

$$\begin{aligned} &3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \left( \nabla v_n - (\nabla v_n)_{\diamond_n} \right) \right]_{\underline{\dot{B}}_{2,1}^{-1/2}(\diamond_n)}^2 \right] \\ &\leq \frac{C \Theta^{1/2}}{\varepsilon} \left| \mathbf{E}_0^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2 \sum_{k=n-l}^n 3^{(1-\varepsilon)(k-n)} \mathbb{E} \left[ \left| \mathbf{E}_0^{-1/2} (\mathbf{A}(\diamond_k) - \mathbf{A}(\diamond_n)) \mathbf{E}_0^{-1/2} \right|^2 \right] \\ &\quad + \frac{C \Theta^{1/2}}{\varepsilon} \left| \mathbf{E}_0^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2 \sum_{k=n-l}^n 3^{(1-\varepsilon)(k-n)} \mathbb{E} \left[ \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbf{E}_0^{-1/2} (\mathbf{A}(z + \diamond_k) - \mathbf{A}(\diamond_n)) \mathbf{E}_0^{-1/2} \right|^2 \right] \\ &\quad + \frac{C \Theta^{1/2}}{1-\gamma} \left| \mathbf{E}_0^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2 \left( 3^{-(1-\gamma)l} + \mathbb{E} [\mathcal{M}_{n,\gamma}^2 \mathbf{1}_{\{\mathcal{M}_{n,\gamma} > 1\}}] \right). \end{aligned} \quad (3.80)$$



We will take  $\varepsilon = 1/2$  in the above display. Using (3.21), (3.24), (3.25) and the triangle inequality, the first term on the right is bounded by  $C\delta\sigma^2\Theta^{1/2}\Theta_m^{1/2}$ . Similarly, by (3.21) and (3.24),

$$\begin{aligned} \sum_{k=n-l}^n 3^{(1-\varepsilon)(k-n)} \mathbb{E} \left[ \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbf{E}_0^{-1/2} (\mathbf{A}(z + \diamond_k) - \mathbf{A}(\diamond_n)) \mathbf{E}_0^{-1/2} \right| \right] \\ \leq \sum_{k=n-l}^n 3^{(1-\varepsilon)(k-n)} \text{trace} (\mathbf{E}_0^{-1/2} (\overline{\mathbf{A}}(\diamond_k) - \overline{\mathbf{A}}(\diamond_n)) \mathbf{E}_0^{-1/2}) \leq C\delta\sigma^2. \end{aligned} \quad (3.81)$$

The second inequality is valid since, by subadditivity,  $\sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \mathbf{A}(z + \diamond_k) \geq \mathbf{A}(\diamond_n)$ . Furthermore, letting  $\mathcal{S}_{h'}$  be as in in Lemma 2.8 (with  $h'$  being the smallest integer such that (2.101) is valid with  $\delta = 1$  and  $h = 0$ ), and recalling the definition in (2.132), we have the implication

$$3^n \geq \mathcal{S}_{h'} \implies \mathcal{M}_{n,\gamma} \leq \sup_{k \in \mathbb{Z} \cap (-\infty, n]} 3^{-\gamma(n-k)} \max_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{E}_0^{-1} \mathbf{A}(y + \diamond_k)| \leq 1.$$

Therefore, by (C.5) and (3.76),

$$\frac{1}{1-\gamma} \mathbb{E}[\mathcal{M}_{n,\gamma}^2 \mathbf{1}_{\{\mathcal{M}_{n,\gamma} > 1\}}] \leq \frac{CK_{\Psi_S}^6 \Pi^3 3^{-n}}{(1-\gamma)^4} \leq C\delta\sigma^2 \quad (3.82)$$

and

$$\frac{1}{1-\gamma} 3^{-(1-\gamma)l} \leq C\delta\sigma^2. \quad (3.83)$$

By combining the above displays we obtain that

$$3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \left( \begin{array}{c} \nabla v(\cdot, \diamond_n, p, q) - (\nabla v_n)_{\diamond_n} \\ \mathbf{a} \nabla v(\cdot, \diamond_n, p, q) - (\mathbf{a} \nabla v_n)_{\diamond_n} \end{array} \right) \right]_{\hat{B}_{2,1}^{-1/2}(\diamond_n)}^2 \right] \leq C\delta\sigma^2 \Theta_m^{1/2} \Theta^{1/2}.$$

The last thing to check is that

$$\mathbb{E} \left[ \left| \mathbf{M}_0^{1/2} \left( \begin{array}{c} P - (\nabla v_n)_{\diamond_n} \\ Q - (\mathbf{a} \nabla v_n)_{\diamond_n} \end{array} \right) \right|^2 \right] \leq C\delta\sigma^2 \Theta_m^{1/2} \Theta^{1/2}.$$

To see this, we write  $\mathbf{M}_0^{1/2} = \mathbf{R}(\mathbf{R}\mathbf{M}_0^{1/2}\mathbf{R}\mathbf{E}_0^{1/2})\mathbf{E}_0^{-1/2}\mathbf{R} = \mathbf{R}(\mathbf{M}_0^{-1/2}\mathbf{E}_0^{1/2})\mathbf{E}_0^{-1/2}\mathbf{R}$  with  $\mathbf{R}$  as in (2.32), and estimate the matrix in the middle using (3.20) as

$$|\mathbf{M}_0^{-1/2}\mathbf{E}_0^{1/2}|^2 = |\mathbf{M}_0^{-1/2}\mathbf{E}_0\mathbf{M}_0^{-1/2}| \leq 2(|\mathbf{m}_0^{-1/2}\mathbf{b}_0\mathbf{m}_0^{-1/2}| + |\mathbf{s}_{*,0}^{-1/2}\mathbf{m}_0\mathbf{s}_{*,0}^{-1/2}|) \leq C\Theta^{1/2}.$$

Thus, we obtain that, by (2.33) (3.78), (3.21) and (3.25),

$$\begin{aligned} \mathbb{E} \left[ \left| \mathbf{M}_0^{1/2} \left( \begin{array}{c} P - (\nabla v_n)_{\diamond_n} \\ Q - (\mathbf{a} \nabla v_n)_{\diamond_n} \end{array} \right) \right|^2 \right] &\leq |\mathbf{M}_0^{-1/2}\mathbf{E}_0^{1/2}|^2 \mathbb{E} \left[ |\mathbf{E}_0^{-1/2} (\mathbf{A}(\diamond_n) - \overline{\mathbf{A}}_m) \mathbf{E}_0^{-1/2}|^2 \right] \left| \mathbf{E}_0^{1/2} \left( \begin{array}{c} -p \\ q \end{array} \right) \right|^2 \\ &\leq C\Theta_m^{1/2} \Theta^{1/2} |\mathbf{E}_0^{-1/2} \overline{\mathbf{A}}_m \mathbf{E}_0^{-1/2}| \mathbb{E} \left[ |\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}(\diamond_n) \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}|^2 \right] \\ &\leq C\delta\sigma^2 \Theta_m^{1/2} \Theta^{1/2}. \end{aligned} \quad (3.84)$$

The proof is complete.  $\square$

We turn to the proof of Proposition 3.3.

*Proof of Proposition 3.3.* We just need to assemble the estimates we have proved above and choose the parameters appropriately. As above, we set  $n := m - l$ .

By combining Lemma 3.7 and 3.8, and the analogous estimate for  $J^*$ , we obtain, for a constant  $C(d) < \infty$ ,

$$\left| \mathbb{E}[J(\diamond_n, p, q)] - \frac{1}{2}P \cdot Q \right| + \left| \mathbb{E}[J^*(\diamond_n, p, q)] - \frac{1}{2}P^* \cdot Q^* \right| \leq C\delta^{1/2}\sigma\Theta_m^{1/2}\Theta^{1/2}. \quad (3.85)$$

We now assume that

$$\delta_0 \leq (8C_{(3.85)})^{-2}.$$

Applying Young's inequality and using  $\delta \leq \delta_0$ , we obtain

$$\left| \mathbb{E}[J(\diamond_n, p, q)] - \frac{1}{2}P \cdot Q \right| + \left| \mathbb{E}[J^*(\diamond_n, p, q)] - \frac{1}{2}P^* \cdot Q^* \right| \leq \frac{\sigma}{16}\Theta_m + \frac{\sigma}{16}\Theta.$$

Since  $l = m - n$ , we have by (3.17) that, if  $\delta_0(d)$  is sufficiently small, then

$$C\Pi 3^{n-m} \leq C\delta\sigma^2 \leq \frac{1}{8}\sigma \leq \frac{1}{8}\sigma\Theta.$$

Turning back to (3.46), and combining it with (3.47) and the previous display, we get that

$$\begin{aligned} \Theta_m - 1 &\leq 2d\mathbb{E}[\tilde{J}(\square_m, p, q) + \tilde{J}^*(\square_m, p, q)] \\ &\leq 2d\left(\mathbb{E}[J(\diamond_n, p, q) + J^*(\diamond_n, p, q)] - \frac{1}{2}P \cdot Q - \frac{1}{2}P^* \cdot Q^*\right) + C\Pi 3^{n-m} \\ &\leq \frac{\sigma}{4}\Theta_m + \frac{\sigma}{4}\Theta. \end{aligned}$$

Reorganizing this, we obtain

$$\left(1 - \frac{\sigma}{4}\right)(\Theta_m - 1) \leq \frac{\sigma}{2}\Theta,$$

and since  $1 - \frac{1}{4}\sigma > \frac{1}{2}$ , we consequently deduce that

$$\Theta_m - 1 \leq \sigma\Theta. \quad (3.86)$$

This completes the proof.  $\square$

**3.3. The iteration from high contrast to small contrast.** We give the proof of Theorem 3.1, which is based on an iteration of Proposition 3.2, renormalizing between each iteration step by appealing to Proposition 2.6.

*Proof of Theorem 3.1.* We introduce the parameters

$$\begin{cases} \gamma_* := \frac{1}{2}(\min\{\nu, 1\} + \gamma) \\ K_{\Psi_S}^* := \max\{K_{\Psi_S}, K_{\Psi}^{[1/\mu]}\}, \\ \Theta^* := 4\Theta, \\ \Pi^* := 2^{10}\Pi. \end{cases} \quad (3.87)$$

Also denote  $\rho := \frac{1}{2}(\min\{\nu, 1\} + \gamma_*)$  and  $\alpha^* := (\min\{\nu, 1\} - \gamma^*)(1 - \beta)$ . Motivated by (2.86), we let  $l_0 \in \mathbb{N}$  be defined by

$$l_0 := \left\lceil \frac{1}{\rho - \gamma^*} \left(1 + \frac{d}{\alpha^*}\right) (5 + \log(\Theta^*)) + \frac{6}{\alpha^*} (1 + \log K_{\Psi}) \right\rceil, \quad (3.88)$$

Motivated by (3.3), we set  $\sigma_0 := 1/8$  and define

$$l_1 := \left\lceil \frac{C}{\sigma_0^2} \left( \log K_{\Psi_S}^* + \frac{1}{(\alpha^*)^2} \log \left( \frac{\Pi^* K_{\Psi}}{\alpha^* \sigma_0^2} \right) \right) \log(1 + \Theta^*) \right\rceil, \quad (3.89)$$

where  $C(d) < \infty$  is the constant in Proposition 3.2. In terms of the original parameters, we have that

$$l_0 \leq \left\lceil \frac{4}{\alpha} \left( 1 + \frac{2d}{\alpha} \right) (7 + \log(\Theta)) + \frac{12}{\alpha} (1 + \log K_{\Psi}) \right\rceil, \quad (3.90)$$

where  $\alpha = (\min\{\nu, 1\} - \gamma)(1 - \beta)$ ; and, by inflating  $C$  by an additional (universal) factor,

$$l_1 \leq C \left( \log K_{\Psi_S} + \frac{1}{\alpha^2} \log \left( \frac{\Pi K_{\Psi}}{\alpha} \right) \right) \log(1 + \Theta). \quad (3.91)$$

We also denote  $m_0 := l_0 + \lceil \log K_{\Psi} \rceil$ .

For each  $n \in \mathbb{N}$  with  $n \geq m_0$ , we may apply Proposition 2.6 to obtain that the pushforward probability measure  $\mathbb{P}_n$ , defined in (2.83), satisfies the assumptions (P1), (P2) and (P3) with  $\delta \leq 1$  and the new parameters

$$\begin{cases} \mathbf{E}_{\text{new}} := 2\overline{\mathbf{A}}(\square_{n-2l_0}), \\ \gamma_{\text{new}} := \gamma^* \\ K_{\Psi, \text{new}} := K_{\Psi} \\ K_{\Psi_S, \text{new}} := K_{\Psi_S}^*, \\ \Theta_{\text{new}} := 4\Theta_{n-2l_0} \leq 4\Theta = \Theta^*, \\ \Pi_{\text{new}} \leq \Pi^*. \end{cases} \quad (3.92)$$

For such  $n$ , applying Proposition 3.2 with  $\mathbb{P}_n$  in place of  $\mathbb{P}$ , with  $\delta = 1$ , and with  $\sigma = \sigma_0 = 1/8$  as above, we obtain that

$$\Theta_{n+l_1} - 1 \leq \frac{1}{2} \Theta_{n-2l_0}.$$

Rephrasing this a bit, what we have shown is that

$$\Theta_{n+2l_0+l_1} - 2 \leq \frac{1}{2} (\Theta_n - 2), \quad \forall n \geq m_0 + 2l_0.$$

An iteration now yields

$$(\Theta_{m_0+2l_0+k(2l_0+l_1)} - 2) \leq 2^{-k} (\Theta - 2)_+.$$

Since  $n \mapsto \Theta_n$  is monotone decreasing in  $n$ , we conclude that

$$n \geq m_1 := m_0 + 2l_0 + \frac{\log \Theta_{m_0}}{\log 2} (2l_0 + l_1) \implies \Theta_n \leq 3.$$

This gives us a quantitative scale  $m_1$  such that  $\Theta_{m_1} \leq 3$ .

We next intend to apply Propositions 2.6 and 3.2 once more to make the ellipticity ratio as close to one as we like. We now take a small parameter  $\sigma \in (0, 1/2]$ , and we argue as above, applying these propositions with  $\delta = 1$  and the renormalized parameters in (3.92) and with  $\sigma/12$  in place of  $\sigma_0$ , and with  $\lceil (\sigma/12)^{-2} |\log \sigma| l_1 \rceil$  in place of  $l_1$ . We obtain

$$n \geq m_1 + 2l_0 + \lceil (\sigma/12)^{-2} |\log \sigma| l_1 \rceil \implies \Theta_n - 1 \leq \frac{\sigma}{12} \cdot 4\Theta_{m_1} \leq \sigma.$$

It is very straightforward to check that, for a constant  $C(d) < \infty$ , we have

$$m_1 + 2l_0 + \lceil (\sigma/12)^{-2} |\log \sigma| l_1 \rceil \leq \frac{C}{\sigma^2} \left( \log K_{\Psi_S} + \frac{1}{\alpha^2} \log \left( \frac{\Pi K_{\Psi}}{\alpha \sigma^2} \right) \right) \log^2(1 + \Theta).$$

This completes the proof of the theorem.  $\square$

#### 4. Renormalization in small contrast

Theorem 3.1 implies the qualitative limit  $\lim_{m \rightarrow \infty} (\Theta_m - 1) = 0$ , and this allows us to identify the homogenized matrix. To see this, we recall that, by (2.54), each of the maps  $n \mapsto \bar{\mathbf{s}}(\square_n)$  and  $n \mapsto \bar{\mathbf{s}}_*^{-1}(\square_n)$  is nonincreasing and bounded. Thus there exist  $\bar{\mathbf{s}}$  and  $\bar{\mathbf{s}}_*^{-1}$  such that  $\bar{\mathbf{s}}_* \leq \bar{\mathbf{s}}$  and

$$\bar{\mathbf{s}} = \lim_{n \rightarrow \infty} \bar{\mathbf{s}}(\square_n) \quad \text{and} \quad \bar{\mathbf{s}}_*^{-1} = \lim_{n \rightarrow \infty} \bar{\mathbf{s}}_*^{-1}(\square_n).$$

The definition of  $\Theta_n$  implies that, for every  $n \in \mathbb{N}$ ,

$$\bar{\mathbf{s}}_* \leq \bar{\mathbf{s}} \leq \bar{\mathbf{s}}(\square_n) \leq \Theta_n \bar{\mathbf{s}}_*(\square_n) \leq \Theta_n \bar{\mathbf{s}}_*.$$

Therefore the qualitative limit  $\Theta_n \rightarrow 1$  implies that  $\bar{\mathbf{s}} = \bar{\mathbf{s}}_*$ . To obtain a limit for  $\bar{\mathbf{k}}(\square_n)$ , we recall two facts: first, by (2.51) we have that, for every  $m, n \in \mathbb{N}$  with  $m \geq n$ ,

$$\bar{\mathbf{s}}(\square_n) + (\bar{\mathbf{k}}(\square_n) - \bar{\mathbf{k}}(\square_m))^t \bar{\mathbf{s}}_*^{-1}(\square_n) (\bar{\mathbf{k}}(\square_n) - \bar{\mathbf{k}}(\square_m)) \leq \bar{\mathbf{s}}(\square_m) \leq \Theta_m \bar{\mathbf{s}}_*(\square_m) \leq \Theta_m \bar{\mathbf{s}}_*(\square_n).$$

Together with the qualitative limit  $\Theta_m \rightarrow 1$  and  $\bar{\mathbf{s}} = \bar{\mathbf{s}}_*$ , this implies that  $\bar{\mathbf{k}}(\square_n)$  has a limit, which we denote by  $\bar{\mathbf{k}}$ . Second, we recall that, by (2.58), we have

$$|\bar{\mathbf{s}}_*^{-1}(\square_m) (\bar{\mathbf{k}}(\square_m) + \bar{\mathbf{k}}(\square_m)^t)| \leq \Theta_m - 1.$$

Sending  $m \rightarrow \infty$  yields that  $\bar{\mathbf{k}}$  is anti-symmetric. We let  $\bar{\mathbf{A}}$  denote the corresponding  $2d$ -by- $2d$  limiting matrix

$$\bar{\mathbf{A}} := \begin{pmatrix} \bar{\mathbf{s}} + \bar{\mathbf{k}}^t \bar{\mathbf{s}}_*^{-1} \bar{\mathbf{k}} & -\bar{\mathbf{k}}^t \bar{\mathbf{s}}_*^{-1} \\ -\bar{\mathbf{s}}_*^{-1} \bar{\mathbf{k}} & \bar{\mathbf{s}}_*^{-1} \end{pmatrix}. \quad (4.1)$$

It follows that  $\lim_{m \rightarrow \infty} \bar{\mathbf{A}}(\square_m) = \bar{\mathbf{A}}$ . Moreover, due to the ordering  $\bar{\mathbf{A}} \leq \bar{\mathbf{A}}(\square_m)$ , we can apply Lemma 2.1 to obtain

$$0 \leq \bar{\mathbf{A}}(\square_m) - \bar{\mathbf{A}} \leq 4(\Theta_m - 1) \bar{\mathbf{A}}. \quad (4.2)$$

While Theorem 3.1 does give some quantitative information about the rate of convergence of  $\Theta_m - 1$  to zero, and thus, by (4.2), of  $\bar{\mathbf{A}}(\square_m) - \bar{\mathbf{A}}$  to zero, this rate is not particularly useful because it is very slow. The purpose of this section is to improve the rate to algebraic. At the same time, we will obtain *quenched* estimates for the difference between the random matrix  $\mathbf{A}(\square_m)$  and  $\bar{\mathbf{A}}$ . These are stated in the following theorem.

**Theorem 4.1.** *There exist  $C(d) < \infty$  and  $c(d) \in (0, 1/2]$  such that, if we define the parameters*

$$\alpha := (\min\{\nu, 1\} - \gamma)(1 - \beta), \quad \text{and} \quad \kappa := \min\{c, \alpha/3\}, \quad (4.3)$$

*then, for every  $m \in \mathbb{N}$ ,*

$$\Theta_m - 1 \leq 3^{-\kappa m} \exp \left( C \log^2(1 + \Theta) \left( \log K_{\Psi_S} + \frac{1}{\alpha^2} \log \left( \frac{\Pi K_{\Psi}}{\alpha} \right) \right) \right). \quad (4.4)$$

*Moreover, for each  $\delta > 0$  and  $\gamma' \in (\gamma, 1)$ , there exists a random variable  $\mathcal{Y}_{\delta, \gamma'}$  satisfying*

$$\mathcal{Y}_{\delta, \gamma'}^{(\nu - \gamma)(1 - \beta)} = \mathcal{O}_{\Psi} \left( (\kappa \delta)^{-d/\kappa} \exp \left( \frac{C \log K_{\Psi}}{\alpha(\gamma' - \gamma)^2} + \frac{C}{\alpha} \log^2(1 + \Theta) \left( \log K_{\Psi_S} + \frac{1}{\alpha^2} \log \left( \frac{\Pi K_{\Psi}}{\alpha} \right) \right) \right) \right), \quad (4.5)$$

*such that, for  $\theta := \frac{1}{8} \min\{\kappa, \gamma' - \gamma\}$  and for every  $m \in \mathbb{N}$ , we have that*

$$\begin{aligned} 3^m &\geq \mathcal{Y}_{\delta, \gamma'} \vee \mathcal{S} \\ \implies \mathbf{A}(z + \square_k) &\leq \left( 1 + \delta 3^{\gamma'(m-k)} \left( \frac{\mathcal{Y}_{\delta, \gamma'} \vee \mathcal{S}}{3^m} \right)^{\theta} \right) \bar{\mathbf{A}}, \quad \forall k \in \mathbb{Z} \cap (-\infty, m], \quad z \in 3^k \mathbb{Z}^d \cap \square_m. \end{aligned} \quad (4.6)$$

**4.1. Iteration from small contrast to homogenization.** The first statement of Theorem 4.1, on the convergence of  $\Theta_m - 1$  to zero, is proved in this subsection. In fact, we present a more general, perturbative statement, given in Proposition 4.2 below, which is independent of the main results of Section 3, and asserts that, if  $\Theta_0 - 1$  is sufficiently small, then the convergence of  $\Theta_m - 1$  is algebraic in the length scale *as well as in the ellipticity ratios and other parameters*. The estimate (4.4) is then reduced to this statement, with the aid of Theorem 3.1 and a renormalization.<sup>10</sup>

This result is of interest beyond its application to the proof of Theorem 4.1, because many problems exhibiting “high ellipticity contrast” can be shown to actually be of small ellipticity contrast—with the notion of ellipticity we are using in this paper.

**Proposition 4.2** (Algebraic convergence rate in small contrast). *There exist constants  $C(d) < \infty$  and  $\sigma_0(d), c(d) \in (0, 1/2]$  such that, for every  $\sigma \in (0, \sigma_0]$ ,*

$$\Theta_0 - 1 \leq \sigma \implies \Theta_m - 1 \leq \min \left\{ \sigma, \frac{C\Pi^2 \max\{K_{\Psi_S}, K_{\Psi}\}^8}{(1 - \gamma)^2} 3^{-\kappa m} \right\}, \quad \forall m \in \mathbb{N}, \quad (4.7)$$

where the exponent  $\kappa > 0$  is given explicitly by

$$\kappa := \min \left\{ c, \frac{1 - \gamma}{2}, \frac{\nu - \gamma}{1 + \nu - \gamma}, \frac{(1 - \beta)(\nu - \gamma)}{\beta + (1 - \beta)(\nu - \gamma)} \right\}. \quad (4.8)$$

The proof of Proposition 4.2 is based on a second iteration, which uses some of the same ingredients as the one in the proof of Theorem 3.1, but takes advantage of the smallness of  $\Theta_0 - 1$  to accelerate the convergence of  $\Theta_m - 1$  to zero. For this iteration to work without picking up constants which depend on  $\Theta$  or  $\Pi$ , we must work with the adapted cubes  $\Diamond_n$  instead of the Euclidean cubes  $\square_n$ . For this reason it is natural to define an analogue of  $\Theta_m$  in terms of the adapted cubes. We denote this by

$$\hat{\Theta}_m := \frac{1}{d} \text{trace} \left( \mathfrak{S}_*^{-1/2}(\Diamond_m) \mathfrak{S}(\Diamond_m) \mathfrak{S}_*^{-1/2}(\Diamond_m) \right). \quad (4.9)$$

For reasons of convenience we also use the trace in (4.9), rather than the spectral norm of the matrices. By subadditivity, the mapping  $m \mapsto \hat{\Theta}_m$  is monotone decreasing.

We begin the proof of Proposition 4.2 with an alternative to Lemma 3.5. In addition to the fact that the estimate is quenched, rather than an estimate for the variance as in (3.28), the statement here is actually very close to (3.28).

There is one significant improvement which should be highlighted. In (3.28), there are three error terms on the right side: the first term is analogous to  $\hat{\Theta}_k - \hat{\Theta}_m$ , up to constant factors, which appears on the right side of (4.11) below. The difference here is that the error is *quadratic*, that is, the term is squared in (4.11) compared to (3.28). (Note that the left side of (3.28) has a square but the left side of (4.11) does not.) This may seem like a subtle and technical point at first glance, but it is the main reason for the accelerated convergence in small contrast. The reader can also consult the arguments of [AK24, Sections 4.2 & 5.1] for more discussion of this point.

**Lemma 4.3** (Fluctuation estimate). *Suppose  $k, n, m \in \mathbb{N}$  satisfy  $k \leq n \leq m$  and*

$$\hat{\Theta}_k - 1 \leq (80d)^{-1}. \quad (4.10)$$

---

<sup>10</sup>This makes it clear that the obstacle to obtaining better quantitative results in high contrast homogenization, even up to power-like dependence in ellipticity, lies in improving the length scale given in Theorem 3.1, rather than any of the arguments which come in this or later sections.

Then we have the estimates

$$|\bar{\mathbf{A}}^{-1/2}(\diamond_m) \bar{\mathbf{A}}(\diamond_k) \bar{\mathbf{A}}^{-1/2}(\diamond_m) - \mathbf{I}_{2d}| \leq 4d(\hat{\Theta}_k - \hat{\Theta}_m) \quad (4.11)$$

and

$$\begin{aligned} & |\bar{\mathbf{A}}^{-1/2}(\diamond_m) \mathbf{A}(\diamond_n) \bar{\mathbf{A}}^{-1/2}(\diamond_m) - \mathbf{I}_{2d}| \\ & \leq 10d(\hat{\Theta}_k - 1) + 4 \left| \sum_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \bar{\mathbf{A}}^{-1/2}(\diamond_m) (\mathbf{A}(z + \diamond_k) - \bar{\mathbf{A}}(\diamond_k)) \bar{\mathbf{A}}^{-1/2}(\diamond_m) \right|. \end{aligned} \quad (4.12)$$

*Proof.* Fix  $n, m, k \in \mathbb{N}$  with  $k \leq n \leq m$ . We slightly deviate from the notation in the previous section, and set, for every  $j \in \mathbb{N}$ ,

$$\bar{\mathbf{A}}_j := \bar{\mathbf{A}}(\diamond_j), \quad \bar{\mathbf{s}}_j := \bar{\mathbf{s}}(\diamond_j), \quad \bar{\mathbf{s}}_{*,j} := \bar{\mathbf{s}}_*(\diamond_j), \quad \bar{\mathbf{k}}_j := \bar{\mathbf{k}}(\diamond_j) \quad \text{and} \quad \bar{\mathbf{b}}_j := \bar{\mathbf{b}}(\diamond_j).$$

We assume without loss of generality that  $\bar{\mathbf{k}}_k$  is symmetric. Otherwise we recenter by subtracting the anti-symmetric part of  $\bar{\mathbf{k}}_k$  from the coefficient field  $\mathbf{a}(\cdot)$ , using the observations from Section 2.3. Note that  $\hat{\Theta}_j$  as well as the matrix ratios in (4.11) and (4.12) are invariant with respect to recenterings of the anti-symmetric part of the coefficient field, as explained in Section 2.3, and therefore so are the assumptions and conclusions of the lemma.

*Step 1.* The proof of (4.11). We argue as in the proof of Lemma 2.1. Denote, for  $\eta \in [0, 1]$ ,

$$\mathbf{G} := \begin{pmatrix} \bar{\mathbf{k}}_k + (1 - \eta)(\bar{\mathbf{k}}_m - \bar{\mathbf{k}}_k) & 0 \\ 0 & \mathbf{I}_d \end{pmatrix}$$

and compute the top left block of the matrix  $\mathbf{G}^t(\bar{\mathbf{A}}_k - \bar{\mathbf{A}}_m)\mathbf{G}$ , which is

$$\begin{aligned} & \text{the top left block of } \mathbf{G}^t(\bar{\mathbf{A}}_k - \bar{\mathbf{A}}_m)\mathbf{G} \\ & = (\bar{\mathbf{s}}_k - \bar{\mathbf{s}}_m) + (1 - \eta)^2(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m)^t \bar{\mathbf{s}}_{*,k}^{-1}(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m) - \eta^2(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m)^t \bar{\mathbf{s}}_{*,m}^{-1}(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m). \end{aligned}$$

The nonnegativity of  $\mathbf{G}^t(\bar{\mathbf{A}}_k - \bar{\mathbf{A}}_m)\mathbf{G}$  implies that the matrix above is also nonnegative, and thus

$$\begin{aligned} \eta^2(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m)^t \bar{\mathbf{s}}_{*,m}^{-1}(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m) & \leq (\bar{\mathbf{s}}_k - \bar{\mathbf{s}}_m) + (1 - \eta)^2(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m)^t \bar{\mathbf{s}}_{*,k}^{-1}(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m) \\ & \leq (\bar{\mathbf{s}}_k - \bar{\mathbf{s}}_m) + |\bar{\mathbf{s}}_{*,m}^{1/2} \bar{\mathbf{s}}_{*,k}^{-1} \bar{\mathbf{s}}_{*,m}^{1/2}| (1 - \eta)^2(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m)^t \bar{\mathbf{s}}_{*,m}^{-1}(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m). \end{aligned}$$

After rearranging this, we get

$$\left( \eta^2 - |\bar{\mathbf{s}}_{*,m}^{1/2} \bar{\mathbf{s}}_{*,k}^{-1} \bar{\mathbf{s}}_{*,m}^{1/2}| (1 - \eta)^2 \right) (\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m)^t \bar{\mathbf{s}}_{*,m}^{-1}(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m) \leq (\bar{\mathbf{s}}_k - \bar{\mathbf{s}}_m).$$

Optimizing in  $\eta$  leads to the choice  $\eta = |\bar{\mathbf{s}}_{*,m}^{1/2} \bar{\mathbf{s}}_{*,k}^{-1} \bar{\mathbf{s}}_{*,m}^{1/2}| / |\bar{\mathbf{s}}_{*,m}^{1/2} \bar{\mathbf{s}}_{*,k}^{-1} \bar{\mathbf{s}}_{*,m}^{1/2} - \mathbf{I}_d|$  and we obtain

$$\begin{aligned} (\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m)^t \bar{\mathbf{s}}_{*,m}^{-1}(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m) & \leq |\bar{\mathbf{s}}_{*,m}^{1/2} \bar{\mathbf{s}}_{*,k}^{-1} \bar{\mathbf{s}}_{*,m}^{1/2} - \mathbf{I}_d| (\bar{\mathbf{s}}_k - \bar{\mathbf{s}}_m) \\ & \leq |\bar{\mathbf{s}}_{*,m}^{1/2} \bar{\mathbf{s}}_{*,k}^{-1} \bar{\mathbf{s}}_{*,m}^{1/2} - \mathbf{I}_d| |\bar{\mathbf{s}}_m^{-1/2} \bar{\mathbf{s}}_k \bar{\mathbf{s}}_m^{-1/2} - \mathbf{I}_d| \bar{\mathbf{s}}_m. \end{aligned} \quad (4.13)$$

Observe furthermore that

$$\begin{aligned} d(\hat{\Theta}_k - \hat{\Theta}_m) & = \text{trace}(\bar{\mathbf{s}}_k \bar{\mathbf{s}}_{*,k}^{-1} - \bar{\mathbf{s}}_m \bar{\mathbf{s}}_{*,m}^{-1}) = \text{trace} \left( (\bar{\mathbf{s}}_k - \bar{\mathbf{s}}_m) \bar{\mathbf{s}}_{*,k}^{-1} + \bar{\mathbf{s}}_m (\bar{\mathbf{s}}_{*,k}^{-1} - \bar{\mathbf{s}}_{*,m}^{-1}) \right) \\ & \geq \text{trace} \left( (\bar{\mathbf{s}}_k - \bar{\mathbf{s}}_m) \bar{\mathbf{s}}_m^{-1} + \bar{\mathbf{s}}_{*,m} (\bar{\mathbf{s}}_{*,k}^{-1} - \bar{\mathbf{s}}_{*,m}^{-1}) \right) \\ & = \text{trace} \left( \bar{\mathbf{s}}_k \bar{\mathbf{s}}_m^{-1} + \bar{\mathbf{s}}_{*,k}^{-1} \bar{\mathbf{s}}_{*,m} \right) - 2d \\ & \geq (|\bar{\mathbf{s}}_m^{-1/2} \bar{\mathbf{s}}_k \bar{\mathbf{s}}_m^{-1/2} - \mathbf{I}_d| + |\bar{\mathbf{s}}_{*,m}^{1/2} \bar{\mathbf{s}}_{*,k}^{-1} \bar{\mathbf{s}}_{*,m}^{1/2} - \mathbf{I}_d|). \end{aligned}$$

Next we take  $\eta = 0$  and notice that then we have

$$\mathbf{G}^t \bar{\mathbf{A}}_m \mathbf{G} = \begin{pmatrix} \bar{\mathbf{s}}_m & 0 \\ 0 & \bar{\mathbf{s}}_{*,m}^{-1} \end{pmatrix} \quad \text{and} \quad \mathbf{G}^t \bar{\mathbf{A}}_k \mathbf{G} = \begin{pmatrix} \bar{\mathbf{s}}_k + (\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m)^t \bar{\mathbf{s}}_{*,k}^{-1} (\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m) & -(\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m)^t \bar{\mathbf{s}}_{*,k}^{-1} \\ -\bar{\mathbf{s}}_{*,k}^{-1} (\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m) & \bar{\mathbf{s}}_{*,k}^{-1} \end{pmatrix}.$$

We therefore compute, using (2.49) and the assumption in (4.10) that  $d(\hat{\Theta}_k - \hat{\Theta}_m) \leq 80^{-1}$ ,

$$\begin{aligned} |\bar{\mathbf{A}}_m^{-1/2} \bar{\mathbf{A}}_k \bar{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}| &= |(\mathbf{G}^t \bar{\mathbf{A}}_m \mathbf{G})^{-1/2} (\mathbf{G}^t \bar{\mathbf{A}}_k \mathbf{G}) (\mathbf{G}^t \bar{\mathbf{A}}_m \mathbf{G})^{-1/2} - \mathbf{I}_{2d}| \\ &\leq |\bar{\mathbf{s}}_m^{-1/2} \bar{\mathbf{s}}_k \bar{\mathbf{s}}_m^{-1/2} - \mathbf{I}_d| + |\bar{\mathbf{s}}_{*,m}^{-1/2} (\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m) \bar{\mathbf{s}}_m^{-1/2}|^2 \\ &\quad + |\bar{\mathbf{s}}_{*,m}^{-1/2} (\bar{\mathbf{k}}_k - \bar{\mathbf{k}}_m) \bar{\mathbf{s}}_m^{-1/2}| |\bar{\mathbf{s}}_{*,k}^{-1} \bar{\mathbf{s}}_{*,m}|^{1/2} + |\bar{\mathbf{s}}_{*,m}^{1/2} \bar{\mathbf{s}}_{*,k}^{-1} \bar{\mathbf{s}}_{*,m}^{1/2} - \mathbf{I}_d| \\ &\leq 4d(\hat{\Theta}_k - \hat{\Theta}_m). \end{aligned}$$

This completes the proof of (4.11).

*Step 2.* We show (4.12). By the triangle inequality, we have that

$$\begin{aligned} |\bar{\mathbf{A}}_m^{-1/2} \mathbf{A}(\diamond_n) \bar{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}| &\leq \left| \bar{\mathbf{A}}_m^{-1/2} \left( \mathbf{A}(\diamond_n) - \sum_z \mathbf{A}(z + \diamond_k) \right) \bar{\mathbf{A}}_m^{-1/2} \right| \\ &\quad + \left| \bar{\mathbf{A}}_m^{-1/2} \sum_z (\mathbf{A}(z + \diamond_k) - \bar{\mathbf{A}}_k) \bar{\mathbf{A}}_m^{-1/2} \right| \\ &\quad + |\bar{\mathbf{A}}_m^{-1/2} \bar{\mathbf{A}}_k \bar{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}|. \end{aligned} \quad (4.14)$$

To lighten the notation, we will drop the index set in the sums over  $z$  in this step, which is in every instance over  $z \in 3^k \mathbb{L}_0 \cap \diamond_n$ . The last term can be estimated using (4.11) and the second last term on the right is the second term on the right in (4.12). We thus focus on estimating the first term on the right side of (4.14).

Fix any matrix  $\mathbf{B} \in \mathbb{R}_{\text{sym}}^{2d \times 2d}$  and consider the following string of inequalities:

$$\begin{aligned} \mathbf{A}(\diamond_n) &\leq \sum_z \mathbf{A}(z + \diamond_k) \\ &= \sum_z \mathbf{A}_*(z + \diamond_k) + \sum_z (\mathbf{A} - \mathbf{A}_*)(z + \diamond_k) \\ &\leq \left( \sum_z \mathbf{A}_*^{-1}(z + \diamond_k) \right)^{-1} + \sum_z ((\mathbf{A}_* - \mathbf{B}) \mathbf{A}_*^{-1} (\mathbf{A}_* - \mathbf{B}))(z + \diamond_k) + \sum_z (\mathbf{A} - \mathbf{A}_*)(z + \diamond_k) \\ &\leq \mathbf{A}_*(\diamond_n) + \sum_z ((\mathbf{A}_* - \mathbf{B}) \mathbf{A}_*^{-1} (\mathbf{A}_* - \mathbf{B}))(z + \diamond_k) + \sum_z (\mathbf{A} - \mathbf{A}_*)(z + \diamond_k) \\ &\leq \mathbf{A}(\diamond_n) + \sum_z ((\mathbf{A} - \mathbf{B}) + \mathbf{B}(\mathbf{A}_*^{-1} - \mathbf{B}^{-1}) \mathbf{B})(z + \diamond_k). \end{aligned} \quad (4.15)$$

The first and fourth lines in the display above are valid by the subadditivity of  $\mathbf{A}$  and  $\mathbf{A}_*^{-1}$ , respectively. The second line is just a rearrangement. The fifth line is valid by the ordering  $\mathbf{A}_* \leq \mathbf{A}$ . The third line is the key step which says roughly that the sample mean and harmonic mean are separated by at most the sample variance. The inequality we used here can be derived as follows. Denoting the sample mean and the harmonic mean by

$$\mathbf{M} := \sum_z \mathbf{A}_*(z + \diamond_k) \quad \text{and} \quad \mathbf{H} := \left( \sum_z \mathbf{A}_*^{-1}(z + \diamond_k) \right)^{-1},$$

respectively, then we have the following identity, which can be checked by a direct computation: for every symmetric matrix  $\tilde{\mathbf{B}} \in \mathbb{R}_{\text{sym}}^{2d \times 2d}$ ,

$$\mathbf{M} = \mathbf{H} + \sum_z (\mathbf{A}_*(z + \diamond_k) - \tilde{\mathbf{B}}) \mathbf{A}_*^{-1}(z + \diamond_k) (\mathbf{A}_*(z + \diamond_k) - \tilde{\mathbf{B}}) - (\mathbf{H} - \tilde{\mathbf{B}}) \mathbf{H}^{-1} (\mathbf{H} - \tilde{\mathbf{B}}).$$

Discarding the last term yields the inequality in the third line of (4.15), above.

Next, by comparing the first and last lines of (4.15) and inserting  $\mathbf{B} = \overline{\mathbf{A}}_k$ , we obtain that

$$0 \leq \sum_z \mathbf{A}(z + \diamond_k) - \mathbf{A}(\diamond_n) \leq \sum_z \left( (\mathbf{A} - \overline{\mathbf{A}}_k) + \overline{\mathbf{A}}_k (\mathbf{A}_*^{-1} - \overline{\mathbf{A}}_k^{-1}) \overline{\mathbf{A}}_k \right) (z + \diamond_k). \quad (4.16)$$

Let  $h \in \mathbb{N}$  with  $h \geq k_0$ . By (2.64) (applied with  $\mathbf{E}_1 = \overline{\mathbf{A}}_h$ ), the condition  $\hat{\Theta}_h - 1 \leq (80d)^{-1}$  implies

$$|\overline{\mathbf{A}}_h^{-1/2} \overline{\mathbf{A}}_{*,h} \overline{\mathbf{A}}_h^{-1/2} - \mathbf{I}_{2d}| \leq 4d(\hat{\Theta}_h - 1) \leq \frac{1}{20} \quad (4.17)$$

and

$$|\overline{\mathbf{A}}_h^{-1/2} \overline{\mathbf{A}}_{*,h}^{-1} \overline{\mathbf{A}}_h^{1/2} - \mathbf{I}_{2d}| \leq 5d(\hat{\Theta}_h - 1) \leq \frac{1}{16}. \quad (4.18)$$

Therefore, using the fact that  $\mathbf{A}_{*,h}^{-1} = \mathbf{R} \mathbf{A}_h \mathbf{R}$  and  $\overline{\mathbf{A}}_{*,h}^{-1} = \mathbf{R} \overline{\mathbf{A}}_h \mathbf{R}$  with  $\mathbf{R}$  as in (2.32), we get

$$\begin{aligned} & \left| \sum_z \overline{\mathbf{A}}_m^{-1/2} \left( \overline{\mathbf{A}}_k (\mathbf{A}_*^{-1}(z + \diamond_k) - \overline{\mathbf{A}}_k^{-1}) \overline{\mathbf{A}}_k \right) \overline{\mathbf{A}}_m^{-1/2} \right| \\ & \leq |\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_{*,m}^{-1/2}|^2 \left( \left| \sum_z \overline{\mathbf{A}}_m^{-1/2} (\mathbf{A}(z + \diamond_k) - \overline{\mathbf{A}}_k) \overline{\mathbf{A}}_m^{-1/2} \right| + |\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}| \right) \\ & \quad + |\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2}|^2 |\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_{*,m}^{-1} \overline{\mathbf{A}}_m^{1/2} - \mathbf{I}_{2d}| + |\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2}| |\overline{\mathbf{A}}_k^{-1/2} \overline{\mathbf{A}}_m^{-1} \overline{\mathbf{A}}_k^{1/2} - \mathbf{I}_{2d}|. \end{aligned}$$

Indeed, we decompose the matrix on the left using

$$\begin{aligned} & \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \mathbf{A}_*^{-1}(z + \diamond_k) \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2} - \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2} \\ & = \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \mathbf{R} \mathbf{A}(z + \diamond_k) \mathbf{R} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2} - \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2} \\ & = \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_{*,m}^{-1/2} \left( \mathbf{R} \overline{\mathbf{A}}_m^{-1/2} (\mathbf{A}(z + \diamond_k) - \overline{\mathbf{A}}_k) \overline{\mathbf{A}}_m^{-1/2} \mathbf{R} \right) \overline{\mathbf{A}}_{*,m}^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2} \\ & \quad + \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_{*,m}^{-1/2} \left( \mathbf{R} (\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}) \mathbf{R} \right) \overline{\mathbf{A}}_{*,m}^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2} \\ & \quad + \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2} (\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_{*,m}^{-1} \overline{\mathbf{A}}_m^{1/2} - \mathbf{I}_{2d}) \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2} \\ & \quad + \overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k^{-1/2} (\overline{\mathbf{A}}_k^{-1/2} \overline{\mathbf{A}}_m^{-1} \overline{\mathbf{A}}_k^{1/2} - \mathbf{I}_{2d}) \overline{\mathbf{A}}_k^{-1/2} \overline{\mathbf{A}}_m^{-1/2}. \end{aligned}$$

By (4.11), the condition  $\hat{\Theta}_k - 1 \leq (80d)^{-1}$  implies

$$|\overline{\mathbf{A}}_k^{-1/2} \overline{\mathbf{A}}_m^{-1} \overline{\mathbf{A}}_k^{1/2} - \mathbf{I}_{2d}| = |\overline{\mathbf{A}}_m^{-1/2} \overline{\mathbf{A}}_k \overline{\mathbf{A}}_m^{-1/2} - \mathbf{I}_{2d}| \leq 4d(\hat{\Theta}_k - \hat{\Theta}_m) \leq \frac{1}{20},$$

and it follows that

$$\begin{aligned} & \left| \sum_z \overline{\mathbf{A}}_m^{-1/2} \left( \overline{\mathbf{A}}_k (\mathbf{A}_*^{-1}(z + \diamond_k) - \overline{\mathbf{A}}_k^{-1}) \overline{\mathbf{A}}_k \right) \overline{\mathbf{A}}_m^{-1/2} \right| \\ & \leq 6d(\hat{\Theta}_k - 1) + 2 \left| \sum_z \overline{\mathbf{A}}_m^{-1/2} (\mathbf{A}(z + \diamond_k) - \overline{\mathbf{A}}_k) \overline{\mathbf{A}}_m^{-1/2} \right|. \end{aligned}$$

Combining this with (4.16), (4.11) and (4.14) yields (4.12), completing the proof.  $\square$



**Lemma 4.4.** Assume that  $\Theta - 1 \leq \sigma \leq 1/10$ . Then there exist  $\sigma_0(d) > 0$  and  $C(d) < \infty$  such that, if  $\sigma \leq \sigma_0$  then, for every  $n \in \mathbb{N}$  satisfying

$$n \geq 2n_0, \quad n_0 := \left\lceil \frac{50}{1-\gamma} \log \frac{CK_{\Psi_S}\Pi}{1-\gamma} \right\rceil, \quad (4.19)$$

we have the estimate

$$\hat{\Theta}_n - 1 \leq C \sum_{k=n_0}^n 3^{\frac{1}{2}(k-n)} \left( \mathbb{E} \left[ |\mathbf{E}_0^{-1/2}(\mathbf{A}(\diamond_k) - \overline{\mathbf{A}}(\diamond_k))\mathbf{E}_0^{-1/2}|^2 \right] + (\hat{\Theta}_k - \hat{\Theta}_n) \right) + C3^{-\frac{1}{2}(1-\gamma)n}. \quad (4.20)$$

*Proof.* We denote  $n_1 \in \mathbb{N}$  by

$$n_1 := \left\lceil 2 \log K_{\Psi_S} + \log \Pi + k_0 + A \right\rceil,$$

where  $k_0(d)$  is assumed to be so large that  $3^{k_0}\mathbb{L}_0 \subseteq \mathbb{Z}^d$  (see Section 2.7), and  $A(d) > 0$  is sufficiently large that, by (2.72),

$$\overline{\mathbf{A}}(\square_{n_1}) \leq \frac{201}{200} \mathbf{E}_0.$$

and, consequently, by Lemma 2.10,

$$\overline{\mathbf{A}}(\diamond_{2n_1}) \leq \frac{101}{100} \mathbf{E}_0.$$

Moreover, by Lemmas 2.1 and 2.2 and the assumption  $\sigma \leq 1/10$ , we deduce that, for every  $m \geq 2n_1$ ,

$$|\mathbf{E}_0^{-1/2} \overline{\mathbf{A}}(\diamond_m) \mathbf{E}_0^{-1/2} - \mathbf{I}_{2d}| \leq 1/2. \quad (4.21)$$

Note that  $2n_1 \leq n_0$  with  $n_0$  defined in (4.19), provided the  $C$  in (4.19) is sufficiently large.

Fix  $n \in \mathbb{N}$  with  $n \geq 2n_0$ . We recenter the coefficient field by subtracting the constant anti-symmetric matrix  $\frac{1}{2}(\bar{\mathbf{k}} - \bar{\mathbf{k}}^t)(\diamond_n)$  from  $\mathbf{a}(\cdot)$  and relabel the new matrix field as  $\mathbf{a}(\cdot)$ . This allows us to assume, without loss of generality, that  $\bar{\mathbf{k}}(\diamond_n)$  is symmetric.<sup>11</sup> We therefore have that, for any  $e \in \mathbb{R}^d$ ,

$$\begin{aligned} & \mathbb{E}[J(\diamond_n, \bar{\mathbf{s}}_*^{-1/2}(\diamond_n)e, \bar{\mathbf{s}}_*^{1/2}(\diamond_n)e)] \\ &= \frac{1}{2}e \cdot (\bar{\mathbf{s}}_*^{-1/2} \overline{\mathbf{s}\mathbf{s}}_*^{-1/2} - \mathbf{I}_d)(\diamond_n)e + \frac{1}{2}e \cdot (2\bar{\mathbf{s}}_*^{-1/2} \bar{\mathbf{k}} \bar{\mathbf{s}}_*^{-1/2} + \bar{\mathbf{s}}_*^{-1/2} \bar{\mathbf{k}} \bar{\mathbf{s}}_*^{-1} \bar{\mathbf{k}} \bar{\mathbf{s}}_*^{-1/2})(\diamond_n)e \\ &\geq \frac{1}{2}e \cdot (\bar{\mathbf{s}}_*^{-1/2} \overline{\mathbf{s}\mathbf{s}}_*^{-1/2} - \mathbf{I}_d)(\diamond_n)e + e \cdot \bar{\mathbf{s}}_*^{-1/2} \bar{\mathbf{k}} \bar{\mathbf{s}}_*^{-1/2} e \end{aligned}$$

and, likewise,

$$\mathbb{E}[J^*(\diamond_n, \bar{\mathbf{s}}_*^{-1/2}(\diamond_n)e, \bar{\mathbf{s}}_*^{1/2}(\diamond_n)e)] \geq \frac{1}{2}e \cdot (\bar{\mathbf{s}}_*^{-1/2} \overline{\mathbf{s}\mathbf{s}}_*^{-1/2} - \mathbf{I}_d)(\diamond_n)e - e \cdot \bar{\mathbf{s}}_*^{-1/2} \bar{\mathbf{k}} \bar{\mathbf{s}}_*^{-1/2} e.$$

Consequently, we deduce that

$$\hat{\Theta}_n - 1 \leq C \max_{|e|=1} \left( \mathbb{E}[J(\diamond_n, \bar{\mathbf{s}}_*^{-1/2}(\diamond_n)e, \bar{\mathbf{s}}_*^{1/2}(\diamond_n)e)] + \mathbb{E}[J^*(\diamond_n, \bar{\mathbf{s}}_*^{-1/2}(\diamond_n)e, \bar{\mathbf{s}}_*^{1/2}(\diamond_n)e)] \right). \quad (4.22)$$

---

<sup>11</sup>Note that this recentering may, as we have seen above in (3.23), change the value of  $\Pi$  by at most a factor of 100. Because this factor of 100 can be absorbed into the constant  $C$  in the statement of the proposition, we will ignore this issue in the argument.

We fix a unit direction  $e \in \mathbb{R}^d$  with  $|e| = 1$  and set, for the remainder of the argument,

$$p := \bar{\mathbf{s}}_*^{-1/2}(\diamond_n)e \quad \text{and} \quad q := \bar{\mathbf{s}}_*^{1/2}(\diamond_n)e. \quad (4.23)$$

Having fixed  $(p, q)$ , we will denote  $v_m := v(\cdot, \diamond_m, p, q)$  for short, for every  $m \in \mathbb{N}$ .

Applying (3.65) with  $(P, Q) = (0, 0)$  and  $\varepsilon = 1$ , we obtain the existence of  $C(d) < \infty$  such that

$$\begin{aligned} \mathbb{E}[J(\diamond_n, p, q)] &\leq C3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \begin{pmatrix} \nabla v_n \\ \mathbf{a} \nabla v_n \end{pmatrix} \right]_{\bar{\mathbf{B}}_{2,1}^{-1/2}(\diamond_n)}^2 \right] + 100(\mathbb{E}[J(\diamond_{n-4}, p, q)] - \mathbb{E}[J(\diamond_n, p, q)]) \\ &\leq C3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \begin{pmatrix} \nabla v_n \\ \mathbf{a} \nabla v_n \end{pmatrix} \right]_{\bar{\mathbf{B}}_{2,1}^{-1/2}(\diamond_n)}^2 \right] + C(\hat{\Theta}_{n-4} - \hat{\Theta}_n). \end{aligned} \quad (4.24)$$

To estimate the first term on the right side, we need to apply Lemma 2.14. To prepare, we make some preliminary estimates. First, by (2.23) and (2.58) (applied for  $\mathbf{E}_1 = \bar{\mathbf{A}}(\diamond_n)$ ), we have

$$\begin{aligned} \left| \bar{\mathbf{A}}^{1/2}(\diamond_n) \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2 &\leq 2(|\bar{\mathbf{s}}_*^{-1/2} \bar{\mathbf{b}} \bar{\mathbf{s}}_*^{-1/2}| \vee 1)(\diamond_n) \\ &\leq 2 + |\bar{\mathbf{s}}_*^{-1/2} \bar{\mathbf{s}} \bar{\mathbf{s}}_*^{-1/2} - \text{Id}|(\diamond_n) + 2|\bar{\mathbf{s}}_*^{-1/2} \bar{\mathbf{k}} \bar{\mathbf{s}}_*^{-1/2}|^2(\diamond_n) \leq 3. \end{aligned} \quad (4.25)$$

We note next that, by  $\Theta - 1 \leq 1/10$ , we obtain

$$|\mathbf{M}_0^{-1/2} \mathbf{E}_0 \mathbf{M}_0^{-1/2}| \leq 2 \quad (4.26)$$

and, by (4.25) and (4.21),

$$|\mathbf{E}_0^{1/2} \begin{pmatrix} -p \\ q \end{pmatrix}| \leq 4. \quad (4.27)$$

We now square and take the expectation of (2.133). Some of the appearing terms have been already analyzed in the proof of Lemma 3.8, such as (3.82) and (3.83), and they will suffice for our purposes as such. As in (3.80), we obtain the existence of a constant  $C(d) < \infty$  such that, for every  $\varepsilon \in (0, 1)$ ,

$$\begin{aligned} &3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \begin{pmatrix} \nabla v_n - (\nabla v_n)_{\diamond_n} \\ \mathbf{a} \nabla v_n - (\mathbf{a} \nabla v_n)_{\diamond_n} \end{pmatrix} \right]_{\bar{\mathbf{B}}_{2,1}^{-1/2}(\diamond_n)}^2 \right] \\ &\leq \frac{C}{\varepsilon} \sum_{k=n_0}^n 3^{(1-\varepsilon)(k-n)} \left( \mathbb{E} \left[ |\mathbf{E}_0^{-1/2}(\mathbf{A}(\diamond_k) - \bar{\mathbf{A}}(\diamond_k)) \mathbf{E}_0^{-1/2}|^2 \right] + |\mathbf{E}_0^{-1/2}(\bar{\mathbf{A}}(\diamond_n) - \bar{\mathbf{A}}(\diamond_k)) \mathbf{E}_0^{-1/2}| \right) \\ &\quad + \frac{CK_{\Psi_S}^6 \Pi^3}{(1-\gamma)^4} 3^{-3n} + \frac{CK_{\Psi_S} \Pi}{(1-\gamma)^2} 3^{-(1-\gamma)(n-n_0)}. \end{aligned} \quad (4.28)$$

To obtain (4.28), we also used (4.21), which gives us  $|\mathbf{E}_0^{-1}(\bar{\mathbf{A}}(\diamond_n) - \bar{\mathbf{A}}(\diamond_k))| \leq 1$ , and thus, by the triangle inequality,

$$\begin{aligned} \mathbb{E} \left[ |\mathbf{E}_0^{-1/2}(\mathbf{A}(\diamond_k) - \mathbf{A}(\diamond_n)) \mathbf{E}_0^{-1/2}|^2 \right] &\leq 3 \mathbb{E} \left[ |\mathbf{E}_0^{-1/2}(\mathbf{A}(\diamond_k) - \bar{\mathbf{A}}(\diamond_k)) \mathbf{E}_0^{-1/2}|^2 \right] \\ &\quad + 3 \mathbb{E} \left[ |\mathbf{E}_0^{-1/2}(\mathbf{A}(\diamond_n) - \bar{\mathbf{A}}(\diamond_n)) \mathbf{E}_0^{-1/2}|^2 \right] \\ &\quad + 3 |\mathbf{E}_0^{-1/2}(\bar{\mathbf{A}}(\diamond_k) - \bar{\mathbf{A}}(\diamond_n)) \mathbf{E}_0^{-1/2}|. \end{aligned}$$

We will take  $\varepsilon = 1/2$  in (4.28). The two terms on the last line of (4.28) can be brutally estimated using (4.19), as

$$\frac{K_{\Psi_S}^6 \Pi^3}{(1-\gamma)^4} 3^{-2n} + \frac{K_{\Psi_S} \Pi}{(1-\gamma)^2} 3^{-(1-\gamma)(n-n_0)} \leq 3^{-\frac{1}{2}(1-\gamma)n}. \quad (4.29)$$

The second term on the right side can be estimated by

$$\sum_{k=n_0}^n 3^{(1-\varepsilon)(k-n)} |\mathbf{E}_0^{-1/2}(\overline{\mathbf{A}}(\diamond_n) - \overline{\mathbf{A}}(\diamond_k)) \mathbf{E}_0^{-1/2}| \leq C \sum_{k=n_0}^n 3^{\frac{1}{2}(k-n)} (\hat{\Theta}_k - \hat{\Theta}_n). \quad (4.30)$$

Finally, we can remove the constant from the left side of (4.28) using the choice of  $p$  and  $q$ , which yield, in view of (2.31),

$$\begin{pmatrix} \overline{\mathbf{s}}_*^{1/2}(\diamond_n) \mathbb{E}[(\nabla v_n)_{\diamond_n}] \\ \overline{\mathbf{s}}_*^{-1/2}(\diamond_n) \mathbb{E}[(\mathbf{a} \nabla v_n)_{\diamond_n}] \end{pmatrix} = \begin{pmatrix} (\overline{\mathbf{s}}_*^{-1/2} \overline{\mathbf{k}} \mathbf{s}_*^{-1/2})(\diamond_n) e \\ -(\overline{\mathbf{s}}_*^{-1/2} \overline{\mathbf{k}} \mathbf{s}_*^{-1} \overline{\mathbf{k}} \mathbf{s}_*^{-1/2} + \overline{\mathbf{s}}_*^{-1/2} \overline{\mathbf{k}} \mathbf{s}_*^{-1/2})(\diamond_n) e \end{pmatrix}$$

It follows from this, (2.33), (2.58) and (4.25) that

$$\mathbb{E} \left[ \left| \mathbf{M}_0^{1/2} \begin{pmatrix} (\nabla v_n)_{\diamond_n} \\ (\mathbf{a} \nabla v_n)_{\diamond_n} \end{pmatrix} \right|^2 \right] \leq C(\hat{\Theta}_n - 1)^2 + C \mathbb{E} \left[ |\mathbf{E}_0^{-1/2}(\mathbf{A}(\diamond_n) - \overline{\mathbf{A}}(\diamond_n)) \mathbf{E}_0^{-1/2}|^2 \right]$$

and therefore

$$\begin{aligned} & \mathbb{E} \left[ \left| \mathbf{M}_0^{1/2} \begin{pmatrix} (\nabla v_n)_{\diamond_n} \\ (\mathbf{a} \nabla v_n)_{\diamond_n} \end{pmatrix} \right|^2 \right] \\ & \leq 2 \mathbb{E} \left[ \left| \mathbf{M}_0^{1/2} \begin{pmatrix} (\nabla v_n)_{\diamond_n} - (\nabla v_n)_{\diamond_n} \\ (\mathbf{a} \nabla v_n)_{\diamond_n} - (\mathbf{a} \nabla v_n)_{\diamond_n} \end{pmatrix} \right|^2 \right] + 2 \mathbb{E} \left[ \left| \mathbf{M}_0^{1/2} \begin{pmatrix} (\nabla v_n)_{\diamond_n} \\ (\mathbf{a} \nabla v_n)_{\diamond_n} \end{pmatrix} \right|^2 \right] \\ & \leq C \mathbb{E} \left[ |\mathbf{E}_0^{-1/2}(\mathbf{A}(\diamond_n) - \overline{\mathbf{A}}(\diamond_n)) \mathbf{E}_0^{-1/2}|^2 \right] + C(\hat{\Theta}_n - 1)^2 + C 3^{-\frac{1}{2}(1-\gamma)n}. \end{aligned}$$

Combining the above, we obtain

$$\begin{aligned} \mathbb{E}[J(\diamond_n, p, q)] & \leq C \sum_{k=n_0}^n 3^{\frac{1}{2}(k-n)} \left( \mathbb{E} \left[ |\mathbf{E}_0^{-1/2}(\mathbf{A}(\diamond_k) - \overline{\mathbf{A}}(\diamond_k)) \mathbf{E}_0^{-1/2}|^2 \right] + (\hat{\Theta}_k - \hat{\Theta}_n) \right) \\ & \quad + C(\hat{\Theta}_n - 1)^2 + C 3^{-\frac{1}{2}(1-\gamma)n}. \end{aligned}$$

By symmetry, we obtain the same bound for  $J^*$  in place of  $J$ . By (4.22), we obtain (4.20) with an additional term of  $C(\hat{\Theta}_n - 1)^2$  on the right side. This term can however now be absorbed by the left side if we require  $\sigma_0$  to be sufficiently small. The proof is complete.  $\square$

We are now ready to prove the proposition.

*Proof of Proposition 4.2.* We combine the previous two lemmas, and then iterate the result. Using Lemma 4.3 and (4.21), we have that, for every  $k_1 \geq n_0$ ,

$$\begin{aligned} & \mathbb{E} \left[ |\mathbf{E}_0^{-1/2}(\mathbf{A}(\diamond_k) - \overline{\mathbf{A}}(\diamond_k)) \mathbf{E}_0^{-1/2}|^2 \right] \\ & \leq C(\hat{\Theta}_k - 1)^2 + C \mathbb{E} \left[ \left| \mathbf{E}_0^{-1/2} \sum_{z \in 3^l \mathbb{L}_0 \cap \diamond_k} (\mathbf{A}(z + \diamond_l) - \overline{\mathbf{A}}(\diamond_l)) \mathbf{E}_0^{-1/2} \right|^2 \right]. \end{aligned} \quad (4.31)$$

By (2.104), we have that, for every  $k, l \in \mathbb{N}$  with  $\beta k < l \leq k$ ,

$$\mathbb{E} \left[ \left| \mathbf{E}_0^{-1/2} \sum_{z \in 3^l \mathbb{L}_0 \cap \diamond_k} (\mathbf{A}(z + \diamond_l) - \overline{\mathbf{A}}(\diamond_l)) \mathbf{E}_0^{-1/2} \right|^2 \right] \leq \frac{C \Pi^2 K_{\Psi_S}^8}{(1 - \gamma)^2} 3^{2(\gamma(k-l)-k)} + C \Pi K_{\Psi}^8 3^{-2(\nu-\gamma)(k-l)}. \quad (4.32)$$

For each  $k \in \mathbb{N}$ , we choose  $l_k$  by

$$l = l_k := \left\lceil \frac{\nu - \gamma}{\kappa + \nu - \gamma} k \right\rceil + 1 \implies \kappa l_k \geq (\nu - \gamma)(k - l_k) \geq \kappa(l_k - 2). \quad (4.33)$$

By the definition of  $\kappa$  in (4.8),

$$\begin{aligned} \kappa &\leq \min \left\{ \frac{1}{2}(1 - \gamma), \nu - \gamma, \frac{1 - \beta}{\beta}(\nu - \gamma) \right\} \\ &\implies 2\kappa l_k \geq \kappa k, \quad l_k > \beta k \quad \text{and} \quad 2(k - \gamma(k - l)) \geq \kappa k - 4. \end{aligned} \quad (4.34)$$

We deduce that

$$\mathbb{E} \left[ \left| \mathbf{E}_0^{-1/2} (\mathbf{A}(\diamond_k) - \overline{\mathbf{A}}(\diamond_k)) \mathbf{E}_0^{-1/2} \right|^2 \right] \leq C(\hat{\Theta}_k - 1)^2 + CH 3^{-\kappa k},$$

where we set

$$H := \frac{C \Pi^2 \max\{K_{\Psi_S}, K_{\Psi}\}^8}{(1 - \gamma)^2}.$$

Inserting this estimate into the result of Lemma 4.4 yields

$$\begin{aligned} \hat{\Theta}_n - 1 &\leq C \sum_{k=n_0}^n 3^{\frac{1}{2}(k-n)} \left( (\hat{\Theta}_k - 1)^2 + H 3^{-\kappa k} + (\hat{\Theta}_k - \hat{\Theta}_n) \right) + C 3^{-\frac{1}{2}(1-\gamma)n} \\ &\leq C \sum_{k=n_0}^n 3^{\frac{1}{2}(k-n)} \left( (\hat{\Theta}_k - 1)^2 + (\hat{\Theta}_k - \hat{\Theta}_n) \right) + CH 3^{-\kappa n}. \end{aligned} \quad (4.35)$$

This inequality can now be iterated to obtain the result (see for instance [AK24, Lemma 4.8]).  $\square$

We next use Theorem 3.1 and the renormalization lemma (Proposition 2.6) to remove the restriction  $\Theta_0 - 1 \leq \sigma$  from the statement of Proposition 4.2. This yields the first statement of Theorem 4.1.

*Proof of the first statement of Theorem 4.1.* Using Theorem 3.1, we find a scale  $n_0$  with

$$n_0 \leq C \left( \log K_{\Psi_S} + \frac{1}{\alpha^2} \log \left( \frac{\Pi K_{\Psi}}{\alpha} \right) \right) \log^2(1 + \Theta)$$

such that, if we let  $\sigma_0(d)$  be the constant given in the statement of Proposition 4.2, then

$$\Theta_{n_0} \leq 1 + \sigma_0.$$

Let  $l_0$  be defined as in (3.87)–(3.88) and let  $m_0 := n_0 + 2l_0 + \lceil \log K_{\Psi} \rceil$ . As in the argument in the beginning of the proof of Theorem 3.1, we may apply Proposition 2.6 to obtain that the pushforward

probability measure  $\mathbb{P}_{m_0-2l_0}$ , defined in (2.83), satisfies the assumptions (P1), (P2) and (P3) with the new parameters

$$\begin{cases} \mathbf{E}_{\text{new}} := 2\overline{\mathbf{A}}(\square_{m_0-2l_0}) \leq 2\overline{\mathbf{A}}(\square_{n_0}), \\ \gamma_{\text{new}} := \frac{1}{2}(\min\{1, \nu\} + \gamma) \\ K_{\Psi, \text{new}} := K_{\Psi} \\ K_{\Psi_S, \text{new}} := K_{\Psi_S}^*, \\ \Theta_{\text{new}} := 4\Theta_{m_0-2l_0} \leq 4\Theta_{n_0} \leq 4 + 4\sigma_0 \leq 8, \\ \Pi_{\text{new}} \leq \Pi^*. \end{cases} \quad (4.36)$$

Observe that  $\Theta_0$  for  $\mathbb{P}_{m_0-2l_0}$  is the same as  $\Theta_{m_0-2l_0}$  for  $\mathbb{P}$ , and we have  $\Theta_{m_0-2l_0} \leq \Theta_{n_0} \leq 1 + \sigma_0$ . We may therefore apply Proposition 4.2 with  $\mathbb{P}_{m_0-2l_0}$  in place of  $\mathbb{P}$ , to obtain that, for every  $n \in \mathbb{N}$ ,

$$\Theta_{n_0 + \lceil \log K_{\Psi} \rceil + n} = \Theta_{m_0-2l_0+n} \leq \frac{C(\Pi^*)^2 \max\{K_{\Psi_S}^*, K_{\Psi}\}^8}{(1 - \gamma^*)^2} 3^{-\kappa n} \leq \frac{C\Pi^2}{\alpha^2} \max\{K_{\Psi_S}, K_{\Psi}^{\lceil 1/\alpha \rceil}\}^8 3^{-\kappa n},$$

where  $\kappa$  is as defined in (4.8) and  $\alpha = (\min\{\nu, 1\} - \gamma)(1 - \beta)$ . Reindexing this, we obtain

$$\Theta_m \leq \frac{C\Pi^2}{\alpha^2} \max\{K_{\Psi_S}, K_{\Psi}^{\lceil 1/\alpha \rceil}\}^8 3^{-\kappa(m-n_0-\lceil \log K_{\Psi} \rceil)}, \quad \forall m \geq n_0 + \lceil \log K_{\Psi} \rceil,$$

We next observe that

$$\frac{C\Pi^2 \max\{K_{\Psi_S}, K_{\Psi}^{\lceil 1/\alpha \rceil}\}^8}{\alpha^2} 3^{\kappa(n_0 + \lceil \log K_{\Psi} \rceil)} \leq \exp\left(C\left(\log K_{\Psi_S} + \frac{1}{\alpha^2} \log\left(\frac{\Pi K_{\Psi}}{\alpha}\right)\right) \log^2(1 + \Theta)\right).$$

Substituting this into the previous display gives (4.4) for every  $m \geq n_0 + \lceil \log K_{\Psi} \rceil$ . For  $m \in \mathbb{N}$  with  $m \leq n_0 + \lceil \log K_{\Psi} \rceil$ , the inequality is obtained trivially from  $\Theta_m \leq \Pi$ , after possibly enlarging the constant  $C$ . This completes the proof of the corollary.  $\square$

**4.2. Quenched convergence of the coarse-grained matrices.** We now turn to the task of proving the second statement of Theorem 4.1, that is, obtaining the (quenched) convergence of the random variables  $\mathbf{A}(\square_m)$  to the constant matrix  $\overline{\mathbf{A}}$ , not just their means. This will be obtained by using the first statement of Theorem 4.1, proved above, and a subadditivity argument which is more or less standard (for instance, see [AK24, Lemma 4.12] for essentially the same argument in a simpler situation, without the explicit constants).

The idea is to estimate each coarse-grained matrix  $\mathbf{A}(z + \square_k)$ , for  $z \in 3^k \mathbb{Z}^d \cap \square_m$ , using subadditivity as follows:

$$\mathbf{A}(z + \square_k) \leq \sum_{z' \in z + 3^l \mathbb{Z}^d \cap \square_k} \mathbf{A}(z' + \square_l) \leq \overline{\mathbf{A}}(\square_l) + \sum_{z' \in z + 3^l \mathbb{Z}^d \cap \square_k} (\mathbf{A}(z' + \square_l) - \overline{\mathbf{A}}(\square_l)). \quad (4.37)$$

Here  $l$  represents a scale that is smaller than  $k$ , and needs to be chosen to balance the two terms on the right side. The first term, namely  $\overline{\mathbf{A}}(\square_l)$ , is upper bounded by  $(1 + 4(\Theta_l - 1))\overline{\mathbf{A}}$ , by (4.2), and so we already have an explicit estimate for it in (4.4); this error becomes smaller for larger choices of  $l$ . The second term is a sum of random variables to which we will apply the mixing condition (P3). In fact, the needed estimate is already found in Lemma 2.5. It becomes *larger* for larger values of  $l$ , because smaller values of  $l$  create more subcubes, leading to greater stochastic cancellations. The argument is only tedious due to the need to carefully select  $l$  in terms of  $k$  and  $m$  to balance these errors, which of course involve many of the parameters.

*Proof of the second statement of Theorem 4.1.* Fix  $\delta \in (0, 1]$  and  $\gamma' \in (\gamma, 1)$ . The goal is to identify a random scale  $\mathcal{Y} = \mathcal{Y}_{\delta, \gamma'}$  such that

$$3^m \geq \mathcal{Y} \vee \mathcal{S} \implies (4.6). \quad (4.38)$$

We  $\mu$  denote be the exponent

$$\mu := (\nu - \gamma)(1 - \beta), \quad (4.39)$$

which will be related to the stochastic integrability exponent  $\mathcal{Y}$ . For convenience, we also let  $\Upsilon$  be the constant defined by

$$\Upsilon := \exp\left(C\left(\log K_{\Psi_S} + \frac{1}{\alpha} \log\left(\frac{\Pi K_{\Psi}}{\alpha}\right)\right) \log^2(1 + \Theta)\right), \quad (4.40)$$

where  $C(d) < \infty$  is a large constant, which is no smaller than  $C_{(4.4)}$ , and which may be further inflated in the course of this argument. We fix parameters  $h, n_1 \in \mathbb{N}$  which are given explicitly below in (4.43) and depend only on  $d$  and the other parameters.

Let  $n \in \mathbb{N}$  be such that  $n \geq n_1 + h$ . As in the discussion around (4.37) above, we use subadditivity, (4.2) and Lemma 2.5, to find that, for every  $k, l, m \in \mathbb{N}$  with  $\beta k < l \leq k \leq m$  and  $z \in 3^k \mathbb{Z}^d \cap \square_m$ ,

$$\begin{aligned} \bar{\mathbf{A}}^{-1/2} \mathbf{A}(z + \square_k) \bar{\mathbf{A}}^{-1/2} \mathbf{1}_{\{S \leq 3^m\}} &\leq \sum_{z' \in z + 3^l \mathbb{Z}^d \cap \square_k} \bar{\mathbf{A}}^{-1/2} \mathbf{A}(z' + \square_l) \bar{\mathbf{A}}^{-1/2} \mathbf{1}_{\{S \leq 3^m\}} \\ &\leq \bar{\mathbf{A}}^{-1/2} \bar{\mathbf{A}}(\square_l) \bar{\mathbf{A}}^{-1/2} \mathbf{1}_{\{S \leq 3^m\}} + \mathcal{O}_{\Psi}(C 3^{\gamma(m-l)} 3^{-\nu(k-l)} \mathbf{I}_{2d}) \\ &\leq (1 + 4(\Theta_l - 1)) \mathbf{I}_{2d} \mathbf{1}_{\{S \leq 3^m\}} + \mathcal{O}_{\Psi}(C 3^{\gamma(m-k)} 3^{-(\nu-\gamma)(k-l)} \mathbf{I}_{2d}). \end{aligned} \quad (4.41)$$

We apply the previous display in the case that  $k \geq n - h$  and with the parameter  $l = \lfloor l_k \rfloor$ , where

$$l_k := \beta k + \frac{1 - \beta}{2}(k - n + h)_+ + (1 - \beta)n_1.$$

It is clear that  $\lfloor l_k \rfloor > \beta k$ . We need to check that  $\lfloor l_k \rfloor$  is smaller or equal to  $k$ . To see this, we use  $k \geq n - h$  and  $n \geq n_1 + h$  to find that

$$l_k \leq \beta k - \frac{1 - \beta}{2}(k + n - h) + (1 - \beta)n_1 \leq k - (1 - \beta)(n - h) + (1 - \beta)n_1 \leq k.$$

The first term on the right side of (4.41) is estimated using (4.4). In view of the choice of  $l = \lfloor l_k \rfloor$ , we obtain

$$4(\Theta_l - 1) \leq \Upsilon 3^{-\kappa \beta k - \frac{\kappa}{2}(1 - \beta)(k - n + h) - \kappa(1 - \beta)n_1}.$$

Before estimating the second term, we note that

$$(\nu - \gamma)(k - l_k) = \mu k + \frac{\mu}{2}(k - n + h) - \mu n_1 \geq \mu n + \frac{\mu}{2}(k - n + h) - \mu(n_1 + h).$$

Substituting this into the second term of (4.41) yields that

$$3^{\gamma(m-k)} 3^{-(\nu-\gamma)(k-l)} \leq C 3^{\mu(n_1+h)} 3^{\gamma(m-k)} 3^{-\frac{\mu}{2}(k-n+h)} 3^{-\mu n}.$$

Putting the above together yields, for every  $k \geq n - h$ ,

$$\begin{aligned} \bar{\mathbf{A}}^{-1/2} (\mathbf{A}(z + \square_k) - \bar{\mathbf{A}}) \bar{\mathbf{A}}^{-1/2} \mathbf{1}_{\{S \leq 3^m\}} \\ \leq \Upsilon 3^{-\kappa \beta k - \frac{\kappa}{2}(1 - \beta)(k - n + h) - \kappa(1 - \beta)n_1} \mathbf{I}_{2d} + \mathcal{O}_{\Psi}(C 3^{\mu(n_1+h)} 3^{\gamma(m-k)} 3^{-\frac{\mu}{2}(k-n+h)} 3^{-\mu n} \mathbf{I}_{2d}). \end{aligned} \quad (4.42)$$

This gives us a good estimate in every cube of the form  $z + \square_k$ , but now we need to use it to make a union bound over all subcubes within a larger cube  $\square_m$ . To that end, we introduce the exponent  $\theta := \frac{1}{4} \min\{\kappa, \gamma' - \gamma\}$ , as defined in the statement of the theorem, and define the composite quantity

$$\mathbf{F}_n := \frac{6}{\delta} \sum_{m=n}^{\infty} 3^{\theta(m-n)} \sum_{k=-\infty}^m 3^{-\gamma'(m-k)} \max_{z \in 3^k \mathbb{Z}^d \cap \square_m} \left| (\overline{\mathbf{A}}^{-1/2} (\mathbf{A}(z + \square_k) - \overline{\mathbf{A}}) \overline{\mathbf{A}}^{-1/2})_+ \right| \mathbf{1}_{\{S \leq 3^m\}}.$$

Here  $(A)_+$  denotes the nonnegative part of a symmetric matrix  $A$ . Notice that  $\mathbf{F}_n$  depends in particular on the parameters  $(\delta, \gamma')$ . We cut the summation in  $k$  for large and small  $k$ . For large  $k$  we use a union bound together with (C.4), (C.6) and (4.42) to obtain

$$\begin{aligned} \mathbf{F}_n^+ &:= \frac{6}{\delta} \sum_{m=n}^{\infty} 3^{\theta(m-n)} \sum_{k=n-h}^m 3^{-\gamma'(m-k)} \max_{z \in 3^k \mathbb{Z}^d \cap \square_m} \left| (\overline{\mathbf{A}}^{-1/2} (\mathbf{A}(z + \square_k) - \overline{\mathbf{A}}) \overline{\mathbf{A}}^{-1/2})_+ \right| \mathbf{1}_{\{S \leq 3^m\}} \\ &\leq \frac{\Upsilon}{\delta} 3^{-\kappa(1-\beta)n_1} \sum_{m=n}^{\infty} 3^{\theta(m-n)} \sum_{k=n-h}^m 3^{-\gamma'(m-k)} 3^{-\beta\kappa k - \frac{\kappa}{2}(1-\beta)(k-n+h)} \mathbf{I}_{2d} \\ &\quad + \mathcal{O}_{\Psi} \left( (CK_{\Psi})^{16d^2(\gamma'-\gamma)^{-2}} \delta^{-1} \theta^{-1} \mu^{-1} 3^{\kappa(n_1+h)} 3^{-\mu n} \mathbf{I}_{2d} \right). \end{aligned}$$

We use above the following elementary observation: for random variables  $\{X_k\}_{k=1}^N$  with  $X_k = \mathcal{O}_{\Psi}(a)$ , we have by a union bound and (C.4) that, for every  $t \geq 1$  and  $\sigma \in (0, 1]$ ,

$$\mathbb{P} \left[ \max_{k \in \{1, \dots, N\}} X_k > N^{\sigma} a t \right] \leq \sum_{k=1}^N \mathbb{P}[X_k > N^{\sigma} a t] \leq \frac{N}{\Psi(N^{\sigma} a t)} \leq \frac{1}{\Psi(K_{\Psi}^{-3[\sigma^{-1}]^2} N^{\sigma} a t)}.$$

Since  $\theta \leq \frac{1}{8} \min\{\kappa, \gamma' - \gamma\}$ , the first sum on the right can be estimated as

$$\begin{aligned} &\sum_{m=n}^{\infty} 3^{\theta(m-n)} \sum_{k=n-h}^m 3^{-\gamma'(m-k)} 3^{-\beta\kappa k - \frac{\kappa}{2}(1-\beta)(k-n+h)} \\ &\leq \frac{C}{\kappa + \gamma'} \sum_{m=n}^{\infty} 3^{\theta(m-n)} (3^{-\gamma'(m-n)} + 3^{-\frac{\kappa}{2}(m-n)}) \leq \frac{C}{(\kappa + \gamma')^2}. \end{aligned}$$

It follows that

$$\mathbf{F}_n^+ \leq \frac{\Upsilon 3^{-\kappa(1-\beta)n_1}}{\delta(\kappa + \gamma')^2} \mathbf{I}_{2d} + \mathcal{O}_{\Psi} \left( (CK_{\Psi})^{4d^2(\gamma'-\gamma)^{-2}} \delta^{-1} \theta^{-1} \mu^{-1} 3^{\mu(n_1+h)} 3^{-\mu n} \mathbf{I}_{2d} \right).$$

On the other hand, since  $\theta \leq \frac{1}{8}(\gamma' - \gamma)$ , we have by (P2) that

$$\begin{aligned} \mathbf{F}_n^- &:= \frac{6}{\delta} \sum_{m=n}^{\infty} 3^{\theta(m-n)} \sum_{k=-\infty}^{n-h-1} 3^{-\gamma'(m-k)} \sup_{z \in 3^k \mathbb{Z}^d \cap \square_m} \overline{\mathbf{A}}^{-1/2} (\mathbf{A}(z + \square_k) - \overline{\mathbf{A}}) \overline{\mathbf{A}}^{-1/2} \mathbf{1}_{\{S \leq 3^m\}} \\ &\leq \frac{C}{\delta} \sum_{m=n}^{\infty} 3^{\theta(m-n)} \sum_{k=-\infty}^{n-h-1} 3^{-(\gamma'-\gamma)(m-k)} \mathbf{I}_{2d} \leq \frac{C 3^{-\frac{1}{2}(\gamma'-\gamma)h}}{\delta(\gamma' - \gamma)^2}. \end{aligned}$$

We then choose  $n_1$  and  $h$  to be, respectively, the smallest integers satisfying

$$\frac{\Upsilon 3^{-\kappa(1-\beta)n_1}}{\delta \kappa^2 (\gamma' - \gamma)^2} \leq \frac{1}{4} \quad \text{and} \quad \frac{C 3^{-\frac{1}{2}(\gamma'-\gamma)h}}{\delta (\gamma' - \gamma)^2} \leq \frac{1}{4}. \quad (4.43)$$

As a consequence we deduce that

$$\mathbf{F}_n = \mathbf{F}_n^+ + \mathbf{F}_n^- \leq \frac{1}{2} \mathbf{I}_{2d} + \mathcal{O}_\Psi(T 3^{-\mu n} \mathbf{I}_{2d}) \quad \text{with} \quad T := (CK_\Psi)^{4d^2(\gamma' - \gamma)^{-2}} \left( \frac{\Upsilon}{\kappa \delta} \right)^{d/\kappa}. \quad (4.44)$$

Set

$$\mathcal{Y}_{\delta, \gamma'} = \mathcal{Y} := \sum_{n=n_1+h}^{\infty} 3^n \min \left\{ 1, \sup_{|e| \leq 1} e \cdot (\mathbf{F}_n - \mathbf{I}_{2d}) e \right\}.$$

We find, a.s.,  $n \in \mathbb{N}$  such that  $3^{n-1} < \mathcal{Y} \leq 3^n$ . We have that  $3^n > \mathcal{Y}$  implies  $\mathbf{F}_n \leq 2\mathbf{I}_{2d}$ , and, hence,  $3^{n-1} \leq \mathcal{Y} < 3^n$  implies that, for every  $m \in \mathbb{N}$ ,

$$\sum_{k=-\infty}^m 3^{-\gamma'(m-k)} \max_{z \in 3^k \mathbb{Z}^d \cap \square_m} \overline{\mathbf{A}}^{-1/2} (\mathbf{A}(z + \square_k) - \overline{\mathbf{A}}) \overline{\mathbf{A}}^{-1/2} \mathbf{1}_{\{S \leq 3^m\}} \leq \delta \left( \frac{\mathcal{Y}}{3^m} \right)^\theta \mathbf{I}_{2d}.$$

We therefore deduce that

$$3^m \geq \mathcal{Y} \vee S \implies \sup_{k \in \mathbb{Z} \cap (-\infty, m]} \sup_{z \in 3^k \mathbb{Z}^d \cap \square_m} \mathbf{A}(z + \square_k) \leq \left( 1 + \delta 3^{\gamma'(m-k)} \left( \frac{\mathcal{Y} \vee S}{3^m} \right)^\theta \right) \overline{\mathbf{A}}.$$

The desired integrability of  $\mathcal{Y}$  is a consequence of (4.44). Indeed, by (4.44) and (C.6), we have

$$\mathbb{P}[\mathcal{Y} > 3^n] \leq \sum_{m=n-1}^{\infty} \mathbb{P}[\mathbf{F}_m \not\leq \mathbf{I}_{2d}] \leq \sum_{m=n-1}^{\infty} \left( \Psi(CT 3^{\mu m}) \right)^{-1} \leq \left( \Psi(CK_\Psi^4 T 3^{\mu n}) \right)^{-1}$$

This yields

$$\mathcal{Y}^\mu \leq \mathcal{O}_\Psi(CK_\Psi^4 T),$$

which gives us (4.5). This completes the proof of (4.6).  $\square$

In Section 6, we will need an estimate which is slightly different from the one in (4.6), given in terms of the defined for every  $s \in (0, 1]$ ,  $y \in \mathbb{R}^d$  and  $n \in \mathbb{N}$ :

$$\tilde{\mathcal{E}}_s(y + \diamond_n) := \sum_{k=-\infty}^n 3^{s(k-n)} \max_{z \in y + 3^k \mathbb{L}_0 \cap \diamond_n} \left| (\overline{\mathbf{A}}^{-1/2} (\mathbf{A}(z + \diamond_k) - \overline{\mathbf{A}}) \overline{\mathbf{A}}^{-1/2})_+ \right|^{1/2}. \quad (4.45)$$

We also need a version of (4.45) for *adapted simplexes*, which we will define next. Let  $\mathcal{P}$  denote the set of permutations of  $\{1, \dots, d\}$ . Given any permutation  $\pi \in \mathcal{P}$ , we define, for every  $n \in \mathbb{Z}$  and  $z \in \mathbb{R}^d$ ,

$$\triangle_n^\pi(z) := z + 3^n \mathbf{q}_0 \left\{ (x_1, \dots, x_d) \in \mathbb{R}^d : -\frac{1}{2} < x_{\pi(1)} < x_{\pi(2)} < \dots < x_{\pi(d)} < \frac{1}{2} \right\}.$$

Like cubes, the simplexes are Lipschitz domains and they have the nice property that any triadic simplex  $\triangle_n^\pi(z)$  is the disjoint union (up to a set of Lebesgue measure zero) of  $3^{d(m-n)}$  many simplexes of the form  $\triangle_n^\sigma(z)$ . Moreover, the adapted cube  $\diamond_n(z)$  is the disjoint union of  $\{\triangle_n^\pi(z) : \pi \in \mathcal{P}\}$ . Note that, since  $|\mathcal{P}| = d!$ , this implies  $|\triangle_n^\pi| = 3^n/d!$ . We call  $S_n := \{\triangle_n^\pi(z) : n \in \mathbb{Z}, z \in 3^n \mathbb{L}_0, \triangle_n^\pi(z) \subseteq \diamond_m\}$  the collection of *triadic adapted simplexes* in  $\diamond_m$ . We call  $3^n$  the *side length* of  $\triangle_n^\pi(z)$ . We also denote  $\triangle_n = \triangle_n^\pi$  if  $\pi$  is the trivial permutation. Each simplex has  $d+1$  many extreme points, which we call the *vertices* of the simplex. For  $k, n \in \mathbb{N}$  with  $k < n$  and  $\pi \in \mathcal{P}$ , denote  $\mathcal{Z}_k^\triangle(\triangle_n^\pi) := \{\triangle \in S_k : \triangle \subseteq \triangle_n^\pi\}$ . Analogously to (4.45), set, for  $\triangle \in S_n$ ,

$$\tilde{\mathcal{E}}_s^\triangle(\triangle) := \sum_{k=-\infty}^n 3^{s(k-n)} \max_{\triangle' \in \mathcal{Z}_k^\triangle(\triangle)} \left| (\overline{\mathbf{A}}^{-1/2} (\mathbf{A}(\triangle') - \overline{\mathbf{A}}) \overline{\mathbf{A}}^{-1/2})_+ \right|^{1/2}. \quad (4.46)$$



**Corollary 4.5.** *Let  $s \in (\gamma/2, 1]$ ,  $\delta \in (0, 1]$  and set  $\theta := \frac{1}{32} \min\{\kappa, 2s - \gamma\}$ , where  $\kappa$  is as in (4.3). There exist a constant  $C(d) < \infty$  and a random scale  $\mathcal{Z}_{\delta,s}$  satisfying*

$$\mathcal{Z}_{\delta,s} = \mathcal{O}_\Psi \left( \exp \left( C \left( \alpha^{-1} |\log(\theta\delta)| + \alpha^{-3} \log^2(1 + \Theta) \log(K_\Psi K_{\Psi_S} \Pi) \right) \right) \right), \quad (4.47)$$

with  $\alpha := (\min\{\nu, 1\} - \gamma)(1 - \beta)$ , such that, for every  $m \in \mathbb{N}$ ,

$$\begin{aligned} 3^m &\geq \mathcal{Z}_{\delta,s} \vee \mathcal{S}, \quad n \in \mathbb{Z} \cap (-\infty, m], \quad z \in 3^n \mathbb{L}_0 \cap \Diamond_m, \quad \Delta \in \{\Delta_n^\pi : \pi \in \mathcal{P}\} \\ \implies \quad \tilde{\mathcal{E}}_s(z + \Diamond_n) + \tilde{\mathcal{E}}_s^\Delta(z + \Delta) &\leq \delta 3^{\frac{1}{2}(\gamma+\theta)(m-n)} \left( \frac{\mathcal{Z}_{\delta,s} \vee \mathcal{S}}{3^m} \right)^\theta. \end{aligned} \quad (4.48)$$

*Proof.* Fix  $s \in (\gamma/2, 1]$ ,  $\delta \in (0, 1]$ ,  $\theta := \frac{1}{32} \min\{\kappa, 2s - \gamma\}$  and  $\gamma' = \gamma + \theta$ . Let

$$\delta_1 := c\delta^2(1 - \gamma)(2s - \gamma)^2 \Pi^{-1/2}, \quad (4.49)$$

and let  $\mathcal{Z}_{\delta,s} := \mathcal{Y}_{\delta_1, \gamma'}$  with  $\mathcal{Y}_{\delta_1, \gamma'}$  being as in the statement of Theorem 4.1. For  $s > 1/2$ , we let  $\mathcal{Z}_{\delta,s} = \mathcal{Z}_{\delta, 1/2}$ . We follow the construction in the proof of Lemma 2.8, with the same notation introduced there. We obtain, by subadditivity and (4.6),

$$\begin{aligned} \left| (\bar{\mathbf{A}}^{-1/2}(\mathbf{A}(z + \Diamond_k) - \bar{\mathbf{A}})\bar{\mathbf{A}}^{-1/2})_+ \right| &\leq \sum_{j=-\infty}^k \frac{|V_j(y)|}{|\Diamond_k|} \left| (\bar{\mathbf{A}}^{-1/2}(\mathbf{A}(V_j(y)) - \bar{\mathbf{A}})\bar{\mathbf{A}}^{-1/2})_+ \right| \\ &\leq \delta_1 3^{\gamma'(m-k)} \left( \frac{\mathcal{Y}_{\delta_1} \vee \mathcal{S}}{3^m} \right)^{2\theta} \sum_{j=-\infty}^k \frac{|V_j(y)|}{|\Diamond_k|} 3^{\gamma'(k-j)} \\ &\leq \frac{C\Pi^{1/2}\delta_1}{1 - \gamma'} 3^{\gamma'(m-k)} \left( \frac{\mathcal{Y}_{\delta_1} \vee \mathcal{S}}{3^m} \right)^{2\theta}. \end{aligned}$$

Therefore,

$$\begin{aligned} \mathcal{E}_s(z + \Diamond_n) &\leq \left( \frac{C\Pi^{1/2}\delta_1}{1 - \gamma'} 3^{\gamma'(m-n)} \left( \frac{\mathcal{Y}_{\delta_1} \vee \mathcal{S}}{3^m} \right)^{2\theta} \right)^{1/2} \sum_{k=-\infty}^n 3^{-(n-k)(s-\gamma'/2)} \\ &\leq \left( \frac{C\Pi^{1/2}\delta_1}{\delta^2(1 - \gamma')(2s - \gamma')^2} \right)^{1/2} \delta 3^{\frac{\gamma'}{2}(m-n)} \left( \frac{\mathcal{Y}_{\delta_1} \vee \mathcal{S}}{3^m} \right)^\theta. \end{aligned}$$

Taking thus  $c$  small enough in the definition of  $\delta_1$ , the result follows for adapted cubes. With the same proof, by changing the definition  $V_j$ , we obtain the result also for the adapted simplexes.  $\square$

## 5. Equations with weaker mixing assumptions

We do not require a general mixing condition to hold, we just need a linear concentration for sums of  $\mathbf{A}(U_i)$  indexed over certain finite, disjoint  $\{U_i\}$  (as in (P3)). This turns out to be essential for the application considered in [ABRK23], in which the field  $\mathbf{a}(\cdot)$  has very weak decay of correlations in a literal sense, but faster decay of correlations in a coarse-grained sense.

(P2') *Ellipticity of coarse-grained coefficients.* There exists  $\gamma \in [0, 1)$ ,  $H \in [1, \infty)$ ,  $D \in [0, \infty)$ ,  $m_2 \in \mathbb{N}$ , a nondecreasing function  $\Psi_S : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  with constants  $K_{\Psi_S} \in [1, \infty)$  and  $p_{\Psi_S} \in (2, \infty)$  satisfying, for every  $p \in [2, p_{\Psi_S}]$ ,

$$s^p \leq K_{\Psi_S}^{3[p]^2} \frac{\Psi_S(ts)}{\Psi_S(t)}, \quad \forall t, s \in [1, \infty), \quad (5.1)$$

such that, for every  $m \in \mathbb{N}$  with  $m \geq m_2$ , we have

$$\sup_{k \in \mathbb{Z} \cap (-\infty, m]} 3^{\gamma(k-m)} \max_{z \in 3^k \mathbb{Z}^d \cap \square_m} |(\bar{\mathbf{A}}^{-1/2}(\square_m) \mathbf{A}(z + \square_k) \bar{\mathbf{A}}^{-1/2}(\square_m) - \mathbf{I}_{2d})_+| \leq \mathcal{O}_{\Psi_S}(Hm^D). \quad (5.2)$$

(P3') *Concentrations for sum (CFS)*. There exist  $\beta \in [0, 1)$ ,  $L_1, L_2 \in [1, \infty)$ ,  $m_3 \in \mathbb{N}$ , a nonincreasing, positive sequence  $\mathbb{N} \ni n \mapsto \omega_n$  with  $\lim_{n \rightarrow \infty} \omega_n = 0$ , an increasing function  $\Psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  and constants  $K_\Psi \in [1, \infty)$  and  $p_\Psi \in (d, \infty)$  satisfying the growth condition, for every  $p \in (1, p_\Psi]$ ,

$$s^p \leq K_\Psi^{3[p]^2} \frac{\Psi(ts)}{\Psi(t)}, \quad \forall t, s \in [1, \infty), \quad (5.3)$$

such that, for every  $m, n \in \mathbb{N}$  with  $n \geq m_3$  and  $\beta m < n < m - L_1 \log(L_2 n)$ ,

$$\left| \sum_{z \in 3^n \mathbb{Z}^d \cap \square_m} \bar{\mathbf{A}}^{-1/2}(\square_n) \mathbf{A}(z + \square_n) \bar{\mathbf{A}}^{-1/2}(\square_n) - \mathbf{I}_{2d} \right| \leq \mathcal{O}_\Psi(\omega_n). \quad (5.4)$$

In this section, we also make an isotropy assumption for simplicity.

(P4) *Dihedral symmetry*: the law  $\mathbb{P}$  is invariant under negation, reflections and permutations across the coordinate planes. That is, for every matrix  $R$  with exactly one  $\pm 1$  in each row and column and 0s elsewhere, the law of the conjugated coefficient  $R^t \mathbf{a}(R \cdot) R$  is the same as that of  $\mathbf{a}$ .

**Theorem 5.1.** *Assume that  $\mathbb{P}$  satisfies (P1), (P2'), (P3') and (P4). There exist constants  $C(d) < \infty$  and  $c(d), \alpha(d) \in (0, 1)$  such that, by defining*

$$\Upsilon_1 := \frac{CK_\Psi^{4d^2}}{\min\{d+1, p_\Psi\} - d}, \quad \Upsilon_2 := \frac{C}{\min\{3, p_{\Psi_S}\} - 2} \exp\left(C\left(L_1 \log L_2 + \frac{D + \log(H + K_{\Psi_S})}{1 - \gamma}\right)\right),$$

then, for every  $m, m_0 \in \mathbb{N}$  satisfying the conditions  $m \geq \max\{m_2, m_3\}$ ,

$$\omega_m^2 \leq \Upsilon_1^{-1} \quad \text{and} \quad m_0 \geq \frac{CD}{(1 - \beta)(1 - \gamma)^2} \log((\Upsilon_1 + \Upsilon_2)(m_2 \vee m_3 + m_0)\Theta_0) \log^2 6\Theta_0, \quad (5.5)$$

we have, for  $\kappa := \min\{\alpha, 1 - \gamma\}$  and for every  $n \in \mathbb{N}$  with  $n \geq m + m_0$ ,

$$\Theta_n - 1 \leq \Upsilon_1 \omega_m^2 + C 3^{-\kappa(n-m-m_0)}. \quad (5.6)$$

We have done most of the work for this theorem already in Sections 3 and 4. For this reason we will be somewhat brief with the details.

We first demonstrate how the assumptions (P1), (P2'), (P3') and (P4) yield information about the weak norms. The main technical tool is Lemma 2.14, together with already obtained estimates such as (3.80) and (3.81).

Due to isotropy, we observe that, for every  $n \in \mathbb{N}$ , the coarse-grained matrices  $\bar{\mathbf{s}}(\square_n)$ ,  $\bar{\mathbf{s}}_*(\square_n)$  and  $\bar{\mathbf{k}}(\square_n)$  are all scalar multiples of  $\mathbf{I}_d$ . This means that  $\bar{\mathbf{A}}(\square_n)$  has scalar block matrices. In particular, if  $\mathbf{E}_0 = \bar{\mathbf{A}}(\square_n)$ , then  $\mathbf{m}_0$  is a multiple of the identity and therefore so is  $\mathbf{q}_0$ . Thus the adapted cubes play no role in this case, and we will work using just normal triadic cubes.

**Lemma 5.2.** *There exists a constant  $C(d) < \infty$  such that for every  $p, q \in \mathbb{R}^d$  and for every  $n, h \in \mathbb{N}$  with  $h \geq \max\{(\beta \vee 1/2)n, m_2, m_3\}$ , and for  $\mathbf{E}_0 := \overline{\mathbf{A}}(\square_h)$  and  $\mathbf{m}_0, \mathbf{M}_0$  defined by (2.93), we have that*

$$\begin{aligned} & 3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \left( \frac{\nabla v_n - (\nabla v_n)_{\square_n}}{\mathbf{a} \nabla v_n - (\mathbf{a} \nabla v_n)_{\square_n}} \right) \right]_{\underline{B}_{2,1}^{-1/2}(\square_n)}^2 \right] \\ & \leq C \Theta_h^{1/2} \left| \overline{\mathbf{A}}^{1/2}(\square_h) \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2 \sum_{k=h}^n 3^{\frac{1}{2}(k-n)} \mathbb{E} \left[ \left| \overline{\mathbf{A}}^{-1/2}(\square_h) (\mathbf{A}(\square_k) - \mathbf{A}(\square_n)) \overline{\mathbf{A}}^{-1/2}(\square_h) \right|^2 \right] \\ & \quad + C \Theta_h^{1/2} \left| \overline{\mathbf{A}}^{1/2}(\square_h) \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2 \sum_{k=h}^n 3^{\frac{1}{2}(k-n)} \left| \overline{\mathbf{A}}^{-1/2}(\square_h) (\overline{\mathbf{A}}(\square_k) - \overline{\mathbf{A}}(\square_n)) \overline{\mathbf{A}}^{-1/2}(\square_h) \right| \\ & \quad + C \Theta_h^{1/2} \left| \overline{\mathbf{A}}^{1/2}(\square_h) \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2 \left( \frac{K_{\Psi}^{4d^2} \omega_h^2}{\min\{d+1, p_{\Psi}\} - d} + \frac{3^{-\frac{1}{2}(1-\gamma)(n-h-K \log h - h_c)}}{\min\{3, p_{\Psi_S}\} - 2} \right), \end{aligned} \quad (5.7)$$

where

$$K := L_1 + \frac{2D}{1-\gamma} \quad \text{and} \quad h_c := L_1 \log L_2 + \frac{100}{1-\gamma} (D + \log H + K_{\Psi_S}). \quad (5.8)$$

*Proof.* We will follow parts of the proof of Lemma 3.8. Fix  $n, h \in \mathbb{N}$  satisfying  $h \geq \max\{\beta n, m_2, m_3\}$  and  $p, q \in \mathbb{R}^d$ . Set

$$\eta := \min\{p_{\Psi}, d+1\} \quad \text{and} \quad \theta := \min\{p_{\Psi_S}, 3\}.$$

Set also  $\rho' := \gamma \vee d/\eta \vee 2/\theta$ , and let  $\rho := \frac{1+\rho'}{2}$ . Let  $\mathbf{E} = \mathbf{E}_0 := \overline{\mathbf{A}}(\square_h)$  and  $\mathbf{M} := \mathbf{M}_0$ . We have that  $|\mathbf{M}^{-1} \mathbf{E}| \leq C \Theta_h^{1/2}$ . Lemma 2.14 is applicable<sup>12</sup> with  $\diamond_k = \square_k$  for every  $k \in \mathbb{Z}$ , and we will estimate different terms appearing on the right in (2.133).

We first show that there exists a constant  $C(d) < \infty$  such that

$$\begin{aligned} & \mathbb{E} \left[ \left( \max_{k \in \mathbb{Z} \cap (-\infty, n]} 3^{-\rho(n-k)} \max_{z \in 3^k \mathbb{L}_0 \cap \diamond_n} \left| (\overline{\mathbf{A}}^{-1/2}(\square_h) (\mathbf{A}(z + \square_k) - \overline{\mathbf{A}}(\square_h)) \overline{\mathbf{A}}^{-1/2}(\square_h))_+ \right| \right)^2 \right] \\ & \leq C \left( \frac{K_{\Psi}^{4d^2} \omega_h^2}{\min\{d+1, p_{\Psi}\} - d} + \frac{K_{\Psi_S}^{36} H^2 n^{2D} 3^{-\frac{1}{2}(1-\gamma)(n-h-L_1 \log(L_2 h))}}{\min\{3, p_{\Psi_S}\} - 2} \right). \end{aligned} \quad (5.9)$$

On the one hand, by subadditivity and (P3') we have that, for every  $k \in \mathbb{N}$  with  $\beta k \leq h \leq k$  and  $k \geq h + L_1 \log(L_2 h)$ ,

$$\begin{aligned} & \left| (\overline{\mathbf{A}}^{-1/2}(\square_h) (\mathbf{A}(z + \square_k) - \overline{\mathbf{A}}(\square_h)) \overline{\mathbf{A}}^{-1/2}(\square_h))_+ \right| \\ & \leq \left| \sum_{z' \in z + 3^h \mathbb{L}_0 \cap \square_k} \overline{\mathbf{A}}^{-1/2}(\square_h) (\mathbf{A}(z + \square_h) - \overline{\mathbf{A}}(\square_h)) \overline{\mathbf{A}}^{-1/2}(\square_h) \right| \leq \mathcal{O}_{\Psi}(\omega_h). \end{aligned}$$

By a union bound, it follows that, for every  $t \geq 1$  and  $k \in \mathbb{N}$  with  $\beta k \leq h \leq k$  and  $k \geq h + L_1 \log(L_2 h)$ ,

$$\begin{aligned} & \mathbb{P} \left[ 3^{-\rho(n-k)} \max_{z \in 3^k \mathbb{L}_0 \cap \square_n} \left| (\overline{\mathbf{A}}^{-1/2}(\square_h) (\mathbf{A}(z + \square_k) - \overline{\mathbf{A}}(\square_h)) \overline{\mathbf{A}}^{-1/2}(\square_h))_+ \right| > \omega_h t \right] \\ & \leq \frac{3^{d(n-k)}}{\Psi(3^{\rho(n-k)} t)} \leq K^{4d^2} 3^{-(\rho\eta-d)(n-k)} t^{-\eta}. \end{aligned}$$

---

<sup>12</sup>The geometry of the adapted cubes does not play any particular role in the proof of Lemma 2.14.

Thus, by another union bound and (5.3), with  $h' := h + L_1 \log(L_2 h)$ ,

$$\mathbb{P} \left[ \max_{k \in \mathbb{N} \cap [h', n]} 3^{-\rho(n-k)} \max_{z \in 3^k \mathbb{L}_0 \cap \square_n} \left| (\overline{\mathbf{A}}^{-1/2}(\square_h) (\mathbf{A}(z + \square_k) - \overline{\mathbf{A}}(\square_h)) \overline{\mathbf{A}}^{-1/2}(\square_h))_+ \right| > \omega_h t \right] \leq \frac{2K_\Psi^{4d^2}}{\eta\rho - d} t^{-\eta}.$$

On the other hand, since  $\overline{\mathbf{A}}(\square_h) \geq \overline{\mathbf{A}}(\square_n)$ , we have by (P2') that

$$\begin{aligned} & \mathbb{P} \left[ \sup_{k \in \mathbb{Z} \cap (-\infty, h')} 3^{-\rho(n-k)} \max_{z \in 3^k \mathbb{Z}^d \cap \square_n} \left| (\overline{\mathbf{A}}^{-1/2}(\square_h) \mathbf{A}(z + \square_k) \overline{\mathbf{A}}^{-1/2}(\square_h) - \mathbf{I}_{2d})_+ \right| > H n^D 3^{-(\rho-\gamma)(n-h')} t \right] \\ & \leq \mathbb{P} \left[ \sup_{k \in \mathbb{Z} \cap (-\infty, n]} 3^{-\gamma(n-k)} \max_{z \in 3^k \mathbb{Z}^d \cap \square_n} \left| (\overline{\mathbf{A}}^{-1/2}(\square_n) \mathbf{A}(z + \square_k) \overline{\mathbf{A}}^{-1/2}(\square_n) - \mathbf{I}_{2d})_+ \right| > H n^D t \right] \\ & \leq \frac{1}{\Psi_S(t)} \leq C K_{\Psi_S}^{36} t^{-\theta}. \end{aligned}$$

Combining the previous two displays yields (5.9) in view of the definition of  $\rho$ .

We may now conclude the proof by applying Lemma 2.14 with  $\mathbf{E} = \overline{\mathbf{A}}(\square_h)$  and  $\mathbf{M} := \mathbf{M}_0$ , proceeding as in (3.80), but using this time (5.9).  $\square$

*Proof of Theorem 5.1. Step 1.* Define, for  $m \in \mathbb{Z}$ ,

$$\Theta_m := \min_{\mathbf{h} \in \mathbb{R}_{\text{skew}}^{d \times d}} \left| (\overline{\mathbf{s}}_*^{-1} (\overline{\mathbf{s}} + (\overline{\mathbf{k}} - \mathbf{h})^t \overline{\mathbf{s}}_*^{-1} (\overline{\mathbf{k}} - \mathbf{h}))) (\square_m) \right|, \quad (5.10)$$

and recall that  $\mathbb{N} \ni m \mapsto \Theta_m$  is non-increasing. We let  $\delta, \sigma \in (0, (80d)^{-2}]$  to be parameters to be fixed by means of  $d$ , and let  $K$  and  $h_c$  be as in (5.8). We take  $m_1 \in \mathbb{N}$  so large that

$$m_1 \geq \max\{m_2, m_3\} \quad \text{and} \quad \frac{\omega_{m_1}^2 K_\Psi^{4d^2}}{\min\{d+1, p_\Psi\} - d} \leq \delta^2 \sigma \quad (5.11)$$

are both valid. We define the universal constants  $\Upsilon_1, \Upsilon_2, \Upsilon$  by

$$\Upsilon_1 := \frac{K_\Psi^{4d^2}}{\min\{d+1, p_\Psi\} - d}, \quad \Upsilon_2 := \frac{3^{h_c}}{\min\{3, p_{\Psi_S}\} - 2} \quad \text{and} \quad \Upsilon := \Upsilon_1 + \Upsilon_2.$$

Assume that  $m_0 = m_0(\delta, \sigma)$  is the smallest integer satisfying the condition

$$m_0(\delta, \sigma) \geq \frac{1}{\delta^3 \sigma^2 (1-\beta)(1-\gamma)} \log \left( \frac{6\Upsilon(m_0(\delta, \sigma) + m_1)^D \Theta_0}{\delta} \right) \log^2 6\Theta_0. \quad (5.12)$$

*Step 1.* We claim that there exists  $\delta_0(d) \in (0, 1)$  such that if  $\delta \leq \delta_0$ , then

$$\Theta_{m_1+2m_0} \leq 1 + \sigma. \quad (5.13)$$

To show this, we follow the outline of Section 3. Define

$$L = L(\sigma, \Theta_m) := \frac{K}{\delta(1-\gamma)} \log \left( \frac{(m_1 + 2m_0)^D \Upsilon \Theta_m}{\delta \sigma^2} \right) + 8h_c \quad (5.14)$$

and

$$M = M(\sigma, \Theta_m) := \left\lceil \frac{6d \log(6\Theta_m) L(\sigma, \Theta_m)}{\delta \sigma^2} \right\rceil. \quad (5.15)$$

Notice that, for every  $h \in \mathbb{N}$  with  $m_1 \leq h \leq m_1 + 2m_0$ ,

$$L_1 \log(L_2 h) + K \log h + h_c \leq \frac{1}{4} L. \quad (5.16)$$

By (5.11) and (5.14), we have

$$\Upsilon_1 \omega_{m_1}^2 + \Upsilon 3^{-\frac{1}{8}(1-\gamma)L} \leq 2\delta \sigma^2. \quad (5.17)$$

As in Section 3, we first show that  $\Theta_m \leq 2$  for some  $m \in \mathbb{N}$  with  $m_1 + m_0 \leq m \leq m_1 + \frac{3}{2}m_0$  using  $\sigma = 1$ , and then conclude with  $\sigma \in (0, 1]$ . Notice that if  $m \in \mathbb{N}$  is such that  $m \geq m_1 + m_0$ , then  $2M(1, \Theta_0) \leq (1 - \beta)m$ , and if  $\Theta_m \leq 2$ , then  $2M(\sigma, 2) \leq (1 - \beta)m$ . In particular,  $\beta(m + 2M) \leq m$ .

Take now  $M = M(1, \Theta_0)$ . As in Lemma 3.4, we obtain by a pigeonhole argument that, for every  $m \in \mathbb{N}$  with  $m_1 + m_0 \leq m \leq m_1 + 2m_0 - 2M$ , there exists  $h \in [m + M, m + 2M - 2L]$  such that, for every  $k \in \mathbb{N}$  with  $h \leq k \leq h + 2L$ , we have that  $h \geq \beta k$  and

$$|\bar{\mathbf{A}}^{-1/2}(\square_k) \bar{\mathbf{A}}(\square_h) \bar{\mathbf{A}}^{-1/2}(\square_k) - \mathbf{I}_{2d}| \leq \delta \sigma^2. \quad (5.18)$$

Now, since  $h \geq \beta k$  for every  $k \in \mathbb{N}$  with  $m \leq k \leq h + 2L$ , (5.4), (5.16) and (5.11) yield that, for every  $n \in \mathbb{N}$  with  $h + \frac{1}{4}L \leq n \leq h + 2L$ ,

$$\mathbb{E} \left[ \left| \sum_{z \in 3^h \mathbb{Z}^d \cap \square_n} \bar{\mathbf{A}}^{-1/2}(\square_h) (\mathbf{A}(z + \square_h) - \bar{\mathbf{A}}(\square_h)) \bar{\mathbf{A}}^{-1/2}(\square_h) \right|^2 \right] \leq K_{\Psi}^{27} \omega_h^2 \leq \delta \sigma^2. \quad (5.19)$$

Using (5.18) and (5.19) in (3.28) (with  $m = k = h$ ), we deduce that, for every  $n \in \mathbb{N}$  with  $h + \frac{1}{4}L \leq n \leq h + 2L$ ,

$$\mathbb{E} \left[ |\bar{\mathbf{A}}^{-1/2}(\square_h) \mathbf{A}(\square_n) \bar{\mathbf{A}}^{-1/2}(\square_h) - \mathbf{I}_{2d}|^2 \right] \leq C \delta \sigma^2. \quad (5.20)$$

Set  $\mathbf{E}_0 := \bar{\mathbf{A}}(\square_h)$  and  $\bar{\mathbf{t}}_n := (\bar{\mathbf{b}} \# \bar{\mathbf{s}}_*)(\square_n)$ , and take, for  $e \in \mathbb{R}^d$  with  $|e| = 1$ ,

$$\begin{pmatrix} p \\ q \end{pmatrix} := \begin{pmatrix} \bar{\mathbf{t}}_n & 0 \\ 0 & \bar{\mathbf{t}}_n \end{pmatrix}^{-1/2} \begin{pmatrix} e \\ \bar{\mathbf{t}}_n e \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} P \\ Q \end{pmatrix} := \mathbb{E} \left[ \int_{\square_n} \begin{pmatrix} \nabla v(\cdot, \square_n, p, q) \\ \mathbf{a} \nabla v(\cdot, \square_n, p, q) \end{pmatrix} \right]. \quad (5.21)$$

We insert these choices into (3.65) and obtain that, for  $n := h + 2L$  and  $k := h + L$ , and for every  $\varepsilon \in (0, 1]$ ,

$$\begin{aligned} \left| \mathbb{E}[J(\square_n)] - \frac{1}{2} P \cdot Q \right| &\leq 50(1 + \varepsilon^{-1}) \bar{\tau}_{n,k} + C(\varepsilon + 3^{-L}) (|\mathbf{b}_0^{1/2} P| + |\mathbf{s}_{*,0}^{-1/2} Q|)^2 \\ &\quad + C 3^{-L} (|\mathbf{b}_0^{1/2} P| + |\mathbf{s}_{*,0}^{-1/2} Q|)^2 \sum_{j=-\infty}^k 3^{\frac{3}{2}(j-k)} \sum_{z \in 3^j \mathbb{Z}^d \cap \square_k} |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}(z + \square_j) \mathbf{E}_0^{-1/2}| \\ &\quad + C 3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \begin{pmatrix} \nabla v_n - P \\ \mathbf{a} \nabla v_n - Q \end{pmatrix} \right]^2 \right]_{\dot{\bar{B}}_{2,1}^{-1/2}(\square_n)}. \end{aligned} \quad (5.22)$$

We take above  $\varepsilon := \delta^{1/2} \sigma \Theta_n^{-1/2}$ . Completely analogously to (3.69) and (3.71), we have that

$$\left| \bar{\mathbf{A}}^{1/2}(\square_n) \begin{pmatrix} -p \\ q \end{pmatrix} \right|^2 \leq C \Theta_n^{1/2} \quad \text{and} \quad |\mathbf{b}_0^{1/2} P| + |\mathbf{s}_{*,0}^{-1/2} Q| \leq C \Theta_h^{1/2} \Theta_n^{1/4} \quad (5.23)$$

and, consequently, as in (3.66), by (5.18) and the previous display, we get that

$$\bar{\tau}_{n,k} \leq C\sigma^2\Theta_n^{1/2}.$$

Furthermore, by (5.18) and (5.9), we have

$$\sum_{j=h}^k 3^{\frac{3}{2}(j-k)} |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}(\square_j) \mathbf{E}_0^{-1/2}| \leq 3 \quad \text{and} \quad \sum_{j=-\infty}^h 3^{\frac{3}{2}(j-k)} \sum_{z \in 3^j \mathbb{Z}^d \cap \square_k} |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}(z + \square_j) \mathbf{E}_0^{-1/2}| \leq C,$$

respectively. Thus,

$$\sum_{j=-\infty}^k 3^{\frac{3}{2}(j-k)} \sum_{z \in 3^j \mathbb{Z}^d \cap \square_k} |\mathbf{E}_0^{-1/2} \bar{\mathbf{A}}(z + \square_j) \mathbf{E}_0^{-1/2}| \leq C. \quad (5.24)$$

Combining the above displays then yields that

$$\begin{aligned} \left| \mathbb{E}[J(\square_n)] - \frac{1}{2} P \cdot Q \right| &\leq C\sigma\delta^{1/2}\Theta_n + C(\sigma\delta^{1/2} + 3^{-L}\Theta_n^{1/2})\Theta_h \\ &\quad + C3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \left( \frac{\nabla v_n - P}{\mathbf{a} \nabla v_n - Q} \right) \right]_{\dot{\bar{B}}_{2,1}^{-1/2}(\square_n)}^2 \right]. \end{aligned}$$

To estimate the last term on the right, we use Lemma 5.2, (5.23), (5.18), (5.20) and (5.17), and  $n \geq h + \frac{1}{4}L$ , to deduce that

$$3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \left( \frac{\nabla v_n - (\nabla v_n)_{\square_n}}{\mathbf{a} \nabla v_n - (\mathbf{a} \nabla v_n)_{\square_n}} \right) \right]_{\dot{\bar{B}}_{2,1}^{-1/2}(\square_n)}^2 \right] \leq C\sigma^2\Theta_h^{1/2}\Theta_n^{1/2}.$$

By similar estimates for  $J^*$ , Lemma 3.6 and (5.14) yield that

$$\Theta_n - 1 \leq C\sigma\delta^{1/2}\Theta_n + C(\sigma\delta^{1/2} + 3^{-L}\Theta_n^{1/2})\Theta_h \leq \frac{\sigma}{8}(\Theta_n - 1) + \frac{\sigma}{8}(\Theta_h - 1) + \frac{\sigma}{4}.$$

After reabsorption and using the monotonicity of  $m \mapsto \Theta_m$ , we then deduce that

$$\Theta_{m+2M} - 1 \leq \frac{\sigma}{4}(\Theta_m - 1) + \frac{\sigma}{2}.$$

Thus, by iterating this  $K = 1 + \lceil \log \Theta_m \rceil$  many times, we deduce that

$$\Theta_{m_1+m_0+2KM} - 1 \leq 4^{-K}(\Theta_{m_0} - 1) + 2/3 \leq 2.$$

Notice that this iteration is done using  $\sigma = 1$ . Iterating once more then yields that

$$\Theta_{m_1+m_0+2KM(1,\Theta_0)+M(\sigma,2)} \leq 1 + \sigma.$$

By the definition of  $m_0$  we see that  $2KM(1,\Theta_0) + M(\sigma,2) \leq m_0$ , and hence we deduce (5.13) by the monotonicity of  $m \mapsto \Theta_m$ .

*Step 2.* We next claim that there exists constants  $\alpha(d), \sigma(d) \in (0, (80d^2)^{-1})$  such that, for  $\kappa := \min\{\alpha, 1 - \gamma\}$  and for every  $n \in \mathbb{N}$  with  $n \geq m_1 + 4m_0(\delta_0, \sigma)$ ,

$$\Theta_n - 1 \leq \min \left\{ \sigma, \sigma^{-1} (\Upsilon_1 \omega_{m_1}^2 + 3^{-\kappa(n-m_1-4m_0)}) \right\}. \quad (5.25)$$

We assume inductively that there exists  $m \in \mathbb{N}$  with  $m > m_1 + 4m_0$  such that (5.25) is valid for every  $n \in \mathbb{N}$  with  $n \in [m_1 + 4m_0, m - 1]$ . The initial step  $n = m_1 + 4m_0$  is valid by Step 1. Our goal is to show that (5.25) is true also for  $n = m$ , which then proves the induction step.

To prove (5.25) with  $n = m$ , we first go back to the proof of Lemma 4.3. Setting  $\hat{\Theta}_m := (\bar{\mathbf{s}}_*^{-1/2} \bar{\mathbf{s}}_*^{-1/2})(\square_m)$  and repeating Step 1 in the proof of Lemma 4.3, we find that there exists  $C(d) < \infty$  such that, for every  $k, n \in \mathbb{N}$  with  $m_1 + 2m_0 \leq k \leq n$ ,

$$|\bar{\mathbf{A}}^{-1/2}(\square_n) \bar{\mathbf{A}}(\square_k) \bar{\mathbf{A}}^{-1/2}(\square_n) - \mathbf{I}_{2d}| \leq 4d(\hat{\Theta}_k - \hat{\Theta}_n) \leq \frac{1}{40}. \quad (5.26)$$

We then revisit Step 2 in the proof of Lemma 4.3. As a consequence of (4.12) we deduce that, for every  $k, m, n \in \mathbb{N}$  with  $2m_0 \leq k \leq n \leq m$  we have that

$$\begin{aligned} & \mathbb{E} \left[ \left| \bar{\mathbf{A}}^{-1/2}(\square_m) \mathbf{A}(\square_n) \bar{\mathbf{A}}^{-1/2}(\square_m) - \mathbf{I}_{2d} \right|^2 \right] \\ & \leq C(\hat{\Theta}_k - 1)^2 + C \mathbb{E} \left[ \left| \sum_{z \in 3^k \mathbb{Z}^d \cap \square_n} \bar{\mathbf{A}}^{-1/2}(\square_m) (\mathbf{A}(z + \square_k) - \bar{\mathbf{A}}_k) \bar{\mathbf{A}}^{-1/2}(\square_m) \right|^2 \right]. \end{aligned}$$

If  $\max\{m_1 + 2m_0, \lceil \beta m \rceil\} < k \leq m - L_1 \log(L_2 k)$ , then, by (P3') and (5.26),

$$\mathbb{E} \left[ \left| \sum_{z \in 3^k \mathbb{Z}^d \cap \square_n} \bar{\mathbf{A}}^{-1/2}(\square_m) (\mathbf{A}(z + \square_k) - \bar{\mathbf{A}}_k) \bar{\mathbf{A}}^{-1/2}(\square_m) \right|^2 \right] \leq CK_\Psi^{27} \omega_k^2 \leq C\Upsilon_1 \omega_{m_1}^2.$$

We thus obtain, for every  $n \in \mathbb{N}$  with  $m_1 + 3m_0 \leq n \leq m$ , that

$$\begin{aligned} \mathbb{E} \left[ \left| \bar{\mathbf{A}}^{-1/2}(\square_m) \mathbf{A}(\square_n) \bar{\mathbf{A}}^{-1/2}(\square_m) - \mathbf{I}_{2d} \right|^2 \right] & \leq C(\hat{\Theta}_{n - \lfloor L_1 \log(L_2 n) \rfloor} - 1)^2 + CK_\Psi^{27} \omega_{n - \lfloor L_1 \log(L_2 n) \rfloor}^2 \\ & \leq C \left( \Upsilon_1 \omega_{m_1}^2 + 3^{-\kappa(n - m_1 - 4m_0)} \right). \end{aligned} \quad (5.27)$$

We then follow the proof of Proposition 4.2. Let  $\mathbf{E}_0 := \mathbf{A}(\square_m)$  and set

$$\begin{pmatrix} p \\ q \end{pmatrix} = \begin{pmatrix} \bar{\mathbf{s}}_*^{-1/2}(\square_m) e \\ \bar{\mathbf{s}}_*^{1/2}(\square_m) e \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} P \\ Q \end{pmatrix} := \mathbb{E} \left[ \int_{\square_m} \begin{pmatrix} \nabla v(\cdot, \square_m, p, q) \\ \mathbf{a} \nabla v(\cdot, \square_m, p, q) \end{pmatrix} \right].$$

By repeating the argument for (4.22) and (4.24), we deduce that there exists a constant  $C(d) < \infty$

$$\hat{\Theta}_n - 1 \leq C 3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \begin{pmatrix} \nabla v_n - P \\ \mathbf{a} \nabla v_n - Q \end{pmatrix} \right]_{\dot{\mathbf{B}}_{2,1}^{-1/2}(\square_n)}^2 \right] + C(\hat{\Theta}_{n-4} - \hat{\Theta}_n) \quad (5.28)$$

for small enough  $\sigma(d)$ . Lemma 5.2, together with (5.26) and (5.27), yields that

$$\begin{aligned} & 3^{-n} \mathbb{E} \left[ \left[ \mathbf{M}_0^{1/2} \begin{pmatrix} \nabla v_n - (\nabla v_n)_{\square_n} \\ \mathbf{a} \nabla v_n - (\mathbf{a} \nabla v_n)_{\square_n} \end{pmatrix} \right]_{\dot{\mathbf{B}}_{2,1}^{-1/2}(\square_n)}^2 \right] \\ & \leq C \sum_{k=m_1+3m_0}^n 3^{\frac{1}{2}(k-n)} \mathbb{E} \left[ \left| \bar{\mathbf{A}}^{-1/2}(\square_m) (\mathbf{A}(\square_k) - \bar{\mathbf{A}}(\square_k)) \bar{\mathbf{A}}^{-1/2}(\square_m) \right|^2 \right] \\ & \quad + C \sum_{k=m_1+3m_0}^n 3^{\frac{1}{2}(k-n)} \left| \bar{\mathbf{A}}^{-1/2}(\square_m) (\bar{\mathbf{A}}(\square_k) - \mathbf{A}(\square_n)) \bar{\mathbf{A}}^{-1/2}(\square_m) \right| \\ & \quad + C \left( \Upsilon_1 \omega_{m_1}^2 + 3^{-\kappa(n - m_1 - 4m_0)} \right) \\ & \leq C \sum_{k=m_1+3m_0}^n 3^{\frac{1}{2}(k-n)} (\hat{\Theta}_k - \hat{\Theta}_n) + C \left( \Upsilon_1 \omega_{m_1}^2 + 3^{-\kappa(n - m_1 - 4m_0)} \right). \end{aligned}$$

Combining this with (5.27) leads to

$$\hat{\Theta}_n - 1 \leq C \sum_{k=m_1+4m_0}^n 3^{\frac{1}{2}(k-n)} (\hat{\Theta}_k - \hat{\Theta}_m) + C \left( \Upsilon_1 \omega_{m_1}^2 + 3^{-\kappa(n-m_1-4m_0)} \right).$$

We have that

$$\sum_{k=m_1+4m_0}^n 3^{-2\kappa_0(n-k)} 3^{-(1-\gamma)(k-m_1-4m_0)} \leq \frac{2}{\kappa_0} 3^{-\min\{1-\gamma, \kappa_0\}(n-m_1-4m_0)}.$$

As in the proof of Proposition 4.2, we obtain by iteration that there exists constants  $\alpha(d) \in (0, 1)$  and  $C(d) < \infty$  such that, with  $\kappa := \min\{\alpha, \frac{1}{2}(1-\gamma)\}$ ,

$$\Theta_m - 1 \leq \frac{C}{\kappa_0 \alpha} \left( \Upsilon_1 \omega_{m_1}^2 + 3^{-\kappa(n-m_1-4m_0)} \right).$$

Therefore, by taking  $\sigma := C^{-1} \kappa_0 \alpha$ , which then depends solely on  $d$ , we obtain (5.25) for  $n = m$ , proving the induction step.

*Step 4. Conclusion.* The result follows by (5.25) after relabelling  $\Upsilon$  and taking  $m_0$  larger as in (5.5), and using the monotonicity of  $m \mapsto \Theta_m$ . The proof is complete.  $\square$

## 6. Quantitative homogenization

Up to this point, this paper has been focused on elaborating properties of the coarse-grained coefficients  $\mathbf{A}(U)$  and proving, under the assumptions (P1), (P2) and (P3) (and variants of these), quantitative estimates on their convergence to the homogenized matrix  $\bar{\mathbf{A}}$  as the domain  $U$  becomes large. This section has a completely different focus: we show that the coarse-grained coefficients can be used to control general solutions. For instance, we would like to show that an assumption that  $\mathbf{A}(V)$  is quantitatively close to  $\bar{\mathbf{A}}$ , for an appropriate collection of subdomains  $V$  belonging to a larger domain  $U$ , gives us a quantitative estimate on the difference between the solutions of the Dirichlet problems for  $-\nabla \cdot \mathbf{a} \nabla$  and  $-\nabla \cdot \bar{\mathbf{a}} \nabla$ , respectively, in  $U$ . Such a statement can be found below in Proposition 6.7.

The arguments here (and thus most of the statements) are *purely deterministic*—in particular, they are independent of the main results from the previous sections and, indeed, independent of the assumptions (P1), (P2) and (P3). Of course, in the context in which these assumptions are valid, then these deterministic estimates can be combined with results proved in the previous sections, such as Theorem 4.1, to immediately yield quantitative homogenization results, like those stated in Theorem B.

Throughout this section we fix symmetric matrix  $\mathbf{s}_0$  and antisymmetric matrix  $\mathbf{k}_0$ ,  $\mathbf{a}_0 = \mathbf{s}_0 + \mathbf{k}_0$ , and

$$\mathbf{A}_0 := \begin{pmatrix} \mathbf{s}_0 + \mathbf{k}_0^t \mathbf{s}_0^{-1} \mathbf{k}_0 & -\mathbf{k}_0^t \mathbf{s}_0 \\ -\mathbf{s}_0 \mathbf{k}_0 & \mathbf{s}_0^{-1} \end{pmatrix}.$$

Let also  $\lambda_0 := |\mathbf{s}_0^{-1}|^{-1}$ ,  $\Lambda_0 := |\mathbf{s}_0|$  and  $\Pi_0 := \Lambda_0 / \lambda_0 = |\mathbf{s}_0| |\mathbf{s}_0^{-1}|$ . The geometry of adapted cubes  $\diamond_n$  is dictated by  $\mathbf{m}_0 := \mathbf{s}_0$  and  $\mathbf{q}_0$  as in Subsection 2.7 so that  $\frac{1}{2} \mathbf{s}_0^{1/2} \leq \lambda_0^{1/2} \mathbf{q}_0 \leq 2 \mathbf{s}_0^{1/2}$ .



Now, to find the right quantity to measure the homogenization, we first recall (2.27), which gives us, for every Lipschitz domain  $U$ ,

$$\begin{aligned} & \sup_{|e| \leq 1} \left( J(U, \mathbf{s}_0^{-1/2} e, \mathbf{a}_0^t \mathbf{s}_0^{-1/2} e) + J^*(U, \mathbf{s}_0^{-1/2} e, \mathbf{a}_0 \mathbf{s}_0^{-1/2} e) \right) \\ & \leq \left( |\mathbf{s}_0^{-1}(\mathbf{s} - \mathbf{s}_*)(\cdot)| + |\mathbf{s}_0^{-1/2} \mathbf{s}_*^{1/2}(\cdot) - \mathbf{s}_0^{1/2} \mathbf{s}_*^{-1/2}(\cdot)|^2 + |\mathbf{s}_0^{-1/2}(\mathbf{k}(\cdot) - \mathbf{k}_0) \mathbf{s}_*^{-1/2}(\cdot)|^2 \right)(U) \\ & \leq 3 \sup_{|e| \leq 1} \left( J(U, \mathbf{s}_0^{-1/2} e, \mathbf{a}_0^t \mathbf{s}_0^{-1/2} e) + J^*(U, \mathbf{s}_0^{-1/2} e, \mathbf{a}_0 \mathbf{s}_0^{-1/2} e) \right). \end{aligned} \quad (6.1)$$

In view of this, we define

$$\begin{aligned} & \mathcal{E}_s(y + \Diamond_n; \mathbf{a}, \mathbf{a}_0) \\ & := s \sum_{k=-\infty}^n 3^{s(k-n)} \max_{z \in y + 3^k \mathbb{L}_0 \cap \Diamond_n} \sup_{|e| \leq 1} \left( J(\cdot, \mathbf{s}_0^{-1/2} e, \mathbf{a}_0^t \mathbf{s}_0^{-1/2} e) + J^*(\cdot, \mathbf{s}_0^{-1/2} e, \mathbf{a}_0 \mathbf{s}_0^{-1/2} e) \right)^{1/2} (z + \Diamond_k). \end{aligned} \quad (6.2)$$

By the subadditivity of both  $J$  and  $J^*$ , we get, for every  $k, n \in \mathbb{N}$  with  $k \leq n$ ,

$$\mathcal{E}_s(y + \Diamond_n; \mathbf{s}_0, \mathbf{k}_0) \leq \sum_{z \in y + 3^k \mathbb{L}_0 \cap \Diamond_n} \mathcal{E}_s(z + \Diamond_k; \mathbf{s}_0, \mathbf{k}_0). \quad (6.3)$$

We also define a version of  $\mathcal{E}_s$  using simplexes instead of cubes. For this, let  $\Delta \in \mathcal{Z}_n^\Delta$  and set

$$\begin{aligned} & \mathcal{E}_s^\Delta(\Delta; \mathbf{a}, \mathbf{a}_0) \\ & := s \sum_{k=-\infty}^n 3^{s(k-n)} \max_{\Delta' \in \mathcal{Z}_k^\Delta(\Delta)} \sup_{|e| \leq 1} \left( J(\Delta', \mathbf{s}_0^{-1/2} e, \mathbf{a}_0^t \mathbf{s}_0^{-1/2} e) + J^*(\Delta', \mathbf{s}_0^{-1/2} e, \mathbf{a}_0 \mathbf{s}_0^{-1/2} e) \right)^{1/2}. \end{aligned} \quad (6.4)$$

Recall that  $\mathcal{Z}_k^\Delta(U)$  is a collection of adapted simplexes of size  $3^k$  inside of  $U$ . Again, by subadditivity,

$$\mathcal{E}_s(y + \Diamond_n; \mathbf{a}, \mathbf{a}_0) \leq \sum_{\Delta \in \mathcal{Z}_k^\Delta(y + \Diamond_n)} \mathcal{E}_s^\Delta(\Delta; \mathbf{a}, \mathbf{a}_0). \quad (6.5)$$

Notice next that it is immediate that

$$\begin{aligned} & \mathcal{E}_s(y + \Diamond_n; \mathbf{a} - \mathbf{k}_0, \mathbf{a}_0 - \mathbf{k}_0) = \mathcal{E}_s(y + \Diamond_n; \mathbf{a} - \mathbf{k}_0, \mathbf{s}_0) \\ & = s \sum_{k=-\infty}^n 3^{s(k-n)} \max_{z \in y + 3^k \mathbb{L}_0 \cap \Diamond_n} \sup_{|e| \leq 1} \left( J_{\mathbf{k}_0}(z + \Diamond_k, \mathbf{s}_0^{-1/2} e, \mathbf{s}_0^{1/2} e) + J_{\mathbf{k}_0}^*(z + \Diamond_k, \mathbf{s}_0^{-1/2} e, \mathbf{s}_0^{1/2} e) \right)^{1/2}. \end{aligned} \quad (6.6)$$

Next we recall that a solution of the equation  $-\nabla \cdot \mathbf{a} \nabla u = 0$  is also solves the equation  $-\nabla \cdot (\mathbf{a} - \mathbf{k}_0) \nabla u = 0$ , since  $\mathbf{k}_0$  is antisymmetric. Consequently, we often assume in the proofs that, without loss of generality,  $\mathbf{k}_0 = 0$ , and use the right side of (6.6) as a measurement of the homogenization with  $\mathbf{k}_0 = 0$ . We also suppress  $\mathbf{a}, \mathbf{a}_0$  from the notation for  $\mathcal{E}_s$  in the proofs, when they are clear from the context.

In the setting of stochastic homogenization in which assumptions (P1), (P2) and (P3) are valid, and  $\mathbf{A}_0 = \overline{\mathbf{A}}$ , the quantity on the right in (6.2) is a random variable which can be estimated using Corollary 4.5. Indeed, for  $|e| \leq 1$ ,

$$\begin{aligned} & J_{\overline{\mathbf{k}}}(U, \overline{\mathbf{s}}^{-1/2} e, \overline{\mathbf{s}}^{1/2} e) + J_{\overline{\mathbf{k}}}^*(U, \overline{\mathbf{s}}^{-1/2} e, \overline{\mathbf{s}}^{1/2} e) \\ & = \frac{1}{2} \begin{pmatrix} -e \\ e \end{pmatrix} \cdot (\overline{\mathbf{A}}_{\overline{\mathbf{k}}}^{-1/2} \mathbf{A}_{\overline{\mathbf{k}}}(U) \overline{\mathbf{A}}_{\overline{\mathbf{k}}}^{-1/2} - \mathbf{I}_{2d}) \begin{pmatrix} -e \\ e \end{pmatrix} + \frac{1}{2} \begin{pmatrix} e \\ e \end{pmatrix} \cdot (\overline{\mathbf{A}}_{\overline{\mathbf{k}}}^{-1/2} \mathbf{A}_{\overline{\mathbf{k}}}(U) \overline{\mathbf{A}}_{\overline{\mathbf{k}}}^{-1/2} - \mathbf{I}_{2d}) \begin{pmatrix} e \\ e \end{pmatrix} \\ & \leq 2 \left| (\overline{\mathbf{A}}_{\overline{\mathbf{k}}}^{-1/2} \mathbf{A}_{\overline{\mathbf{k}}}(U) \overline{\mathbf{A}}_{\overline{\mathbf{k}}}^{-1/2} - \mathbf{I}_{2d})_+ \right|. \end{aligned}$$

It follows that

$$\mathcal{E}_s(y + \diamond_n; \mathbf{a}, \mathbf{a}_0) \leq 2\tilde{\mathcal{E}}_s(y + \diamond_n; \mathbf{a}, \mathbf{a}_0) \quad \text{and} \quad \mathcal{E}_s^\Delta(\Delta; \mathbf{a}, \mathbf{a}_0) \leq 2\tilde{\mathcal{E}}_s(\Delta; \mathbf{a}, \mathbf{a}_0) \quad (6.7)$$

with  $\tilde{\mathcal{E}}_s$  and  $\tilde{\mathcal{E}}_s^\Delta$  defined by (4.45) and (4.46), respectively and these have been estimated in Corollary 4.5. The only “random” ingredient needed in this section is the estimate (4.48).

**6.1. Coarse-grained Caccioppoli estimate.** The most fundamental estimate, which leads to all regularity for divergence-form elliptic equations, is the Caccioppoli inequality. For a solution  $u \in H^1(B_r)$  of the uniformly elliptic equation

$$-\nabla \cdot \mathbf{a} \nabla u = 0 \quad \text{in } U,$$

where  $\mathbf{a}(x)$  satisfies the uniform ellipticity condition (1.2), then this estimate states that

$$\|\phi \nabla u\|_{\underline{L}^2(U)}^2 \leq 4\Pi \|u \nabla \phi\|_{\underline{L}^2(U)}^2 \quad (6.8)$$

where  $\Pi := \Lambda/\lambda$  denotes the ellipticity ratio and  $\phi$  is any smooth cutoff function.

The purpose of this subsection is to *coarse-grain* the Caccioppoli inequality by getting rid of the factor of  $\Pi$  in (6.8), in domains  $U = \diamond_n$  for which the quantity  $\mathcal{E}_s(\diamond_n)$  is sufficiently small.

We begin by coarse-graining the gradient-to-flux map. This estimate states that, for an arbitrary solution  $u$ , the difference of  $\mathbf{a} \nabla u$  and  $\bar{\mathbf{a}} \nabla u$ , in a weak Besov norm is controlled by  $\mathcal{E}_s$ .

**Lemma 6.1.** *For every  $s \in (0, 1]$ ,  $y \in \mathbb{R}^d$  and  $n \in \mathbb{N}$ , we have the estimates*

$$s3^{-ns} [\mathbf{s}_0^{-1/2} (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u)]_{\dot{B}_{2,1}^{-s}(y + \diamond_n)} \leq 3^{d+2} \mathcal{E}_s(y + \diamond_n; \mathbf{a}, \mathbf{a}_0) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(y + \diamond_n)} \quad (6.9)$$

and

$$s3^{-ns} [\mathbf{s}_0^{1/2} \nabla u]_{\dot{B}_{2,1}^{-s}(y + \diamond_n)} \leq 3^{d+2} (1 + \mathcal{E}_s(y + \diamond_n; \mathbf{a}, \mathbf{a}_0)) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(y + \diamond_n)}. \quad (6.10)$$

*Proof.* First, by (2.37) and (2.38), we have

$$\left| \mathbf{s}_0^{-1/2} \int_{z + \diamond_k} (\mathbf{a} - \mathbf{a}_*(z + \diamond_k)) \nabla u \right|^2 \leq 2 |\mathbf{s}_0^{-1} (\mathbf{s} - \mathbf{s}_*)(z + \diamond_k)| \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(z + \diamond_k)}^2 \quad (6.11)$$

and

$$\left| \mathbf{s}_*^{1/2}(z + \diamond_k) \int_{z + \diamond_k} \nabla u \right|^2 \leq \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(z + \diamond_k)}^2.$$

Using these and the triangle inequality, we also have

$$\left| \mathbf{s}_0^{-1/2} \int_{z + \diamond_k} (\mathbf{a}_*(z + \diamond_k) - \mathbf{a}_0) \nabla u \right|^2 \leq |\mathbf{s}_0^{-1/2} (\mathbf{a}_*(z + \diamond_k) - \mathbf{a}_0) \mathbf{s}_*^{-1/2}(z + \diamond_k)|^2 \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(z + \diamond_k)}^2.$$

It thus follows, by defining

$$\mathcal{U}_{z,k} := |\mathbf{s}_0^{-1} (\mathbf{s} - \mathbf{s}_*)(z + \diamond_k)| + |\mathbf{s}_0^{-1/2} (\mathbf{a}_*(z + \diamond_k) - \mathbf{a}_0) \mathbf{s}_*^{-1/2}(z + \diamond_k)|^2 \quad (6.12)$$

and using (2.126), that

$$\begin{aligned}
[\mathbf{s}_0^{-1/2}(\mathbf{a}\nabla u - \mathbf{a}_0\nabla u)]_{\dot{B}_{2,1}^{-s}(y+\diamond_n)} &= 3^{d+s} \sum_{k=-\infty}^n 3^{sk} \left( \sum_{z \in y+3^k \mathbb{L}_0 \cap \diamond_n} \left| \mathbf{s}_0^{-1/2} \int_{z+\diamond_k} (\mathbf{a} - \mathbf{a}_0) \nabla u \right|^2 \right)^{1/2} \\
&\leq 2 \cdot 3^{d+s} \sum_{k=-\infty}^n 3^{sk} \left( \sum_{z \in y+3^k \mathbb{L}_0 \cap \diamond_n} \mathcal{U}_{z,k} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(z+\diamond_k)}^2 \right)^{1/2} \\
&\leq 2 \cdot 3^{d+s} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(y+\diamond_n)} \sum_{k=-\infty}^n 3^{sk} \max_{z \in y+3^k \mathbb{L}_0 \cap \diamond_n} \mathcal{U}_{z,k}^{1/2} \quad (6.13)
\end{aligned}$$

and

$$\begin{aligned}
[\mathbf{s}_0^{1/2} \nabla u]_{\dot{B}_{2,1}^{-s}(y+\diamond_n)} &\leq 2 \cdot 3^{d+s} \sum_{k=-\infty}^n 3^{sk} \left( 2 \sum_{z \in y+3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{s}_0 \mathbf{s}_*^{-1}(z+\diamond_k)| \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(z+\diamond_k)}^2 \right)^{1/2} \\
&\leq 2 \cdot 3^{d+s} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(y+\diamond_n)} \sum_{k=-\infty}^n 3^{sk} \max_{z \in y+3^k \mathbb{L}_0 \cap \diamond_n} |\mathbf{s}_0^{1/2} \mathbf{s}_*^{-1/2}(z+\diamond_k)|. \quad (6.14)
\end{aligned}$$

Since

$$|\mathbf{s}_0^{1/2} \mathbf{s}_*^{-1/2}(z+\diamond_k)| \leq 1 + |\mathbf{s}_0^{-1/2} \mathbf{s}_*^{1/2}(z+\diamond_k) - \mathbf{s}_0^{1/2} \mathbf{s}_*^{-1/2}(z+\diamond_k)|.$$

Comparing the definition (6.12) to (6.1), then (6.13) and (6.14) complete the proof, recalling the definition of  $\mathcal{E}_s(y+\diamond_n)$  in (6.2).  $\square$

Note that (6.9), (6.10) and the triangle inequality imply the existence of  $C(d) < \infty$  such that, for every  $X \in \mathcal{S}(y+\diamond_n)$ ,

$$s 3^{-sn} [\mathbf{A}_0^{1/2} X]_{\dot{B}_{2,1}^{-s}(y+\diamond_n)} \leq C(1 + \mathcal{E}_s(y+\diamond_n; \mathbf{a}, \mathbf{a}_0)) \|\mathbf{A}^{1/2} X\|_{\underline{L}^2(y+\diamond_n)}. \quad (6.15)$$

As an immediate consequence of the previous lemma we obtain the first version of the large-scale Caccioppoli estimate.

**Lemma 6.2** (Large-scale Caccioppoli inequality, Version 1). *There exists a constant  $C(d) < \infty$  such that, for every  $m \in \mathbb{N}$ ,  $n \in \mathbb{Z}$  with  $n < m - 2$  and  $u \in \mathcal{A}(\diamond_m)$ , we have the estimate*

$$\begin{aligned}
&\|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_{m-1})} \\
&\leq C \left( 1 + \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_{1/2}(y+\diamond_n; \mathbf{a}, \mathbf{a}_0) \right) \left( \lambda_0^{1/2} 3^{-m} \|u\|_{\underline{L}^2(\diamond_m)} + 3^{-(m-n)} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)} \right). \quad (6.16)
\end{aligned}$$

*Proof.* We assume that  $\mathbf{a}_0 = \mathbf{s}_0$  and denote  $\Theta_n := 1 + \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_{1/2}(y+\diamond_n; \mathbf{a}, \mathbf{a}_0)$ . Let  $\varphi \in C_c^\infty(\diamond_m)$  be such that  $0 \leq \varphi \leq 1$ ,  $\varphi = 1$  in  $\diamond_{m-1}$  and  $3^{jm} \|(\mathbf{q}_0 \nabla)^j \varphi\|_{L^\infty(\diamond_m)} \leq C$  for  $j \in \{1, 2\}$ .

Testing the equation of  $u$  with  $u\varphi^2$ , using the duality between  $B_{2,1}^{-1/2}$  and  $B_{2,\infty}^{1/2}$ , and then applying Lemma 6.1 yields

$$\begin{aligned}
\int_{\diamond_m} |\mathbf{s}^{1/2} \nabla u|^2 \varphi^2 &= - \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \int_{z+\diamond_n} \mathbf{a} \nabla u \cdot (u \nabla \varphi^2) \\
&\leq \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} [\mathbf{s}_0^{-1/2} \mathbf{a} \nabla u]_{\dot{B}_{2,1}^{-1/2}(z+\diamond_n)} \|u \mathbf{s}_0^{1/2} \nabla \varphi^2\|_{\dot{B}_{2,\infty}^{1/2}(z+\diamond_n)} \\
&\leq C \Theta_n \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(z+\diamond_n)} 3^{n/2} \|u \mathbf{s}_0^{1/2} \nabla \varphi^2\|_{\dot{B}_{2,\infty}^{1/2}(z+\diamond_n)}.
\end{aligned}$$

To estimate the last norm on the right, we apply (2.123), (2.128) and Lemma 6.1 to get

$$\begin{aligned}
& 3^{n/2} [u s_0^{1/2} \nabla \varphi^2]_{\underline{B}_{2,\infty}^{1/2}(z+\diamond_n)} \\
& \leq \sup_{k \in \mathbb{Z} \cap (-\infty, n]} 3^{\frac{1}{2}(k-n)} \sum_{z' \in z+3^n \mathbb{L}_0 \cap \diamond_n} \|s_0^{1/2} (u \nabla \varphi^2 - (u \nabla \varphi^2)_{z'+\diamond_k})\|_{\underline{L}^2(z'+\diamond_k)} \\
& \leq \|s_0^{1/2} \nabla \varphi^2\|_{L^\infty(z+\diamond_n)} 3^{n/2} [u]_{\underline{B}_{2,\infty}^{1/2}(z+\diamond_n)} + C \lambda_0^{1/2} 3^n \|u\|_{\underline{L}^2(z+\diamond_n)} \|(\mathbf{q}_0^2 \nabla^2) \varphi^2\|_{L^\infty(z+\diamond_n)} \\
& \leq C 3^{-m} 3^{n/2} [s_0^{1/2} \nabla u]_{\underline{B}_{2,1}^{-1/2}(z+\diamond_n)} \|\varphi\|_{L^\infty(z+\diamond_n)} + C \lambda_0^{1/2} 3^{n-2m} \|u\|_{\underline{L}^2(z+\diamond_n)} \\
& \leq C 3^{n-m} \Theta_n \|s^{1/2} \nabla u\|_{\underline{L}^2(z+\diamond_n)} \|\varphi\|_{L^\infty(z+\diamond_n)} + C \lambda_0^{1/2} 3^{n-2m} \|u\|_{\underline{L}^2(z+\diamond_n)}
\end{aligned}$$

and

$$|(u s_0^{1/2} \nabla \varphi^2)_{z+\diamond_n}| \leq C \lambda_0^{1/2} \|\varphi\|_{L^\infty(z+\diamond_n)} 3^{-m} \|u\|_{\underline{L}^2(z+\diamond_n)}.$$

Furthermore, since

$$\|s^{1/2} \nabla u\|_{\underline{L}^2(z+\diamond_n)} \|\varphi\|_{L^\infty(z+\diamond_n)} \leq \|\varphi s^{1/2} \nabla u\|_{\underline{L}^2(z+\diamond_n)} + C 3^{n-m} \|s^{1/2} \nabla u\|_{\underline{L}^2(z+\diamond_n)},$$

we get by Hölder's inequality for sums that

$$\begin{aligned}
& \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \|s^{1/2} \nabla u\|_{\underline{L}^2(z+\diamond_n)} \|\varphi\|_{L^\infty(z+\diamond_n)} 3^{-m} \|u\|_{\underline{L}^2(z+\diamond_n)} \\
& \leq C \|\varphi s^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)} 3^{-m} \|u\|_{\underline{L}^2(\diamond_m)} + C 3^{n-m} \|s^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)} 3^{-m} \|u\|_{\underline{L}^2(\diamond_m)},
\end{aligned}$$

and similarly for the other terms. Combining the above estimates and applying Young's inequality completes the proof.  $\square$

We next show that *mollifying* an arbitrary solution of the equation  $\nabla \cdot \mathbf{a} \nabla u = 0$  yields a solution of the constant-coefficient equation  $\nabla \cdot \mathbf{a}_0 \nabla w = 0$ , up to an error which is controlled by the random variables  $\mathcal{E}_1$ . This can already be seen as a quantitative homogenization result. It is natural to use the convolution in the adapted geometry of  $\mathbf{s}_0$ . To make room for the convolution, we define, for each  $m, n \in \mathbb{N}$  with  $n < m - 2$ ,

$$\diamond_m^\circ := \{x \in \diamond_m : \exists n \in \mathbb{N}, x + \diamond_{n+1} \subseteq \diamond_m\}.$$

Note that  $\diamond_m^\circ$  depends on  $n$ , but this is kept implicit in the notation. For  $r \in (3^{n-1}, 3^n]$ , we let  $\eta_r$  be a mollifier adapted to the  $\mathbf{s}_0$  geometry; that is, we select a smooth function  $\eta$  satisfying, for some  $C(d) < \infty$ ,

$$\eta \in C_c^\infty(\diamond_0), \quad \int_{\mathbb{R}^d} \eta = 1, \quad 0 \leq \eta \leq 1, \quad \|\mathbf{q}_0^{1/2} \nabla \eta\|_{L^\infty(\mathbb{R}^d)} \leq C, \quad (6.17)$$

and then define

$$\eta_r := r^{-d} \eta\left(\frac{\cdot}{r}\right), \quad \forall r > 0. \quad (6.18)$$

The next simple lemma shows the coarse-graining for arbitrary solutions.

**Lemma 6.3.** *Let  $m, n \in \mathbb{N}$  with  $n < m - 2$  and  $r \in (3^{n-1}, 3^n]$ . Let  $\eta_r$  be as in (6.17)–(6.18). There exists a constant  $C(d) < \infty$  such that, for every  $u \in \mathcal{A}(\diamond_m)$  and  $y \in 3^n \mathbb{L}_0 \cap \diamond_m^\circ$ , we have that*

$$\|s_0^{-1/2} (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u) * \eta_r\|_{L^\infty(y+\diamond_n)} \leq C \mathcal{E}_1(y+\diamond_{n+1}; \mathbf{a}, \mathbf{a}_0) \|s^{1/2} \nabla u\|_{\underline{L}^2(y+\diamond_{n+1})} \quad (6.19)$$

and

$$\begin{aligned} & \|s_0^{1/2} \nabla u * \eta_r\|_{L^\infty(y+\diamond_n)} + \lambda_0^{1/2} r^{-1} \|u - u * \eta_r\|_{\underline{L}^2(y+\diamond_n)} \\ & \leq C(1 + \mathcal{E}_1(y+\diamond_{n+1}; \mathbf{a}, \mathbf{a}_0)) \|s^{1/2} \nabla u\|_{\underline{L}^2(y+\diamond_{n+1})}. \end{aligned} \quad (6.20)$$

*Proof.* Fix  $y \in 3^n \mathbb{L}_0 \cap \diamond_m^\circ$ . We apply Lemma 6.1 to obtain that

$$\begin{aligned} & \|s_0^{-1/2} (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u) * \eta_r\|_{L^\infty(y+\diamond_n)} \leq \|\eta_r\|_{\underline{B}_{2,\infty}^1(\diamond_{n+1})} \|s_0^{-1/2} (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u)\|_{\underline{B}_{2,1}^{-1}(y+\diamond_{n+1})} \\ & \leq C 3^{-n} \|s_0^{-1/2} (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u)\|_{\underline{B}_{2,1}^{-1}(y+\diamond_{n+1})} \\ & \leq C \mathcal{E}_1(y+\diamond_{n+1}) \|s^{1/2} \nabla u\|_{\underline{L}^2(y+\diamond_{n+1})}. \end{aligned} \quad (6.21)$$

A similar estimate is valid for the gradient. We then also have that, by (2.128) and (2.130),

$$\begin{aligned} & \|u_r - u\|_{\underline{L}^2(y+\diamond_n)} \leq C \|u - (u)_{y+\diamond_{n+1}}\|_{\underline{L}^2(y+\diamond_{n+1})} \\ & \leq C(1 + \mathcal{E}_1(y+\diamond_{n+1})) \lambda_0^{-1/2} r \|s^{1/2} \nabla u\|_{\underline{L}^2(y+\diamond_{n+1})}. \end{aligned}$$

This completes the proof.  $\square$

**Lemma 6.4.** *Let  $m, n \in \mathbb{N}$  with  $n < m - 2$  and  $r \in (3^{n-1}, 3^n]$ . Let  $\eta_r$  be as in (6.17)–(6.18). There exists a constant  $C(d) < \infty$  such that, for every  $s, t \in (0, 1]$  with  $t > s$  and  $f \in B_{2,\infty}^t(\diamond_m)$ ,*

$$\sup \left\{ \left| \int_{\diamond_m} (f * \eta_r - f) v \right| : v \in C_c^\infty(\diamond_m^\circ), [v]_{\underline{B}_{2,\infty}^t(\diamond_m)} \leq 1 \right\} \leq C 3^{(t-s)n} [f]_{\underline{B}_{2,1}^{-s}(\diamond_m)}. \quad (6.22)$$

*Proof.* Fix  $v \in C_c^\infty(\diamond_m^\circ)$  such that  $[v]_{\underline{B}_{2,\infty}^t(\diamond_m)} \leq 1$ . Then, by Fubini's theorem and duality,

$$\left| \int_{\diamond_m} (f * \eta_r - f) v \right| = \left| \int_{\diamond_m} f(v * \eta_r - v) \right| \leq [f]_{\underline{B}_{2,1}^{-s}(\diamond_m)} \|v * \eta_r - v\|_{\underline{B}_{2,\infty}^s(\diamond_m)}.$$

The seminorm above is of the form

$$\|v * \eta_r - v\|_{\underline{B}_{2,\infty}^s(\diamond_m)} = \sup_{j \in (-\infty, m] \cap \mathbb{N}} 3^{-js} \left( \sum_{z \in 3^j \mathbb{L}_0 \cap \diamond_m} \|v * \eta_r - v - (v * \eta_r - v)_{z+\diamond_j}\|_{\underline{L}^2(z+\diamond_j)}^2 \right)^{1/2}.$$

On the one hand, for  $j, n \in \mathbb{N}$  with  $j \geq n$  we have

$$\begin{aligned} & \sum_{z \in 3^j \mathbb{L}_0 \cap \diamond_m} \|v * \eta_r - v - (v * \eta_r - v)_{z+\diamond_j}\|_{\underline{L}^2(z+\diamond_j)}^2 \\ & \leq C \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \|v - (v)_{z+\diamond_{n+1}}\|_{\underline{L}^2(z+\diamond_{n+1})}^2 \leq C 3^{2tn} [v]_{\underline{B}_{2,\infty}^t(\diamond_m)}^2 \leq C 3^{2tn}. \end{aligned}$$

On the other hand, for  $j < n$  we obtain

$$\sum_{z \in 3^j \mathbb{L}_0 \cap \diamond_m} \|v * \eta_r - v - (v * \eta_r - v)_{z+\diamond_j}\|_{\underline{L}^2(z+\diamond_j)}^2 \leq C 3^{2tj} [v]_{\underline{B}_{2,\infty}^t(\diamond_m)}^2.$$

Therefore, we have the estimate

$$\|v * \eta_r - v\|_{\underline{B}_{2,\infty}^s(\diamond_m)} \leq C 3^{(t-s)n} [v]_{\underline{B}_{2,\infty}^t(\diamond_m)} \leq C 3^{(t-s)n},$$

and the result follows.  $\square$

The next lemma compares the energies of the mollified solution to the energy of the original solution.

**Lemma 6.5** (Coarse-graining the energy). *Let  $m, n \in \mathbb{N}$  with  $n < m - 2$  and  $r \in (3^{n-1}, 3^n]$ . Let  $\eta_r$  be as in (6.17)–(6.18). There exists a constant  $C(d) < \infty$  such that, for every  $u \in \mathcal{A}(\diamond_m)$  and  $\varphi \in C_0^\infty(\diamond_m - \diamond_n)$  satisfying  $\max_{j \in \{0,1,2\}} 3^{jm} \|\mathbf{q}_0^j \nabla^j \varphi\|_{L^\infty(\diamond_m)} \leq 1$  we have, for every  $s \in (0, 1)$ ,*

$$\left| \int_{\diamond_m} \varphi^2 (\nabla u \cdot \mathbf{s} \nabla u - \nabla(u * \eta_r) \cdot \mathbf{s}_0 \nabla(u * \eta_r)) \right| \leq C \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \left( \mathcal{E}_1(y + \diamond_n; \mathbf{a}, \mathbf{a}_0) + 3^{-(1-s)(m-n)} (1 + \mathcal{E}_s(y + \diamond_n; \mathbf{a}, \mathbf{a}_0)) \right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}^2. \quad (6.23)$$

*Proof.* Assume, without loss of generality, that  $\mathbf{k}_0$  vanishes, so that  $\mathbf{a}_0 = \mathbf{s}_0$ . Fix  $m, n \in \mathbb{N}$  with  $n < m - 2$  and  $r \in (3^{n-1}, 3^n]$ . We may assume that

$$\max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \left( \mathcal{E}_1(y + \diamond_n; \mathbf{a}, \mathbf{a}_0) + 3^{-(1-s)(m-n)} (1 + \mathcal{E}_s(y + \diamond_n; \mathbf{a}, \mathbf{a}_0)) \right) \leq 1, \quad (6.24)$$

since otherwise the result follows by the triangle inequality and Lemma 6.3.

Denote  $u_r := \eta_r * u$  and  $\diamond_m^\circ := \diamond_m - \diamond_n$ . Fix also  $u \in \mathcal{A}(\diamond_m)$  and  $\varphi \in C_0^\infty(\diamond_m^\circ)$  such that  $\max_{j \in \{0,1,2\}} 3^{jm} \|\mathbf{q}_0^j \nabla^j \varphi\|_{L^\infty(\diamond_m)} \leq 1$ .

To compare the energy density  $\nabla u \cdot \mathbf{s} \nabla u$  to the “coarse-grained” energy density  $\nabla u_r \cdot \mathbf{s}_0 \nabla u_r$ , we use the following decomposition for their difference:

$$\begin{aligned} \nabla u \cdot \mathbf{s} \nabla u - \nabla u_r \cdot \mathbf{s}_0 \nabla u_r \\ = \nabla u_r \cdot (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u) * \eta_r + \nabla u_r \cdot (\mathbf{a} \nabla u - (\mathbf{a} \nabla u) * \eta_r) + \nabla(u - u_r) \cdot \mathbf{a} \nabla u. \end{aligned} \quad (6.25)$$

First, to estimate the contribution of the first term on the right side of (6.25), we use Lemma 6.3, the Cauchy-Schwarz inequality and (6.3) to get

$$\begin{aligned} \left| \int_{\diamond_m} \varphi^2 \nabla u_r \cdot (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u) * \eta_r \right| &\leq \frac{|\diamond_m^\circ|}{|\diamond_m|} \sum_{y \in 3^n \mathbb{L}_0 \cap \diamond_m^\circ} \left| \int_{y + \diamond_n} \varphi^2 \nabla u_r \cdot (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u) * \eta_r \right| \\ &\leq C \sum_{y \in 3^n \mathbb{L}_0 \cap \diamond_m^\circ} \mathcal{E}_1(y + \diamond_{n+1}) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(y + \diamond_{n+1})}^2 \\ &\leq C \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}^2. \end{aligned} \quad (6.26)$$

Second, we integrate by parts, using that  $\nabla \cdot \mathbf{a} \nabla u = 0 = \nabla \cdot ((\mathbf{a} \nabla u) * \eta_r)$ , which is valid by Lemma 2.13, to obtain that

$$\int_{\diamond_m} \varphi^2 \nabla u_r \cdot (\mathbf{a} \nabla u - (\mathbf{a} \nabla u) * \eta_r) = - \int_{\diamond_m} (u_r - (u_r)_{\diamond_m^\circ}) \nabla \varphi^2 \cdot (\mathbf{a} \nabla u - (\mathbf{a} \nabla u) * \eta_r).$$

By Lemma 6.4 we deduce that

$$\begin{aligned} \left| \int_{\diamond_m} (u_r - (u_r)_{\diamond_m^\circ}) \nabla \varphi^2 \cdot (\mathbf{a} \nabla u - (\mathbf{a} \nabla u) * \eta_r) \right| \\ \leq 2 \|(u_r - (u_r)_{\diamond_m^\circ}) \mathbf{s}_0^{1/2} \nabla \varphi\|_{\underline{B}_{2,\infty}^1(\diamond_m)} [\varphi \mathbf{s}_0^{-1/2} (\mathbf{a} \nabla u - (\mathbf{a} \nabla u) * \eta_r)]_{\underline{B}_{2,1}^{-1}(\diamond_m)} \\ \leq C r^{1-s} \|(u_r - (u_r)_{\diamond_m^\circ}) \mathbf{s}_0^{1/2} \nabla \varphi\|_{\underline{B}_{2,\infty}^1(\diamond_m)} [\mathbf{s}_0^{-1/2} \mathbf{a} \nabla u]_{\underline{B}_{2,1}^{-s}(\diamond_m)}. \end{aligned}$$

By (6.9), (6.10) and (6.24) we have

$$[\mathbf{s}_0^{-1/2} \mathbf{a} \nabla u]_{\underline{B}_{2,1}^s(\diamond_m)} \leq C 3^{sm} (1 + \mathcal{E}_s(\diamond_m)) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \quad (6.27)$$

Since  $u_r \mathbf{s}_0^{1/2} \nabla \varphi$  is smooth, we have by the Poincaré inequality (2.129) and the bound for  $\varphi$  that

$$\begin{aligned} \|(u_r - (u_r)_{\diamond_m^\circ}) \mathbf{s}_0^{1/2} \nabla \varphi\|_{\underline{B}_{2,\infty}^1(\diamond_m)} &\leq C \lambda_0^{1/2} \|(u_r - (u_r)_{\diamond_m^\circ}) \mathbf{q}_0 \nabla \varphi\|_{\underline{B}_{2,\infty}^1(\diamond_m)} \\ &\leq C \lambda_0^{1/2} 3^{-m} (3^{-m} \|u_r - (u_r)_{\diamond_m^\circ}\|_{\underline{L}^2(\diamond_m^\circ)} + \|\mathbf{q}_0 \nabla u_r\|_{\underline{L}^2(\diamond_m^\circ)}) \\ &\leq C 3^{-m} \|\mathbf{s}_0^{1/2} \nabla u_r\|_{\underline{L}^2(\diamond_m^\circ)}. \end{aligned}$$

Hence, by (6.20) and (6.24),

$$\|(u_r - (u_r)_{\diamond_m^\circ}) \mathbf{s}_0^{1/2} \nabla \varphi\|_{\underline{B}_{2,\infty}^1(\diamond_m)} \leq C 3^{-m} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}.$$

Putting the last four displays together yields

$$\left| \int_{\diamond_m} \varphi \nabla u_r \cdot (\mathbf{a} \nabla u - (\mathbf{a} \nabla u) * \eta_r) \right| \leq C 3^{-(1-s)(m-n)} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}^2. \quad (6.28)$$

Third, by integration by parts and duality, we have that, for  $w_r := u - u_r$ ,

$$\left| \int_{\diamond_m} \varphi \nabla(u - u_r) \cdot \mathbf{a} \nabla u \right| = \left| \int_{\diamond_m} w_r \nabla \varphi \cdot \mathbf{a} \nabla u \right| \leq \|w_r \mathbf{s}_0^{1/2} \nabla \varphi\|_{\underline{B}_{2,\infty}^s(\diamond_m)} [\mathbf{s}_0^{-1/2} \mathbf{a} \nabla u]_{\underline{B}_{2,1}^s(\diamond_m)}.$$

Notice that, by (6.20) and (6.24), we have

$$|(w_r \mathbf{s}_0^{1/2} \nabla \varphi)_{\diamond_m^\circ}| \leq C \lambda_0^{1/2} 3^{-m} \|w_r\|_{\underline{L}^2(\diamond_m^\circ)} \leq C 3^{-(m-n)} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \quad (6.29)$$

We finally claim that

$$3^{sm} [w_r \mathbf{s}_0^{1/2} \nabla \varphi]_{\underline{B}_{2,\infty}^s(\diamond_m)} \leq C 3^{-(1-s)(m-n)} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \quad (6.30)$$

Before the proof, we demonstrate how this leads to the result. Together with (6.27), (6.30) yields

$$\left| \int_{\diamond_m} \varphi \nabla(u - u_r) \cdot \mathbf{a} \nabla u \right| \leq C 3^{-(1-s)(m-n)} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}^2. \quad (6.31)$$

Combining (6.25), (6.26), (6.28) and (6.31) then completes the proof of (6.23).

To prove (6.30), we denote  $\mathcal{Z}_k := 3^{k-1} \mathbb{L}_0 \cap \diamond_m^\circ$ , and recall that

$$[w_r \mathbf{s}_0^{1/2} \nabla \varphi]_{\underline{B}_{2,\infty}^s(\diamond_m)} = \sup_{k \in (-\infty, m] \cap \mathbb{Z}} 3^{-sk} \left( \sum_{z \in \mathcal{Z}_k} \|w_r \mathbf{s}_0^{1/2} \nabla \varphi - (w_r \mathbf{s}_0^{1/2} \nabla \varphi)_{z+\diamond_k}\|_{\underline{L}^2(z+\square_k)}^2 \right)^{1/2}.$$

On the one hand, we trivially have that

$$\sup_{k \in [n, m] \cap \mathbb{N}} 3^{-sk} \left( \sum_{z \in \mathcal{Z}_k} \|w_r \mathbf{s}_0^{1/2} \nabla \varphi - (w_r \mathbf{s}_0^{1/2} \nabla \varphi)_{z+\diamond_k}\|_{\underline{L}^2(z+\diamond_k)}^2 \right)^{1/2} \leq C 3^{-m-sn} \lambda_0^{1/2} \|w_r\|_{\underline{L}^2(\diamond_m^\circ)}.$$

On the other hand, following Step 2 of the proof of Lemma A.3, we see that

$$\begin{aligned} & \sup_{k \in (-\infty, n] \cap \mathbb{Z}} 3^{-sk} \left( \sum_{z \in \mathcal{Z}_k} \|w_r \mathbf{s}_0^{1/2} \nabla \varphi - (w_r \mathbf{s}_0^{1/2} \nabla \varphi)_{z+\diamond_k}\|_{\underline{L}^2(z+\diamond_k)}^2 \right)^{1/2} \\ & \leq C \lambda_0^{1/2} 3^{(1-s)n-2m} \|w_r\|_{\underline{L}^2(\diamond_m^\circ)} + C \lambda_0^{1/2} 3^{-m} \sup_{k \in (-\infty, n] \cap \mathbb{Z}} 3^{-sk} \left( \sum_{z \in \mathcal{Z}_k} \|w_r - (w_r)_{z+\diamond_k}\|_{\underline{L}^2(z+\diamond_k)}^2 \right)^{1/2}. \end{aligned}$$

Now (6.20) and (6.24) imply that

$$\|w_r\|_{\underline{L}^2(\diamond_m^\circ)} \leq C \lambda_0^{-1/2} 3^n \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}.$$

By (2.128), (6.10) and (6.24) we get

$$\begin{aligned} 3^{-sk} \|w_r - (w_r)_{z+\diamond_k}\|_{\underline{L}^2(z+\diamond_k)} & \leq C 3^{(1-s)k} \|\mathbf{q}_0^{1/2} \nabla w_r\|_{\underline{L}^2(z+\diamond_k)} \\ & \leq C 3^{-(1-s)k-n} \|u - (u)_{\diamond_{k+1}}\|_{\underline{L}^2(z+\diamond_{k+1})} \\ & \leq C 3^{-(1-s)k-n} [\mathbf{q}_0^{1/2} \nabla u]_{\dot{B}_{2,1}^{-1}(z+\diamond_{k+1})} \\ & \leq C 3^{(2-s)k-n} \lambda_0^{-1/2} (1 + \mathcal{E}_1^*(z+\diamond_{k+1})) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(z+\diamond_{k+1})} \\ & \leq C 3^{-2(1-s)(n-k)+(1-s)n} \lambda_0^{-1/2} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(z+\diamond_{k+1})}. \end{aligned}$$

Therefore, by combining the previous five displays, we obtain

$$[w_r \mathbf{s}_0^{1/2} \nabla \varphi]_{\dot{B}_{2,\infty}^s(\diamond_m)} \leq C 3^{-(1-s)(m-n)} 3^{-sm} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)},$$

which gives (6.30). This completes the proof.  $\square$

We use the previous lemma to obtain a large-scale version of the Caccioppoli inequality which does not lose unnecessary factors of  $\Theta$  or  $\Pi$ .

**Lemma 6.6** (Large-scale Caccioppoli inequality, Version 2). *There exists a constant  $C(d) < \infty$  such that, for every  $m \in \mathbb{N}$ ,  $n \in \mathbb{Z}$  with  $n < m - 2$ ,  $s \in (0, 1)$  and  $u \in \mathcal{A}(\diamond_m)$ , we have the estimate*

$$\begin{aligned} & \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_{m-1})} \leq C \lambda_0^{1/2} 3^{-m} \|u\|_{\underline{L}^2(\diamond_m)} \\ & + C \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \left( \mathcal{E}_1(y+\diamond_n; \mathbf{a}, \mathbf{a}_0) + 3^{-s(m-n)} (1 + \mathcal{E}_{1-s}(y+\diamond_n; \mathbf{a}, \mathbf{a}_0)) \right)^{1/2} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \end{aligned} \quad (6.32)$$

*Proof.* We assume that  $\mathbf{a}_0 = \mathbf{s}_0$  and  $\max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y+\diamond_n; \mathbf{a}, \mathbf{a}_0) \leq 1$ . Both  $\mathbf{a} \nabla u$  and  $(\mathbf{a} \nabla u) * \eta_r$  are divergence-free, and hence we have

$$-\nabla \cdot \mathbf{a}_0 \nabla u_r = \nabla \cdot (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u) * \eta_r \quad \text{in } \diamond_m^\circ.$$

Testing this equation with  $\varphi u_r$ , where  $\varphi$  is as in the statement of Lemma 6.5 satisfying in addition  $\varphi = 1$  in  $\diamond_{m-1}$ , we therefore obtain

$$\begin{aligned} \int_{\diamond_m} \varphi^2 \nabla u_r \cdot \mathbf{s}_0 \nabla u_r & = \int_{\diamond_m} (\nabla(\varphi^2 u_r) - 2u_r \varphi \nabla \varphi) \cdot \mathbf{a}_0 \nabla u_r \\ & = \int_{\diamond_m} (\nabla(\varphi^2 u_r) \cdot (\mathbf{a}_0 \nabla u - \mathbf{a} \nabla u) * \eta_r) - 2u_r \varphi \nabla \varphi \cdot \mathbf{s}_0 \nabla u_r \\ & = \int_{\diamond_m} (\varphi(2u_r \nabla \varphi + \varphi^2 \nabla u_r) \cdot (\mathbf{a}_0 \nabla u - \mathbf{a} \nabla u) * \eta_r - 2(u_r \varphi \mathbf{s}_0^{1/2} \nabla \varphi) \cdot (\varphi \mathbf{s}_0^{1/2} \nabla u_r)). \end{aligned}$$



Applying Cauchy-Schwarz and then Young's inequality, we thus deduce that

$$\begin{aligned} & \|\varphi \mathbf{s}_0^{1/2} \nabla u_r\|_{\underline{L}^2(\diamond_m)}^2 \\ & \leq \frac{1}{2} \|\varphi \mathbf{s}_0^{1/2} \nabla u_r\|_{\underline{L}^2(\diamond_m)}^2 + C \|u_r \mathbf{s}_0^{1/2} \nabla \varphi\|_{\underline{L}^2(\diamond_m)}^2 + C \|\varphi \mathbf{s}_0^{-1/2} (\mathbf{a}_0 \nabla u - \mathbf{a} \nabla u) * \eta_r\|_{\underline{L}^2(\diamond_m)}^2. \end{aligned}$$

By reabsorbing the first term and using (6.19), which says that

$$\|\varphi (\mathbf{a}_0 \nabla u - \mathbf{a} \nabla u) * \eta_r\|_{\underline{L}^2(\diamond_m)}^2 \leq C \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1^2(y + \diamond_n) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}^2,$$

we obtain

$$\|\varphi \mathbf{s}_0^{1/2} \nabla u_r\|_{\underline{L}^2(\diamond_m)} \leq C 3^{-m} \lambda_0^{1/2} \|u_r\|_{\underline{L}^2(\diamond_m)} + C \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}.$$

We obtain (6.32) from the this, Lemma 6.5 and (6.20), completing the proof.  $\square$

**6.2. Harmonic approximation estimates.** In this section we discuss two different harmonic approximation results together with the homogenization estimate for the Dirichlet problem. We start with the simplest of the results, namely that every heterogeneous solution can be approximated by the homogenized solution locally. This result is crucial when proving large-scale regularity results for the solution in the next subsection.

**Proposition 6.7** (Harmonic approximation I). *Let  $n \in \mathbb{N}$  with  $n < m - 2$ ,  $r \in (3^{n-1}, 3^n]$  and  $\eta_r$  be as in (6.17)–(6.18). For every  $u \in \mathcal{A}(\diamond_m)$  there exists an  $\mathbf{a}_0$ -harmonic function  $u_{\text{hom}}$  in  $\diamond_{m-1}$  satisfying the following properties. There exists a constant  $C(d) < \infty$  such that*

$$3^{-m} \lambda_0^{1/2} \|u - u_{\text{hom}}\|_{\underline{L}^2(\diamond_{m-1})} \leq C \left( 3^{-(m-n)} + \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n; \mathbf{a}, \mathbf{a}_0) \right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \quad (6.33)$$

For the weak norms we have, for every  $s \in (0, 1]$  and  $\theta \in (0, s)$ ,

$$\begin{aligned} & s 3^{-sm} [\mathbf{s}_0^{1/2} (\nabla u - \nabla u_{\text{hom}})]_{\underline{B}_{2,1}^{-s}(\diamond_{m-1})} + s 3^{-sm} [\mathbf{s}_0^{-1/2} ((\mathbf{a} - \mathbf{k}_0) \nabla u - \mathbf{s}_0 \nabla u_{\text{hom}})]_{\underline{B}_{2,1}^{-s}(\diamond_{m-1})} \\ & \leq C \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \left( \mathcal{E}_1(y + \diamond_n) + 3^{-\theta(m-n)} (1 + \mathcal{E}_{s-\theta}(y + \diamond_n)) \right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \end{aligned} \quad (6.34)$$

Finally, for every  $p \in [2, \infty)$ , there exists a constant  $C(p, d) < \infty$  such that

$$\begin{aligned} & 3^{-m} \lambda_0^{1/2} \|u * \eta_r - u_{\text{hom}}\|_{L^\infty(\diamond_{m-1})} + \|\mathbf{s}_0^{1/2} \nabla (u * \eta_r - u_{\text{hom}})\|_{\underline{L}^p(\diamond_{m-1})} \\ & \leq C \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n; \mathbf{a}, \mathbf{a}_0) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \end{aligned} \quad (6.35)$$

*Proof.* Without loss of generality, we may take  $\mathbf{k}_0 = 0$ . Denote  $u_r := \eta_r * u$ . Let  $u_{\text{hom}} \in H^1(\diamond_{m-1})$  be the solution of

$$\begin{cases} -\nabla \cdot \mathbf{a}_0 \nabla u_{\text{hom}} = 0 & \text{in } \diamond_{m-1}, \\ u_{\text{hom}} = u_r & \text{on } \partial \diamond_{m-1}. \end{cases}$$

Notice that  $u_r$  solves the equation

$$-\nabla \cdot \mathbf{a}_0 \nabla u_r = \nabla \cdot (\eta_r * (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u)).$$

Now, if  $\tilde{w} \in H_0^1(\diamond_{m-1})$  solves  $-\nabla \cdot \mathbf{q}_0 \nabla \tilde{w} = \nabla \cdot \tilde{\mathbf{f}}$  in  $\diamond_{m-1}$ , we have, after rescaling, by the classical Calderón-Zygmund estimates that, for every  $p \in (1, \infty)$ , there exists  $C(p, d) < \infty$  such that

$$\|\mathbf{q}_0 \nabla \tilde{w}\|_{\underline{L}^p(\diamond_{m-1})} \leq C \|\mathbf{q}_0^{-1} \tilde{\mathbf{f}}\|_{\underline{L}^p(\diamond_{m-1})}. \quad (6.36)$$

Applying this for  $\tilde{w} := u_r - u_{\text{hom}}$ , the above estimate implies that

$$\|\mathbf{s}_0^{1/2} \nabla (u_r - u_{\text{hom}})\|_{\underline{L}^p(\diamond_{m-1})} \leq C \|\mathbf{s}_0^{-1/2} \eta_r * (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u)\|_{\underline{L}^p(\diamond_{m-1})}. \quad (6.37)$$

Now (6.19) yields

$$\|\mathbf{s}_0^{-1/2} \eta_r * (\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u)\|_{L^\infty(\diamond_{m-1})} \leq C \max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(z + \diamond_n) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}$$

By Morrey's inequality we get

$$3^{-m} \lambda_0^{1/2} \|u_r - u_{\text{hom}}\|_{L^\infty(\diamond_{m-1})} \leq C \max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(z + \diamond_n) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)},$$

and thus (6.35) follows by the last three displays. Next, by (6.20) we have

$$\lambda_0^{1/2} \|u - u_r\|_{\underline{L}^2(\diamond_{m-1})} \leq C 3^n \left(1 + \max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(z + \diamond_n)\right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}.$$

The last two displays give us (6.33) by the triangle inequality.

By the triangle inequality, (6.37), (6.15) and Lemma 6.4, we get, for every  $s \in (0, 1]$  and  $\theta \in (0, s)$ ,

$$\begin{aligned} & s 3^{-sm} \left[ \mathbf{A}_0^{1/2} \left( \begin{array}{c} \nabla u - \nabla u_{\text{hom}} \\ \mathbf{a} \nabla u - \mathbf{s}_0 \nabla u_{\text{hom}} \end{array} \right) \right]_{\underline{B}_{2,1}^{-s}(\diamond_{m-1})} \\ & \leq s 3^{-sm} \left[ \left( \begin{array}{c} \mathbf{s}_0^{1/2} (\nabla u_r - \nabla u) \\ (\mathbf{s}_0^{-1/2} (\mathbf{a} - \mathbf{s}_0) \nabla u) * \eta_r \end{array} \right) \right]_{\underline{B}_{2,1}^{-s}(\diamond_{m-1})} + C \|\mathbf{s}_0^{1/2} (\nabla u_r - \nabla u_{\text{hom}})\|_{\underline{L}^2(\diamond_{m-1})} \\ & \leq C \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \left( \mathcal{E}_1(y + \diamond_n) + 3^{-\theta(m-n)} (1 + \mathcal{E}_{s-\theta}(y + \diamond_n)) \right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}, \end{aligned}$$

which gives us (6.34). The proof is complete.  $\square$

With a modification of the previous proof, we obtain a generalization of the previous proposition for more general domains and general boundary values. Notice that in the statement the last term in (6.40) is small only if the convolution of the solution and  $g$  are very close to each other in the boundary layer. This can be guaranteed, for example, if there is some kind of extra quantitative property on the behavior of the coefficients such as pointwise ellipticity. Then if  $u = g$  at the boundary, the last term can be estimated a boundary Poincaré inequality, and the geometric factor  $3^{-(1/d \wedge 1/6)(m-n)}$  provides smallness against for possibly very large pointwise ellipticity ratio. There are many other similar situations, and thus we leave the estimate in such a form that it can be applied to every possible solution  $u$  in  $U$ .

Given a Lipschitz domain  $U$  and  $n \in \mathbb{N}$ , denote

$$Z_n(U) := \{z \in 3^{n+4} \mathbb{L}_0 : z + \diamond_{n+4} \subseteq U\} \quad \text{and} \quad U_n^\circ := \bigcup_{z \in Z_n(U)} \overline{(z + \diamond_n)}. \quad (6.38)$$

**Proposition 6.8** (Homogenization of the Dirichlet problem). *Let  $U$  be a Lipschitz domain and let  $m \in \mathbb{N}$  be the smallest integer larger than 2 such that  $U$  belongs to  $\Diamond_m$ . Let  $n \in \mathbb{N}$  with  $n < m - 2$  and  $r \in (3^{n-1}, 3^n]$ , and let  $\eta_r$  be as in (6.17)–(6.18). There exist a constant  $C(d, \mathbf{s}_0^{-1/2}U) < \infty$  such that, if  $g \in H^1(U)$  and  $u_{\text{hom}}$  solves*

$$\begin{cases} -\nabla \cdot \mathbf{a}_0 \nabla u_{\text{hom}} = 0 & \text{in } U, \\ u_{\text{hom}} = g & \text{on } \partial U, \end{cases}$$

then, for every  $u \in \mathcal{A}(U)$ , we have the estimate

$$\begin{aligned} 3^{-m} \lambda_0^{1/2} \|u - u_{\text{hom}}\|_{\underline{L}^2(U_{n+1}^\circ)} &\leq C \left( 3^{-(1/d \wedge 1/6)(m-n)} + \max_{y \in 3^n \mathbb{L}_0 \cap \Diamond_m} \mathcal{E}_1(y + \Diamond_n) \right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(U_n^\circ)} \\ &\quad + C 3^{-(1/d \wedge 1/6)(m-n)} \left( \|\mathbf{s}_0^{1/2} \nabla g\|_{\underline{L}^2(U)} + \frac{\|u * \eta_r - g\|_{L^2(U_n^\circ \setminus U_{n+1}^\circ)}}{3^n |U|^{1/2}} \right). \end{aligned} \quad (6.39)$$

Moreover, if  $k \in \mathbb{N}$ ,  $y \in \mathbb{R}^d$  are such that  $y + \Diamond_{k+1} \subseteq U_{n+2}^\circ$ , then, for every  $u \in \mathcal{A}(U)$ ,  $s \in (0, 1]$  and  $\theta \in (0, s)$ , we have

$$\begin{aligned} s 3^{-sk} 3^{-(d/2+1)(m-k)} &\left[ \overline{\mathbf{A}}^{1/2} \begin{pmatrix} \nabla u - \nabla u_{\text{hom}} \\ \mathbf{a} \nabla u - \mathbf{s}_0 \nabla u_{\text{hom}} \end{pmatrix} \right]_{\underline{B}_{2,1}^{-s}(y + \Diamond_k)} \\ &\leq C \max_{y \in 3^n \mathbb{L}_0 \cap \Diamond_m} \left( \mathcal{E}_1(y + \Diamond_n; \mathbf{a}, \mathbf{a}_0) + 3^{-(1/d \wedge 1/6 \wedge \theta)(k-n)} (1 + \mathcal{E}_{s-\theta}(y + \Diamond_n; \mathbf{a}, \mathbf{a}_0)) \right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(U_n^\circ)} \\ &\quad + C 3^{-(1/d \wedge 1/6)(m-n)} \left( \|\mathbf{s}_0^{1/2} \nabla g\|_{\underline{L}^2(U)} + \frac{\|u * \eta_r - g\|_{L^2(U_{n+1}^\circ \setminus U_{n+2}^\circ)}}{r |U|^{1/2}} \right). \end{aligned} \quad (6.40)$$

*Proof.* Let  $n \in \mathbb{N}$  and let  $U$  be a smooth or a Lipschitz domain and assume that  $m \in \mathbb{N}$  is the smallest integer larger than  $n + 4$  such that  $U$  belongs to  $\Diamond_m$ . Without loss of generality, by removing the anti-symmetric part of  $\mathbf{a}_0$ , we may assume that  $\mathbf{a}_0 = \mathbf{s}_0$ . Let  $r \in (3^{n-1}, 3^n]$  and denote  $u_r := u * \eta_r$ . Let  $\zeta_r$  be a smooth cut-off function satisfying  $\mathbf{1}_{U_{n+1}^\circ} \leq \zeta_r \leq \mathbf{1}_{U_n^\circ}$  and  $\|\mathbf{q}_0^{1/2} \nabla \zeta_r\|_{L^\infty(U)} \leq C r^{-1}$ . We define

$$w_r := \zeta_r u_r + (1 - \zeta_r) g.$$

Let  $u_{\text{hom}} \in H^1(U)$  solve

$$\begin{cases} -\nabla \cdot \mathbf{a}_0 \nabla u_{\text{hom}} = 0 & \text{in } U, \\ u_{\text{hom}} = g & \text{on } \partial U. \end{cases}$$

We will compare  $w_r$  to  $u_{\text{hom}}$ , and then to  $u$  and deduce estimates on  $u - u_{\text{hom}}$  by the triangle inequality. The difference  $w_r - u_{\text{hom}}$  is estimated using the coarse-graining lemma (Lemma 6.3) and a crude boundary layer estimate. The difference  $w_r - u$  is estimated using the closeness of  $u_r$  to  $u$ , which follows from a scale separation argument like in the proof of Lemma 6.5, and another (similar) boundary layer estimate.

To estimate  $w_r - u_{\text{hom}}$ , we write its equation as follows:

$$\begin{aligned} \nabla \cdot \mathbf{a}_0 \nabla (w_r - u_{\text{hom}}) &= \nabla \cdot (\zeta_r (\mathbf{a}_0 \nabla u - \mathbf{a} \nabla u) * \eta_r) + \nabla \zeta_r \cdot (\mathbf{a} \nabla u) * \eta_r \\ &\quad + \nabla \cdot ((u_r - g) \mathbf{s}_0 \nabla \zeta_r + (1 - \zeta_r) \mathbf{s}_0 \nabla g). \end{aligned} \quad (6.41)$$

Now  $w_r - u_{\text{hom}}$  vanishes on the boundary and can be extended to be zero outside of  $U_n^\circ$ . After rescaling, if  $\tilde{w} \in H_0^1(U)$  solves  $-\nabla \cdot \mathbf{q}_0 \nabla \tilde{w} = \nabla \cdot \tilde{\mathbf{f}} + \tilde{g}$  in  $U$ , we have, after rescaling, by the classical Calderón-Zygmund estimates that

$$\|\mathbf{q}_0^{1/2} \nabla \tilde{w}\|_{\underline{L}^p(U)} \leq C \|\mathbf{q}_0^{-1/2} \tilde{\mathbf{f}}\|_{\underline{L}^p(U)} + C \sup \left\{ \iint_U \tilde{g} \psi : \|\mathbf{q}_0^{1/2} \nabla \psi\|_{\underline{L}^p(U)} \leq 1 \right\}.$$

We denote the last quantity on the right by  $\|\tilde{g}\|_{\underline{W}_{\mathbf{q}_0}^{-1,p}(U)}$ . Above, if  $\partial U$  is smooth or convex, we may take any  $p \in (1, \infty)$  with a constant  $C(p, \mathbf{s}_0^{-1/2} U, d) < \infty$  and if  $\partial U$  is merely Lipschitz, then we have the same conclusion with  $p \in [3/2, 3]$  and  $C(\mathbf{s}_0^{-1/2} U, d) < \infty$ , see [JK95, FMM98]. In the Lipschitz case we take  $p := \frac{3}{2} \vee 2_*$  with  $2_* := \frac{2d}{d+2}$ , and in the smooth or convex case with  $d > 2$  we take  $p = 2_*$  and if  $d = 2$ , any number larger than one. Applying the above estimate for  $w_r - u_{\text{hom}}$ , we deduce that

$$\begin{aligned} \|\mathbf{s}_0^{1/2} \nabla (w_r - u_{\text{hom}})\|_{\underline{L}^p(U)} &\leq C \|\zeta_r \mathbf{s}_0^{-1/2} (\mathbf{a}_0 \nabla u - \mathbf{a} \nabla u) * \eta_r\|_{\underline{L}^p(U)} + C \lambda_0^{-1/2} \|\nabla \zeta_r \cdot (\mathbf{a} \nabla u) * \eta_r\|_{\underline{W}_{\mathbf{q}_0}^{-1,p}(U)} \\ &\quad + C \|(u_r - g) \mathbf{s}_0^{1/2} \nabla \zeta_r\|_{\underline{L}^p(U)} + C \|(1 - \zeta_r) \mathbf{s}_0^{1/2} \nabla g\|_{\underline{L}^p(U)}. \end{aligned} \quad (6.42)$$

We obtain by the Sobolev inequality that

$$\begin{aligned} \|u_r - u_{\text{hom}}\|_{\underline{L}^2(U_{n+1}^\circ)} &\leq |U_{n+2}^\circ|^{-1/2} \|w_r - u_{\text{hom}}\|_{\underline{L}^2(\mathbb{R}^d)} \\ &\leq C(d) |U_{n+2}^\circ|^{-1/2} \|\mathbf{q}_0^{1/2} \nabla (w_r - u_{\text{hom}})\|_{L^{2*}(\mathbb{R}^d)} \\ &\leq C(d) \lambda_0^{-1/2} |U_{n+2}^\circ|^{-1/2} |U|^{1/2*} \|\mathbf{s}_0^{1/2} \nabla (w_r - u_{\text{hom}})\|_{\underline{L}^{2*}(U)} \\ &\leq C(\mathbf{q}_0^{-1/2} U, d) \lambda_0^{-1/2} 3^m \|\mathbf{s}_0^{1/2} \nabla (w_r - u_{\text{hom}})\|_{\underline{L}^p(U)}. \end{aligned} \quad (6.43)$$

Thus, by the triangle inequality, (6.20) and a covering argument, we also obtain that

$$\begin{aligned} &\frac{\lambda_0^{1/2}}{3^m} \|u - u_{\text{hom}}\|_{\underline{L}^2(U_{n+1}^\circ)} \\ &\leq \frac{\lambda_0^{1/2}}{3^m} \|u_r - u_{\text{hom}}\|_{\underline{L}^2(U_{n+1}^\circ)} + \frac{\lambda_0^{1/2}}{3^m} \|u_r - u\|_{\underline{L}^2(U_{n+1}^\circ)} \\ &\leq C \|\mathbf{s}_0^{1/2} \nabla (w_r - u_{\text{hom}})\|_{\underline{L}^p(U)} + C 3^{n-m} \left( 1 + \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n) \right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(U)}. \end{aligned} \quad (6.44)$$

In view of the last three displays, the goal is therefore to estimate the terms on the right in (6.42).

The first term on the right in (6.42) is the main one, which is estimated by Lemma 6.3. Indeed, this is where we use that  $u_r$  is nearly a solution of the equation for  $\mathbf{a}_0$ . By (6.19), we have that

$$\|\zeta_r \mathbf{s}_0^{-1/2} (\mathbf{a}_0 \nabla u - \mathbf{a} \nabla u) * \eta_r\|_{\underline{L}^2(U)} \leq C \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(U)}. \quad (6.45)$$

The other terms represent boundary layer errors. We will handle the last term first, since it is the simplest. We have that

$$\|(1 - \zeta_r) \mathbf{s}_0^{1/2} \nabla g\|_{\underline{L}^p(U)} \leq C |U|^{-1/p} \|\mathbf{s}_0^{1/2} \nabla g\|_{L^p(U \setminus U_{n+1}^\circ)} \leq C 3^{-(1/p-1/2)(m-n)} \|\mathbf{s}_0^{1/2} \nabla g\|_{\underline{L}^2(U)}. \quad (6.46)$$

Similarly,

$$\|(u_r - g) \mathbf{s}_0^{1/2} \nabla \zeta_r\|_{\underline{L}^p(U)} \leq C \lambda_0^{1/2} 3^{-(1/p-1/2)(m-n)} \frac{\|u * \eta_r - g\|_{L^2(U_n^\circ \setminus U_{n+1}^\circ)}}{3^n |U|^{1/2}}. \quad (6.47)$$

Next, by the triangle inequality, (6.19), (6.20) and a covering argument, we have that

$$\|\mathbf{s}_0^{-1/2}(\mathbf{a}\nabla u) * \eta_r\|_{\underline{L}^2(U_{n+1}^\circ)} \leq C \left(1 + \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n)\right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(U)}.$$

Letting  $\psi \in W_{0, \mathbf{q}_0}^{1, p'}(U)$  be such that  $\|\mathbf{q}_0^{1/2} \nabla \psi\|_{\underline{L}^{p'}(U)} \leq 1$ , we then deduce that

$$\begin{aligned} \lambda_0^{-1/2} \left| \int_U \nabla \zeta_r \cdot (\mathbf{a}\nabla u) * \eta_r \psi \right| &\leq C |U|^{-1} \|\mathbf{s}_0^{-1/2}(\mathbf{a}\nabla u) * \eta_r\|_{L^p(U_n^\circ \setminus U_{n+1}^\circ)} \|\mathbf{q}_0^{1/2} \nabla \zeta_r\| \|\psi\|_{L^{p'}(U_n^\circ \setminus U_{n+1}^\circ)} \\ &\leq C |U|^{-1/p} \|\mathbf{s}_0^{-1/2}(\mathbf{a}\nabla u) * \eta_r\|_{L^p(U_n^\circ \setminus U_{n+1}^\circ)} \|\mathbf{q}_0^{1/2} \nabla \psi\|_{\underline{L}^{p'}(U)} \\ &\leq C 3^{-(1/p-1/2)(m-n)} \left(1 + \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n)\right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(U)}. \end{aligned} \quad (6.48)$$

The second last inequality is true by the Poincaré inequality since  $\psi$  can be extended to be zero outside of  $U$  and then using the property that the complement of  $U$  has a positive geometric density. It follows that

$$\lambda_0^{-1/2} \|\nabla \zeta_r \cdot (\mathbf{a}\nabla u) * \eta_r\|_{\underline{W}^{-1, p}(U)} \leq C 3^{-(1/p-1/2)(m-n)} \left(1 + \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n)\right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(U)}.$$

Combining the above estimates with (6.42), (6.43) and (6.44) shows that

$$\begin{aligned} \|\mathbf{s}_0^{1/2} \nabla (w_r - u_{\text{hom}})\|_{\underline{L}^p(U)} &\leq C \left(3^{-(1/p-1/2)(m-n)} + \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n)\right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(U)} \\ &\quad + C 3^{-(1/p-1/2)(m-n)} \left( \|\mathbf{s}_0^{1/2} \nabla g\|_{\underline{L}^2(U)} + \lambda_0^{1/2} \frac{\|u * \eta_r - g\|_{L^2(U_n^\circ \setminus U_{n+1}^\circ)}}{3^n |U|^{1/2}} \right), \end{aligned} \quad (6.49)$$

and hence (6.44) implies (6.39).

To show (6.40), we observe that we have, by interior Calderón-Zygmund estimates, for  $\tilde{w}$  as before, that

$$\|\mathbf{q}_0^{1/2} \nabla \tilde{w}\|_{\underline{L}^p(y + \diamond_k)} \leq C 3^{-k} \|\tilde{w}\|_{\underline{L}^1(y + \diamond_{k+1})} + C \|\mathbf{q}_0^{-1/2} \tilde{\mathbf{f}}\|_{\underline{L}^p(y + \diamond_{k+1})} + C \|\tilde{g}\|_{\underline{W}_{\mathbf{q}_0}^{-1, p}(y + \diamond_{k+1})}$$

If  $y + \diamond_{k+1}$  is as in the statement, then the equation for  $\tilde{w} = u_r - u_{\text{hom}}$  satisfies the above estimate with  $\tilde{\mathbf{f}} := \eta_r * (\mathbf{a}\nabla u - \mathbf{a}_0 \nabla u)$ , and the result follows with the same proof as (6.34) had. The only extra ingredient is the local term  $\|\tilde{w}\|_{\underline{L}^1(y + \diamond_{k+1})}$ , but that can be estimated using (6.39). The proof is complete.  $\square$

An adaptation of the previous proof gives us (1.7) in Theorem A.

*Proof of (1.7) in Theorem A.* Let  $u$  and  $u_{\text{hom}}$  be as in (1.6). Take  $\mathbf{a}_0 = \bar{\mathbf{a}}$ . Observe that by testing the equation of  $u$  with  $u - g$ , using also Sobolev's inequality and Young's inequality, gives us

$$\begin{aligned} \|\bar{\mathbf{s}}^{1/2} \nabla u\|_{\underline{L}^2(U)} &\leq \left(\frac{\Lambda_0}{\lambda}\right)^{1/2} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(U)} \\ &\leq C \left(\frac{\bar{\Lambda}}{\lambda}\right)^{1/2} \|(\mathbf{s} + \mathbf{k}^t \mathbf{s}^{-1} \mathbf{k})^{1/2} \nabla g\|_{\underline{L}^2(U)} + C \left(\frac{\bar{\Lambda}}{\lambda}\right)^{1/2} \lambda^{-1/2} 3^m \|f\|_{\underline{L}^{2*}(U)} \\ &\leq C \frac{\Lambda}{\lambda} \left( \|\bar{\mathbf{s}}^{1/2} \nabla g\|_{\underline{L}^2(U)} + 3^m \|f\|_{\underline{L}^{2*}(U)} \right). \end{aligned} \quad (6.50)$$

This is a crude estimate we will use whenever there is geometric smallness provided by powers of  $3^{-(m-n)}$ .

We next adapt the notation from the proof of Proposition 6.8. Now (6.41) becomes

$$\begin{aligned} \nabla \cdot \bar{\mathbf{a}} \nabla (w_r - u_{\text{hom}}) &= \nabla \cdot (\zeta_r (\bar{\mathbf{a}} \nabla u - \mathbf{a} \nabla u) * \eta_r) + \nabla \zeta_r \cdot (\mathbf{a} \nabla u) * \eta_r + f - (f * \eta_r) \zeta_r \\ &\quad + \nabla \cdot ((u_r - g) \bar{\mathbf{s}} \nabla \zeta_r + (1 - \zeta_r) \bar{\mathbf{s}} \nabla g). \end{aligned} \quad (6.51)$$

From this we have an analogous estimate to (6.42) with  $p = 2_*$  for  $d > 2$  and any number larger than one for  $d = 2$ :

$$\begin{aligned} &\|\mathbf{s}_0^{1/2} \nabla (w_r - u_{\text{hom}})\|_{\underline{L}^p(U)} \\ &\leq C \|\zeta_r \bar{\mathbf{s}}^{-1/2} (\bar{\mathbf{a}} \nabla u - \mathbf{a} \nabla u) * \eta_r\|_{\underline{L}^p(U)} + C \bar{\lambda}^{-1/2} \|\nabla \zeta_r \cdot (\mathbf{a} \nabla u) * \eta_r\|_{\underline{W}_{\mathbf{q}_0}^{-1,p}(U)} \\ &\quad + \|f - (f * \eta_r) \zeta_r\|_{\underline{W}_{\bar{\mathbf{q}}}^{-1,p}(U)} + C \|(u_r - g) \bar{\mathbf{s}}^{1/2} \nabla \zeta_r\|_{\underline{L}^p(U)} + C \|(1 - \zeta_r) \bar{\mathbf{s}}^{1/2} \nabla g\|_{\underline{L}^p(U)}. \end{aligned} \quad (6.52)$$

The last two terms on the right can be treated similarly to (6.46) and (6.47) as

$$\|(1 - \zeta_r) \bar{\mathbf{s}}^{1/2} \nabla g\|_{\underline{L}^p(U)} \leq C 3^{-(1/p-1/2)(m-n)} \|\bar{\mathbf{s}}^{1/2} \nabla g\|_{\underline{L}^2(U)}$$

and

$$\|(u_r - g) \bar{\mathbf{s}}^{1/2} \nabla \zeta_r\|_{\underline{L}^p(U)} \leq C \bar{\lambda}^{1/2} 3^{-(1/p-1/2)(m-n)} \frac{\|u * \eta_r - g\|_{L^2(U_n^\circ \setminus U_{n+1}^\circ)}}{3^n |U|^{1/2}},$$

and further, by Poincaré's inequality and (6.50),

$$\begin{aligned} \bar{\lambda}^{1/2} \frac{\|u * \eta_r - g\|_{L^2(U_n^\circ \setminus U_{n+1}^\circ)}}{3^n |U|^{1/2}} &\leq C \|\bar{\mathbf{s}}^{1/2} \nabla (u - g)\|_{\underline{L}^2(U)} \\ &\leq C \frac{\Lambda}{\lambda} \left( \|\mathbf{s}_0^{1/2} \nabla g\|_{\underline{L}^2(U)} + 3^m \|f\|_{\underline{L}^{2*}(U)} \right). \end{aligned} \quad (6.53)$$

Taking  $\psi \in W_{\bar{\mathbf{q}}}^{1,p'}$  such that  $\|\bar{\mathbf{q}} \nabla \psi\|_{\underline{L}^{p'}(U)} \leq 1$  and by extending it as zero outside of  $U$ , we obtain, by Poincaré's inequality,

$$\begin{aligned} \int_U \psi (f - (f * \eta_r) \zeta_r) &= \int_U f (\psi - \psi * \eta_r) + \int_U f \eta_r * (\psi (1 - \zeta_r)) \\ &\leq C 3^n \|f\|_{\underline{L}^p(U)} \|\bar{\mathbf{q}} \nabla \psi\|_{\underline{L}^{p'}(U)} + C |U|^{1/p'} \|f\|_{\underline{L}^p(U)} \|\psi (1 - \zeta_r)\|_{\underline{L}^{p'}(\mathbb{R}^d)} \\ &\leq C 3^n \|f\|_{\underline{L}^p(U)}. \end{aligned}$$

Combining the last three displays yields

$$\begin{aligned} &\|f - (f * \eta_r) \zeta_r\|_{\underline{W}_{\bar{\mathbf{q}}}^{-1,p}(U)} + C \|(u_r - g) \bar{\mathbf{s}}^{1/2} \nabla \zeta_r\|_{\underline{L}^p(U)} + C \|(1 - \zeta_r) \bar{\mathbf{s}}^{1/2} \nabla g\|_{\underline{L}^p(U)} \\ &\leq C \frac{\Lambda}{\lambda} 3^{-(1/p-1/2)(m-n)} \left( \|\bar{\mathbf{s}}^{1/2} \nabla g\|_{\underline{L}^2(U)} + 3^m \|f\|_{\underline{L}^{2*}(U)} \right). \end{aligned} \quad (6.54)$$

We next seek the replacements for (6.45) and (6.48) in the previous proof, but now in the uniformly elliptic setting. We use a straightforward localization using harmonic approximation. To

this end, let  $z \in 3^{n+1}\mathbb{L}_0 \cap U$  be such that  $z + \diamond_{n+1} \subset U$ . Let  $u_z$  be the unique solution  $u_z \in \mathcal{A}(z + \diamond_{n+1}) \cap (u + H_0^1(z + \diamond_{n+1}))$ . Testing then gives

$$\begin{aligned} \|\mathbf{s}^{1/2} \nabla(u - u_z)\|_{\underline{L}^2(z + \diamond_{n+1})}^2 &\leq \|u - u_z\|_{\underline{L}^{2*}(z + \diamond_{n+1})} \|f\|_{\underline{L}^{2*}(z + \diamond_{n+1})} \\ &\leq C 3^n \|\nabla(u - u_z)\|_{\underline{L}^2(z + \diamond_{n+1})} \|f\|_{\underline{L}^{2*}(z + \diamond_{n+1})} \\ &\leq C \lambda^{-1/2} 3^n \|\mathbf{s}^{1/2} \nabla(u - u_z)\|_{\underline{L}^2(z + \diamond_{n+1})} \|f\|_{\underline{L}^{2*}(z + \diamond_{n+1})}, \end{aligned}$$

which yields

$$\|\mathbf{s}^{1/2} \nabla(u - u_z)\|_{\underline{L}^2(z + \diamond_{n+1})} \leq C \lambda^{-1/2} 3^n \|f\|_{\underline{L}^{2*}(z + \diamond_{n+1})}. \quad (6.55)$$

By (6.19), the above display and the triangle inequality, we obtain

$$\begin{aligned} &\|\eta_r * (\bar{\mathbf{s}}^{-1/2}(\mathbf{a} - \bar{\mathbf{s}}) \nabla u)\|_{L^\infty(z + \diamond_n)} \\ &\leq \|\eta_r * (\bar{\mathbf{s}}^{-1/2}(\mathbf{a} - \bar{\mathbf{s}}) \nabla u_z)\|_{L^\infty(z + \diamond_n)} + \|\eta_r * (\bar{\mathbf{s}}^{-1/2}(\mathbf{a} - \bar{\mathbf{s}}) \nabla(u - u_z))\|_{L^\infty(z + \diamond_n)} \\ &\leq C \mathcal{E}_1(z + \diamond_n; \mathbf{a}, \bar{\mathbf{a}}) \|\mathbf{s}^{1/2} \nabla u_z\|_{\underline{L}^2(z + \diamond_{n+1})} + C \left(\frac{\Lambda}{\lambda}\right)^{1/2} \|\mathbf{s}^{1/2} \nabla(u - u_z)\|_{\underline{L}^2(z + \diamond_{n+1})} \\ &\leq C \mathcal{E}_1(z + \diamond_n; \mathbf{a}, \bar{\mathbf{a}}) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(z + \diamond_{n+1})} + C \frac{\Lambda}{\lambda} 3^n \|f\|_{\underline{L}^{2*}(z + \diamond_{n+1})}. \end{aligned} \quad (6.56)$$

Similarly, using also (6.20), we get

$$\|\eta_r * (\bar{\mathbf{s}}^{1/2} \nabla u)\|_{L^\infty(z + \diamond_n)} \leq C \frac{\Lambda}{\lambda} \left( \|\bar{\mathbf{s}}^{1/2} \nabla g\|_{\underline{L}^2(U)} + 3^m \|f\|_{\underline{L}^{2*}(U)} \right). \quad (6.57)$$

These can be used instead of (6.45) and (6.48) in the previous proof. We obtain

$$\begin{aligned} &\|\eta_r * (\mathbf{s}_0^{-1/2}(\mathbf{a} - \mathbf{s}_0) \nabla u) \zeta_r\|_{\underline{L}^2(U)} \\ &\leq C \frac{\Lambda}{\lambda} \left( \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n; \mathbf{a}, \bar{\mathbf{a}}) + 3^{-(m-n)} \right) \left( \|\bar{\mathbf{s}}^{1/2} \nabla g\|_{\underline{L}^2(U)} + 3^m \|f\|_{\underline{L}^{2*}(U)} \right). \end{aligned}$$

and

$$\bar{\lambda}^{-1/2} \left| \int_U \nabla \zeta_r \cdot (\mathbf{a} \nabla u) * \eta_r \psi \right| \leq C \frac{\Lambda}{\lambda} 3^{-(1/p+1/2)(m-n)} \left( \|\bar{\mathbf{s}}^{1/2} \nabla g\|_{\underline{L}^2(U)} + 3^m \|f\|_{\underline{L}^{2*}(U)} \right).$$

Combining the above two displays with (6.52) and (6.54) implies that

$$\begin{aligned} &\|\mathbf{s}_0^{1/2} \nabla(w_r - u_{\text{hom}})\|_{\underline{L}^p(U)} \\ &\leq C \frac{\Lambda}{\lambda} \left( \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n; \mathbf{a}, \bar{\mathbf{a}}) + 3^{-(1/p+1/2)(m-n)} \right) \left( \|\bar{\mathbf{s}}^{1/2} \nabla g\|_{\underline{L}^2(U)} + 3^m \|f\|_{\underline{L}^{2*}(U)} \right). \end{aligned}$$

Similarly to (6.44), using (6.55), we then obtain

$$\begin{aligned} &\frac{\bar{\lambda}^{1/2}}{3^m} \|u - u_{\text{hom}}\|_{\underline{L}^2(U)} \\ &\leq C \frac{\Lambda}{\lambda} \left( \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(y + \diamond_n; \mathbf{a}, \bar{\mathbf{a}}) + 3^{-(1/p+1/2)(m-n)} \right) \left( \|\bar{\mathbf{s}}^{1/2} \nabla g\|_{\underline{L}^2(U)} + 3^m \|f\|_{\underline{L}^{2*}(U)} \right). \end{aligned}$$

By choosing  $n$  appropriately, an application of Corollary 4.5 with smaller  $\delta$  depending on  $\Lambda/\lambda$  completes the proof of (1.7).  $\square$

The statement of Proposition 6.7 allows us to approximate an arbitrary element of  $\mathcal{A}(U)$  by a solution of the homogenized equation. It does not imply the converse statement—that we can approximate an arbitrary solution of the homogenized equation in  $U$  by an element of  $\mathcal{A}(U)$ . This is particular to our very general setting. Indeed, typically in homogenization theory, the approximation of solutions of the homogenized equation by those of the heterogeneous equation is the “easy” direction, since the homogenized solution is smoother. In our setting, the heterogeneous equation can be very degenerate, so an additional step is required, and we must construct our approximations by brute force. This is the point of the following lemma.

In the proof we need to approximate the given  $\mathbf{a}_0$ -harmonic function by piecewise linear functions, and for that we need to use adapted simplexes.

**Proposition 6.9** (Harmonic approximation II). *Let  $s \in (0, 1]$  and  $m, n \in \mathbb{N}$  with  $n < m - 2$ . For  $\mathcal{E}_s^\Delta$  defined by (6.4), define a composite  $\tilde{\mathcal{E}}_s(\diamond_m; \mathbf{a}, \mathbf{a}_0)$  quantity by*

$$\hat{\mathcal{E}}_{s,n}(\diamond_m; \mathbf{a}, \mathbf{a}_0) := \max_{\Delta \in \mathcal{Z}_n^\Delta(\diamond_m)} \left( \mathcal{E}_1^\Delta(\Delta; \mathbf{a}, \mathbf{a}_0) + 3^{-s(m-n)} (1 + \mathcal{E}_{s,n}^\Delta(\Delta; \mathbf{a}, \mathbf{a}_0)) \right). \quad (6.58)$$

*There exists a constant  $C(d) < \infty$  such that for every  $\mathbf{a}_0$ -harmonic function  $u_{\text{hom}} \in W^{1,\infty}(\diamond_m)$  in  $\diamond_m$  there exists  $\mathcal{F}(\diamond_m)$ -measurable  $u \in \mathcal{A}(\diamond_m)$  such that*

$$\begin{aligned} 3^{-m} \lambda_0^{1/2} \|u - u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)} + s 3^{-sm} \left[ \mathbf{A}_0^{1/2} \left( \frac{\nabla u - \nabla u_{\text{hom}}}{\mathbf{a} \nabla u - \mathbf{a}_0 \nabla u_{\text{hom}}} \right) \right]_{\underline{B}_{2,1}^{-s}(\diamond_m)} \\ \leq C \hat{\mathcal{E}}_{s,n}(\diamond_m; \mathbf{a}, \mathbf{a}_0) \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)} + C 3^n \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{L^2(\diamond_m)} \end{aligned} \quad (6.59)$$

Moreover,

$$\begin{aligned} \left| \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}^2 - \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}^2 \right| \\ \leq C (\hat{\mathcal{E}}_{1,n}(\diamond_m; \mathbf{a}, \mathbf{a}_0))^2 \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}^2 + C 3^{2n} \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{L^\infty(\diamond_m)}^2. \end{aligned} \quad (6.60)$$

*Proof.* Fix  $m, n \in \mathbb{N}$  with  $n \leq m - 2$ . We divide the proof in multiple steps. For reasons which will become apparent below, we work here with simplexes rather than cubes for our partitions.

First, if  $\hat{\mathcal{E}}_1 := \hat{\mathcal{E}}_{1,n}(\diamond_m; \mathbf{a}, \mathbf{a}_0) > 1$ , then we simply take  $u = (u_{\text{hom}})_{\diamond_m}$  and use the straightforward estimate

$$3^{-m} \lambda_0^{1/2} \|u_{\text{hom}} - (u_{\text{hom}})_{\diamond_m}\|_{\underline{L}^2(\diamond_m)} + s 3^{-sm} \left[ \mathbf{A}_0^{1/2} \left( \frac{\nabla u_{\text{hom}}}{\mathbf{a}_0 \nabla u_{\text{hom}}} \right) \right]_{\underline{B}_{2,1}^{-s}(\diamond_m)} \leq C \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}.$$

Also (6.60) is trivial. Thus, for the rest of the argument, we assume that  $\hat{\mathcal{E}}_1 \leq 1$ .

In Step 1 we show the needed Besov estimates for the gradients and fluxes in adapted simplexes. In Step 2 we construct the candidate for the heterogeneous approximation of  $u_{\text{hom}}$ , and in Steps 3-7 we show that it has the desired properties. Throughout the proof, we assume without loss of generality that  $\mathbf{a}_0 = \mathbf{s}_0$  so that

$$\mathbf{A}_0 = \begin{pmatrix} \mathbf{s}_0 & 0 \\ 0 & \mathbf{s}_0^{-1} \end{pmatrix}.$$

*Step 1.* Let  $n, k \in \mathbb{N}$  with  $k \leq n$ . For  $s \in (0, 1]$  and  $\Delta \in S_n$ , we set

$$[f]_{\underline{B}_{2,1}^{-s}(\Delta)} := \sum_{k=-\infty}^n 3^{sk} \left( \sum_{\Delta' \in \mathcal{Z}_k(\Delta)} |(f)_{\Delta'}|^2 \right)^{1/2}.$$



Let  $X \in \mathcal{S}(\Delta)$ . By (2.40) and the identities  $\mathbf{A}_*^{-1}(\Delta) = \mathbf{R}\mathbf{A}(\Delta)\mathbf{R}$  and  $\mathbf{A}_0^{-1/2} = \mathbf{R}\mathbf{A}_0^{1/2}\mathbf{R}$ , with  $\mathbf{R}$  as in (2.32), we get

$$|(\mathbf{A}_0^{1/2}X)_{\Delta'}|^2 \leq |\mathbf{A}_*^{-1/2}(\Delta')\mathbf{A}_0\mathbf{A}_*^{-1/2}(\Delta')| |(\mathbf{A}_*^{1/2}(\Delta')X)_{\Delta'}|^2 \leq |\mathbf{A}_0^{-1/2}\mathbf{A}(\Delta')\mathbf{A}_0^{-1/2}| \|\mathbf{A}^{1/2}X\|_{\underline{L}^2(\Delta')}^2. \quad (6.61)$$

Thus we obtain

$$\begin{aligned} s3^{-sn}[\overline{\mathbf{A}}^{1/2}X]_{\underline{B}_{2,1}^{-s}(\Delta)} &\leq s3^d \|\mathbf{A}^{1/2}X\|_{\underline{L}^2(\Delta)} \sum_{k=-\infty}^n 3^{s(k-n)} \max_{\Delta' \in \mathcal{Z}_k(\Delta)} |\mathbf{A}_0^{-1/2}\mathbf{A}(\Delta')\mathbf{A}_0^{-1/2}|^{1/2} \\ &\leq C \left(1 + \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_s^\Delta(\Delta)\right) \|\mathbf{A}^{1/2}X\|_{\underline{L}^2(\Delta)}. \end{aligned} \quad (6.62)$$

The last inequality is true since we have

$$\begin{aligned} |\mathbf{A}_0^{-1/2}\mathbf{A}(U)\mathbf{A}_0^{-1/2}| &\leq \sup_{|e|^2 \leq 2} \begin{pmatrix} -\mathbf{s}_0^{-1/2}e \\ \mathbf{s}_0^{1/2}e \end{pmatrix} \cdot \mathbf{A}_{\mathbf{k}_0}(U) \begin{pmatrix} -\mathbf{s}_0^{-1/2}e \\ \mathbf{s}_0^{1/2}e \end{pmatrix} + \sup_{|e|^2 \leq 2} \begin{pmatrix} \mathbf{s}_0^{-1/2}e \\ \mathbf{s}_0^{1/2}e \end{pmatrix} \cdot \mathbf{A}_{\mathbf{k}_0}(U) \begin{pmatrix} \mathbf{s}_0^{-1/2}e \\ \mathbf{s}_0^{1/2}e \end{pmatrix} \\ &= 4 \left(1 + \sup_{|e| \leq 1} J(U, \mathbf{s}_0^{-1/2}e, \mathbf{a}_0^t \mathbf{s}_0^{-1/2}e) + \sup_{|e| \leq 1} J^*(U, \mathbf{s}_0^{-1/2}e, \mathbf{a}_0 \mathbf{s}_0^{-1/2}e)\right), \end{aligned} \quad (6.63)$$

and it follows that

$$s \sum_{k=-\infty}^n 3^{s(k-n)} \max_{\Delta' \in \mathcal{Z}_k(\Delta)} |\mathbf{A}_0^{-1/2}\mathbf{A}(\Delta')\mathbf{A}_0^{-1/2}|^{1/2} \leq C(1 + \mathcal{E}_s^\Delta(\Delta)). \quad (6.64)$$

*Step 2.* Construction of heterogeneous solution. Given a solution  $u_{\text{hom}}$  of the homogenized equation, we approximate  $u_{\text{hom}}$  in  $\diamond_n$  by a piecewise affine function, denoted by  $v$ , which is uniquely defined by the conditions:

- $v|_{\Delta}$  is affine for each  $\Delta \in \mathcal{S}_n$ ; and
- $v(x) = u_{\text{hom}}(x)$  for every vertex  $x$  of the simplexes in the partition of  $\diamond_m$ .

It follows that, for every  $\Delta$  in the partition,

$$\|\mathbf{s}_0^{1/2}\nabla(v - u_{\text{hom}})\|_{\underline{L}^2(\Delta)} \leq C3^n \lambda_0^{-1/2} \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{\underline{L}^2(\Delta)}. \quad (6.65)$$

We let  $W_\Delta$  being the maximizer obtained by solving the optimization problem for  $\mathbf{J}(\Delta, P_\Delta, 0)$  in each simplex  $\Delta$  in the partition, where

$$p_\Delta := \nabla v|_{\Delta} \quad \text{and} \quad P_\Delta := - \begin{pmatrix} p_\Delta \\ \mathbf{s}_0 p_\Delta \end{pmatrix}.$$

In particular, since  $W_\Delta$  is the negation of the minimizer of the energy over  $P_\Delta + L_{\mathbf{a}, \text{pot}, 0}^2(\Delta) \times L_{\mathbf{a}, \text{sol}, 0}^2(\Delta)$ , we have, using also the first variation, that

$$\int_{\Delta} W_\Delta = -P_\Delta \quad \text{and} \quad Y \in \mathcal{S}(\Delta) \implies \int_{\Delta} W_\Delta \cdot \mathbf{A}Y = -P_\Delta \cdot (\mathbf{A}Y)_\Delta. \quad (6.66)$$

Let  $W$  be the function glued together:  $W(x) = W_\Delta(x)$  whenever  $x \in \Delta$ . By (2.15), we have that

$$\frac{1}{2} \int_{\Delta} W_\Delta \cdot \mathbf{A}W_\Delta = \mathbf{J}(\Delta, P_\Delta, 0) = \frac{1}{2} J(\Delta, -p_\Delta, \mathbf{s}_0 p_\Delta) + \frac{1}{2} J^*(\Delta, p_\Delta, \mathbf{s}_0 p_\Delta). \quad (6.67)$$

Now, by (2.13), we have

$$J^*(\Delta, p_\Delta, \mathbf{s}_0 p_\Delta) \leq C \left( \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1^\Delta(y + \diamond_n) \right)^2 |\mathbf{s}_0^{1/2} p_\Delta|^2 \quad (6.68)$$

and

$$J(\Delta, -p_\Delta, \mathbf{s}_0 p_\Delta) = 2 |\mathbf{s}_0^{1/2} p_\Delta|^2 + J^*(\Delta, p_\Delta, \mathbf{s}_0 p_\Delta). \quad (6.69)$$

Therefore,

$$\begin{aligned} & \left| \frac{1}{2} \|\mathbf{A}^{1/2} W\|_{\underline{L}^2(\diamond_m)}^2 - \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}^2 \right| \\ & \leq C \left( \max_{y \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1^\Delta(y + \diamond_n) \right)^2 \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}^2 + C 3^{2n} \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}^2. \end{aligned} \quad (6.70)$$

We will show that  $W$  is close to the minimizer of a quadratic minimization problem in  $\diamond_m$ . Let  $X \in \mathcal{S}(\diamond_m)$  be the minimizer of

$$\inf_{X \in \mathcal{S}(\diamond_m)} \int_{\diamond_m} \frac{1}{2} (X - W) \cdot \mathbf{A} (X - W)$$

The above minimization problem is also given over all functions with Dirichlet data given by  $W$  and, in particular,  $X - W \in L_{\mathbf{a}, \text{pot}, 0}^2(\diamond_m) \times L_{\mathbf{a}, \text{sol}, 0}^2(\diamond_m)$  and

$$\int_{\diamond_m} \frac{1}{2} (X - W) \cdot \mathbf{A} (X - W) = \int_{\diamond_m} \frac{1}{2} W \cdot \mathbf{A} W - \int_{\diamond_m} \frac{1}{2} X \cdot \mathbf{A} X. \quad (6.71)$$

Furthermore, since  $X$  and  $W$  are members of  $\mathcal{S}(\diamond_m)$  and  $\mathcal{S}(\Delta)$  in each  $\Delta$ , respectively, we find, for each  $\Delta$ , elements  $(u, u^*) \in \mathcal{A}(\diamond_m) \times \mathcal{A}^*(\diamond_m)$  and  $(w, w^*) \in \mathcal{A}(\Delta) \times \mathcal{A}^*(\Delta)$  such that

$$X = \begin{pmatrix} \nabla u + \nabla u^* \\ \mathbf{a} \nabla u - \mathbf{a}^t \nabla u^* \end{pmatrix} \quad \text{and} \quad W = \begin{pmatrix} \nabla w + \nabla w^* \\ \mathbf{a} \nabla w - \mathbf{a}^t \nabla w^* \end{pmatrix}. \quad (6.72)$$

Moreover,  $(w, w^*)$  are the maximizers of  $J$  and  $J^*$  in (6.67) in each  $\Delta$ , respectively, and since  $W_\Delta$ s agree on the boundaries of the simplexes, there are globally defined potentials such that (6.72) is valid in the whole  $\diamond_m$ . Notice that, by (6.68), the energy of  $w^*$  is small. By a direct computation,

$$\frac{1}{2} \|\mathbf{A}^{1/2} (X - W)\|_{\underline{L}^2(\diamond_m)}^2 = \int_{\diamond_m} (\nabla(u - w) \cdot \mathbf{s} \nabla(u - w) + \nabla(u^* - w^*) \cdot \mathbf{s} \nabla(u^* - w^*)),$$

$$\frac{1}{2} \|\mathbf{A}^{1/2} W\|_{\underline{L}^2(\diamond_m)}^2 = \|\mathbf{s}^{1/2} \nabla w\|_{\underline{L}^2(\diamond_m)}^2 + \|\mathbf{s}^{1/2} \nabla w^*\|_{\underline{L}^2(\diamond_m)}^2$$

and

$$\frac{1}{2} \|\mathbf{A}^{1/2} X\|_{\underline{L}^2(\diamond_m)}^2 = \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}^2 + \|\mathbf{s}^{1/2} \nabla u^*\|_{\underline{L}^2(\diamond_m)}^2.$$

Therefore, by (6.68), (6.70), (6.71), the previous two displays and the triangle inequality, we obtain

$$\begin{aligned} & \left| \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}^2 - \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}^2 \right| \\ & \leq 2 \|\mathbf{A}^{1/2} (X - W)\|_{\underline{L}^2(\diamond_m)}^2 + C \tilde{\mathcal{E}}_1^2 \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}^2 + C 3^{2n} \lambda_0^{-1/2} \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}^2. \end{aligned} \quad (6.73)$$

We will show in Step 8 below that there exists a constant  $C(d) < \infty$  such that

$$\|\mathbf{A}^{1/2}(X - W)\|_{\underline{L}^2(\diamond_m)} \leq C\tilde{\mathcal{E}}_1 \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)} + C3^n \lambda_0^{-1/2} \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}. \quad (6.74)$$

Before proving the above estimate, we show in the next step that this implies the result of the lemma.

*Step 3.* We show (6.59) and (6.60) assuming (6.74). First, (6.60) follows by (6.73) and (6.74). Second, by (6.70) and (6.74), recalling that  $\tilde{\mathcal{E}}_1 \leq 1$ , we obtain

$$\|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)} \leq \frac{1}{2} \|\mathbf{A}^{1/2} X\|_{\underline{L}^2(\diamond_m)} \leq C \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)} + C3^n \lambda_0^{-1/2} \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}.$$

Third, by the triangle inequality and Jensen's inequality we have, for  $\mathbf{F} \in B_{2,1}^{-s}(\diamond_m)$ , that

$$s3^{-sm} [\mathbf{F}]_{\dot{B}_{2,1}^{-s}(\diamond_m)} \leq C \left( \sum_{\Delta \in \mathcal{Z}_n(\diamond_m)} |(\mathbf{F})_\Delta|^2 \right)^{1/2} + 3^{-s(m-n)} \sum_{\Delta \in \mathcal{Z}_n(\diamond_m)} s3^{-sn} [\mathbf{F}]_{\dot{B}_{2,1}^{-s}(\Delta)}. \quad (6.75)$$

Having the above display in mind, we use the triangle inequality, (6.62) and (6.70) to get

$$\begin{aligned} s3^{-sm} \sum_{\Delta \in \mathcal{Z}_n(\diamond_m)} \left[ \mathbf{A}_0^{1/2} \left( \nabla u - \nabla u_{\text{hom}} \right) \right]_{\dot{B}_{2,1}^{-s}(\Delta)} \\ \leq C3^{-s(m-n)} \sum_{\Delta \in \mathcal{Z}_n(\diamond_m)} \left( \left( 1 + \max_{\Delta \in \mathcal{Z}_n(\diamond_m)} \mathcal{E}_s^\Delta(\Delta) \right) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)} + \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)} \right) \\ \leq C\tilde{\mathcal{E}}_s \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)} + C3^n \lambda_0^{-1/2} \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}. \end{aligned} \quad (6.76)$$

Corresponding the first term on the right in (6.75), we decompose the field, in each  $\Delta$ , as

$$\begin{pmatrix} \nabla u - \nabla u_{\text{hom}} \\ \mathbf{a} \nabla u - \mathbf{a}_0 \nabla u_{\text{hom}} \end{pmatrix} = \begin{pmatrix} \nabla u - \nabla w_\Delta \\ \mathbf{a} \nabla u - \mathbf{a} \nabla w_\Delta \end{pmatrix} + \begin{pmatrix} \nabla w_\Delta - p_\Delta \\ \mathbf{a} \nabla w_\Delta - \mathbf{s}_0 p_\Delta \end{pmatrix} + \begin{pmatrix} p_\Delta - \nabla u_{\text{hom}} \\ \mathbf{s}_0 (p_\Delta - \nabla u_{\text{hom}}) \end{pmatrix}. \quad (6.77)$$

The last term in (6.77) can be estimated using (6.65) as

$$\sum_{\Delta \in \mathcal{Z}_n(\diamond_m)} \|\mathbf{s}_0^{1/2} (\nabla u_{\text{hom}} - p_\Delta)_\Delta\|_{\underline{L}^2(\Delta)} \leq C3^n \lambda_0^{-1/2} \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}. \quad (6.78)$$

By (2.40) and (6.64) (applied with  $s = 1$ ) we get

$$\left( \sum_{\Delta \in \mathcal{Z}_n(\diamond_m)} \left| \int_\Delta \mathbf{A}_0^{1/2} \begin{pmatrix} \nabla u - \nabla w_\Delta \\ \mathbf{a} \nabla u - \mathbf{a} \nabla w_\Delta \end{pmatrix} \right|^2 \right)^{1/2} \leq C \|\mathbf{A}^{1/2}(X - W)\|_{\underline{L}^2(\diamond_m)}. \quad (6.79)$$

Furthermore, we have

$$\mathbf{A}_0^{1/2} \int_\Delta \begin{pmatrix} \nabla w_\Delta + \nabla w_\Delta^* \\ \mathbf{a} \nabla w_\Delta - \mathbf{a}^t \nabla w_\Delta^* \end{pmatrix} = \mathbf{A}_0^{1/2} \int_\Delta W_\Delta = -\mathbf{A}_0^{1/2} P_\Delta = \begin{pmatrix} \mathbf{s}_0^{1/2} p_\Delta \\ \mathbf{s}_0^{1/2} p_\Delta \end{pmatrix}.$$

Now (2.40), (6.64) and (6.68) yield that

$$\left| \mathbf{A}_0^{1/2} \int_\Delta \begin{pmatrix} \nabla w_\Delta^* \\ -\mathbf{a}^t \nabla w_\Delta^* \end{pmatrix} \right|^2 \leq C \int_\Delta \nabla w^* \cdot \mathbf{s} \nabla w^* \leq C J^*(\Delta, p_\Delta, \mathbf{s}_0 p_\Delta) \leq C \tilde{\mathcal{E}}_1^2 |\mathbf{s}_0^{1/2} p_\Delta|^2.$$

Therefore, by the triangle inequality and  $\tilde{\mathcal{E}}_1 \leq 1$ ,

$$\begin{aligned} & \left( \sum_{\Delta \in \mathcal{Z}_n(\diamond_m)} \left| \mathbf{A}_0^{1/2} \oint_{\Delta} \left( \mathbf{a} \nabla w_{\Delta} - p_{\Delta} \right) \right|^2 \right)^{1/2} \\ & \leq C \tilde{\mathcal{E}}_1 \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)} + C 3^n \lambda_0^{-1/2} \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)}. \end{aligned} \quad (6.80)$$

Combining now (6.77) with (6.78), (6.79), (6.80) and (6.74), and then inserting the result and (6.76) into (6.75), using also (2.128) by choosing the additive constant in  $u$  to be such that  $(u - u_{\text{hom}})_{\diamond_m} = 0$ , leads to (6.59).

*Step 4.* We begin the proof of (6.74) by estimating  $\mathbf{A}^{1/2}(X - W)$  in  $L^2$ . We claim that

$$\begin{aligned} & \oint_{\diamond_m} (X - W) \cdot \mathbf{A}(X - W) \\ & = \sum_{\Delta} \oint_{\Delta} \mathbf{A}_0 \left( P_{\Delta} + \left( \frac{\nabla u_{\text{hom}}}{\mathbf{s}_0 \nabla u_{\text{hom}}} \right) \right) \cdot (X - W) + \sum_{\Delta} P_{\Delta} \cdot ((\mathbf{A} - \mathbf{A}_0)(X - W))_{\Delta}. \end{aligned} \quad (6.81)$$

Since  $X - W \in L_{\mathbf{a}, \text{pot}, 0}^2(\diamond_m) \times L_{\mathbf{a}, \text{sol}, 0}^2(\diamond_m)$  and  $X \in \mathcal{S}(\diamond_m)$ , we obtain using (6.66) that

$$\begin{aligned} & \oint_{\diamond_m} (X - W) \cdot \mathbf{A}(X - W) = - \oint_{\diamond_m} W \cdot \mathbf{A}(X - W) \\ & = \sum_{\Delta} \mathbf{A}_0 P_{\Delta} \cdot (X - W)_{\Delta} + \sum_{\Delta} P_{\Delta} \cdot ((\mathbf{A} - \mathbf{A}_0)(X - W))_{\Delta}. \end{aligned} \quad (6.82)$$

Since  $X - W \in L_{\mathbf{a}, \text{pot}, 0}^2(\diamond_m) \times L_{\mathbf{a}, \text{sol}, 0}^2(\diamond_m)$  and  $-\nabla \cdot \mathbf{s}_0 \nabla u_{\text{hom}} = 0$ , we also deduce that

$$\oint_{\diamond_m} \mathbf{A}_0 \left( \frac{\nabla u_{\text{hom}}}{\mathbf{s}_0 \nabla u_{\text{hom}}} \right) \cdot (X - W) = \oint_{\diamond_m} \left( \frac{\mathbf{s}_0 \nabla u_{\text{hom}}}{\nabla u_{\text{hom}}} \right) \cdot (X - W) = 0.$$

Here we need also the fact that  $u_{\text{hom}}$  is in  $W^{1, \infty}(\diamond_m)$  since we only have  $X - W \in L^1(\diamond_m)$ . Therefore,

$$\sum_{\Delta} \mathbf{A}_0 P_{\Delta} \cdot (X - W)_{\Delta} = \sum_{\Delta} \oint_{\Delta} \mathbf{A}_0 \left( P_{\Delta} + \left( \frac{\nabla u_{\text{hom}}}{\mathbf{s}_0 \nabla u_{\text{hom}}} \right) \right) (X - W).$$

Combining this with (6.82) gives us (6.81).

*Step 5.* We show that

$$\begin{aligned} & |P_{\Delta} \cdot ((\mathbf{A} - \mathbf{A}_0)(X - W))_{\Delta}| \\ & \leq 4 \sup_{|e|=1} \left( J(\Delta, \mathbf{s}_0^{-1/2} e, \mathbf{s}_0^{1/2} e) + J^*(\Delta, \mathbf{s}_0^{-1/2} e, \mathbf{s}_0^{1/2} e) \right)^{1/2} |\mathbf{s}_0^{1/2} p_{\Delta}| \|\mathbf{A}^{1/2}(X - W)\|_{\underline{L}^2(\Delta)}. \end{aligned} \quad (6.83)$$

By the definition of  $P_{\Delta}$ , we have that

$$|\mathbf{A}_0^{1/2} P_{\Delta}|^2 = 2 |\mathbf{s}_0^{1/2} p_{\Delta}|^2.$$

We have

$$\begin{aligned} & \mathbf{A}_0^{-1/2} (\mathbf{A} - \mathbf{A}_0)(X - W) \\ & = \left( \mathbf{s}_0^{-1/2} (\mathbf{a} \nabla(u - w) - \mathbf{s}_0 \nabla(u - w)) \right) + \left( \mathbf{s}_0^{-1/2} (\mathbf{a}^t \nabla(u^* - w^*) - \mathbf{s}_0 \nabla(u^* - w^*)) \right) \\ & \quad - \left( \mathbf{s}_0^{-1/2} (\mathbf{s}_0 \nabla(u - w) - \mathbf{a} \nabla(u - w)) \right) + \left( \mathbf{s}_0^{-1/2} (-\mathbf{s}_0 \nabla(u^* - w^*) + \mathbf{a}^t \nabla(u^* - w^*)) \right). \end{aligned}$$

By (2.34), we obtain, for every  $e \in \mathbb{R}^d$

$$|\mathbf{s}_0^{-1/2}(\mathbf{a}\nabla(u-w) - \mathbf{s}_0\nabla(u-w))_\Delta|^2 \leq 2 \sup_{|e|=1} J(\Delta, \mathbf{s}_0^{-1/2}e, \mathbf{s}_0^{1/2}e) \oint_\Delta \nabla(u-w) \cdot \mathbf{s}\nabla(u-w).$$

A similar estimates is valid for  $u^* - w^*$ , and hence we obtain (6.83).

*Step 6.* We show that

$$\begin{aligned} & \left| \oint_\Delta \mathbf{A}_0 \left( P_\Delta + \begin{pmatrix} \nabla u_{\text{hom}} \\ \mathbf{s}_0 \nabla u_{\text{hom}} \end{pmatrix} \right) \cdot (X - W) \right| \\ & \leq C 3^n \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{\underline{L}^2(\Delta)} \|\mathbf{A}^{1/2}(X - W)\|_{\underline{L}^2(\Delta)} \sum_{k=-\infty}^n 3^{k-n} \max_{\Delta' \in \mathcal{Z}_k(\Delta)} |\mathbf{A}_0^{-1/2} \mathbf{A}(\Delta') \mathbf{A}_0^{-1/2}|^{1/2}. \end{aligned} \quad (6.84)$$

By (6.65) we have that

$$\left\| \mathbf{A}_0^{1/2} \left( P_\Delta + \begin{pmatrix} \nabla u_{\text{hom}} \\ \mathbf{s}_0 \nabla u_{\text{hom}} \end{pmatrix} \right) \right\|_{\underline{B}_{2,\infty}^1(\Delta)} \leq C 3^n \lambda_0^{-1/2} \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{\underline{L}^2(\Delta)}$$

and, by (6.62),

$$[\mathbf{A}_0^{1/2}(X - W)]_{\underline{B}_{2,1}^{-1}(\Delta)} \leq 3^d \|\mathbf{A}^{1/2}(X - W)\|_{\underline{L}^2(\Delta)} \sum_{k=-\infty}^n 3^k \max_{\Delta' \in \mathcal{Z}_k(\Delta)} |\mathbf{A}_0^{-1/2} \mathbf{A}(\Delta') \mathbf{A}_0^{-1/2}|^{1/2}.$$

Therefore, (6.84) follows by combining the above displays.

*Step 7.* We prove (6.74), which then completes the proof by Step 3. Combining (6.81), (6.83) and (6.84) with (6.64) (applied with  $s = 1$ ) provides us, after applying Young's inequality and reabsorption, that

$$\|\mathbf{A}^{1/2}(X - W)\|_{\underline{L}^2(\Delta)} \leq 2\mathcal{E}_1^\Delta(\Delta) \|\mathbf{s}_0^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\Delta)} + C(1 + \mathcal{E}_1^\Delta(\Delta)) 3^n \|\mathbf{s}_0 \nabla^2 u_{\text{hom}}\|_{\underline{L}^2(\Delta)}.$$

Summing over the simplexes yields (6.74) and completes the proof.  $\square$

We conclude this subsection by describing the properties of the finite volume correctors, defined, for every  $m \in \mathbb{N}$  and  $e \in \mathbb{R}^d$ , by

$$v(\cdot, \diamond_m, e) := v(\cdot, \diamond_m, 0, \mathbf{s}_*(\diamond_m)e).$$

**Lemma 6.10.** *For every  $m, n \in \mathbb{N}$  with  $n \leq m - 2$  and  $s \in (0, 1]$ , denote*

$$\begin{aligned} & \hat{\mathcal{E}}_{s,n}(\diamond_m; \mathbf{a}, \mathbf{a}_0) \\ & := (1 + \mathcal{E}_1(\diamond_m; \mathbf{a}, \mathbf{a}_0)) \max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \left( \mathcal{E}_1(z + \diamond_n; \mathbf{a}, \mathbf{a}_0) + 3^{-s(m-n)} (1 + \mathcal{E}_s(z + \diamond_n; \mathbf{a}, \mathbf{a}_0)) \right). \end{aligned} \quad (6.85)$$

*There exists a constant  $C(d) < \infty$  such that, we have, for every  $e \in \mathbb{R}^d$  and  $s \in (0, 1]$ , the estimates*

$$\begin{aligned} & s 3^{-ms} \left( [\mathbf{s}_0^{1/2}(\nabla v(\cdot, \diamond_m, e) - e)]_{\underline{B}_{2,1}^{-s}(\diamond_m)} + [\mathbf{s}_0^{-1/2}(\mathbf{a}\nabla v(\cdot, \diamond_m, e) - \mathbf{a}_0 e)]_{\underline{B}_{2,1}^{-s}(\diamond_m)} \right) \\ & \leq C \hat{\mathcal{E}}_{s,n}(\diamond_m; \mathbf{a}, \mathbf{a}_0) |\mathbf{s}_0^{1/2} e| \end{aligned} \quad (6.86)$$

*and, for the affine function  $\ell_e := e \cdot x + (v(\cdot, \diamond_m, e))_{\diamond_m}$ ,*

$$3^{-m} \|v(\cdot, \diamond_m, e) - \ell_e\|_{\underline{L}^2(\diamond_m)} \leq C \hat{\mathcal{E}}_{1,n}(\diamond_m; \mathbf{a}, \mathbf{a}_0) |\mathbf{q}_0 e|. \quad (6.87)$$

*Moreover, the energy of  $v(\cdot, \diamond_m, e)$  satisfies*

$$\left| \|\mathbf{s}^{1/2} \nabla v(\cdot, \diamond_m, e)\|_{\underline{L}^2(\diamond_m)}^2 - |\mathbf{s}_0^{1/2} e|^2 \right| \leq C \hat{\mathcal{E}}_{1,n}(\diamond_m; \mathbf{a}, \mathbf{a}_0) |\mathbf{s}_0^{1/2} e|^2. \quad (6.88)$$

*Proof.* Again, without loss of generality, we assume that  $\mathbf{k}_0 = 0$ . We first prove (6.86). Denote

$$p = 0, \quad q = \mathbf{s}_*(\diamond_m)e, \quad P = \begin{pmatrix} e \\ \mathbf{a}_0 e \end{pmatrix} \quad \text{and} \quad X_{n,z} = \begin{pmatrix} \nabla v(\cdot, z + \diamond_n, 0, q) \\ \mathbf{a} \nabla v(\cdot, z + \diamond_n, 0, q) \end{pmatrix},$$

and  $X_m := X_{m,0}$ . For any Lipschitz domain  $U$ , we also denote, for short,

$$H(U) := \left( \sup_{|e'| \leq 1} J(U, \mathbf{s}_0^{-1/2} e', \mathbf{a}_0^t \mathbf{s}_0^{-1/2} e') + \sup_{|e'| \leq 1} J^*(U, \mathbf{s}_0^{-1/2} e', \mathbf{a}_0 \mathbf{s}_0^{-1/2} e') \right)^{1/2}.$$

Notice that, by a direct computation using (6.1), we have

$$\begin{aligned} \left| \frac{1}{2} \|\mathbf{A}^{1/2} X_m\|_{\underline{L}^2(\diamond_m)}^2 - |\mathbf{s}_0^{1/2} e|^2 \right| &= |e \cdot (\mathbf{s}_*(\diamond_m) - \mathbf{s}_0) e| \leq |\mathbf{s}_0^{-1/2} \mathbf{s}_*(\diamond_m) \mathbf{s}_0^{-1/2} - \text{Id}| |\mathbf{s}_0^{1/2} e|^2 \\ &\leq C(H(\diamond_m) + H^2(\diamond_m)) |\mathbf{s}_0^{1/2} e|^2, \end{aligned}$$

which yields (6.88).

We then show the weak norm bound (6.86). Notice that (6.87) follows immediately from this by the Poincaré inequality in (2.128) (applied with  $s = 0$ ). To show (6.86), we first get, similarly to (6.75),

$$\begin{aligned} s3^{-sm} [\mathbf{A}_0^{1/2} (X_m - P)]_{\underline{\dot{B}}_{2,1}^{-s}(\diamond_m)} &\leq C \left( \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} |(\mathbf{A}_0^{1/2} (X_m - P))_{z + \diamond_n}|^2 \right)^{1/2} \\ &\quad + s3^{-sm} \left( \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} [\mathbf{A}_0^{1/2} (X_m - P)]_{\underline{\dot{B}}_{2,1}^{-s}(z + \diamond_n)}^2 \right)^{1/2}. \end{aligned} \quad (6.89)$$

For the second term in (6.89) we have, by a similar estimate as in (6.64),

$$\begin{aligned} s3^{-sn} [\mathbf{A}_0^{1/2} (X_m - P)]_{\underline{\dot{B}}_{2,1}^{-s}(z + \diamond_n)} &\leq C |\mathbf{s}_0^{1/2} e| + C \|\mathbf{A}^{1/2} X_m\|_{\underline{L}^2(z + \diamond_n)} \left( s \sum_{k=-\infty}^n 3^{-s(n-k)} \max_{z \in 3^k \mathbb{L}_0 \cap \diamond_m} |\mathbf{A}_0^{-1/2} \mathbf{A}(z + \diamond_n) \mathbf{A}_0^{-1/2}|^{1/2} \right) \\ &\leq C |\mathbf{s}_0^{1/2} e| + C(1 + \mathcal{E}_s(z + \diamond_n)) \|\mathbf{A}^{1/2} X_m\|_{\underline{L}^2(z + \diamond_n)}, \end{aligned}$$

and thus

$$\begin{aligned} s3^{-sm} \left( \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} [\mathbf{A}_0^{1/2} (X_m - P)]_{\underline{\dot{B}}_{2,1}^{-s}(z + \diamond_n)}^2 \right)^{1/2} &\leq C(1 + H(\diamond_m)) 3^{-s(m-n)} \left( 1 + \max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_s(z + \diamond_n) \right) |\mathbf{s}_0^{1/2} e|. \end{aligned} \quad (6.90)$$

For the first term on the right in (6.89), if  $\max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(z + \diamond_n) \geq 1$ , we get, as above,

$$\left( \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} |(\mathbf{A}_0^{1/2} (X_m - P))_{z + \diamond_n}|^2 \right)^{1/2} \leq C(1 + H(\diamond_m)) \left( 1 + \max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_s(z + \diamond_n) \right) |\mathbf{s}_0^{1/2} e|.$$

Combining the last three displays yields (6.86) if  $\max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(z + \diamond_n) > 1$ .

Assume then that  $\max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(z + \diamond_n) \leq 1$ , so that, by subadditivity, also  $H(\diamond_m) \leq 1$ . By the triangle inequality we have

$$\begin{aligned} & \frac{1}{3} |\mathbf{A}_0^{1/2} (X_m - P)_{z + \diamond_n}|^2 \\ & \leq |\mathbf{A}_0^{1/2} (X_m - P)_{\diamond_m}|^2 + |\mathbf{A}_0^{1/2} ((X_{n,z})_{z + \diamond_n} - (X_m)_{\diamond_m})|^2 + |\mathbf{A}_0^{1/2} (X_{n,z} - X_m)_{z + \diamond_n}|^2. \end{aligned} \quad (6.91)$$

By (2.31) and (6.1), the first term can be estimated as

$$|\mathbf{A}_0^{1/2} (X_m - P)_{\diamond_m}| = \left| \begin{pmatrix} 0 \\ \mathbf{s}_0^{-1/2} (\mathbf{a}_*(\diamond_m) - \mathbf{a}_0) e \end{pmatrix} \right| \leq CH(\diamond_m) |\mathbf{s}_0^{1/2} e|. \quad (6.92)$$

To estimate the second term in (6.91), we write, using (2.33) and  $\mathbf{R} \mathbf{A}_0^{1/2} \mathbf{R} = \mathbf{A}_0^{-1/2}$ ,

$$\begin{aligned} \mathbf{A}_0^{1/2} ((X_{n,z})_{z + \diamond_n} - (X_m)_{\diamond_m}) &= \mathbf{A}_0^{1/2} \mathbf{R} (\mathbf{A}(z + \diamond_n) - \mathbf{A}(\diamond_m)) \begin{pmatrix} 0 \\ q \end{pmatrix} \\ &= \mathbf{R} (\mathbf{A}_0^{-1/2} (\mathbf{A}(z + \diamond_n) - \mathbf{A}(\diamond_m)) \mathbf{A}_0^{-1/2}) \mathbf{A}_0^{1/2} \begin{pmatrix} 0 \\ q \end{pmatrix} \end{aligned}$$

By a direct computation using (6.1), we deduce that

$$\begin{aligned} & |\mathbf{A}_0^{-1/2} \mathbf{A}(U) \mathbf{A}_0^{-1/2} - \mathbf{I}_{2d}| \\ & \leq 2 |\mathbf{s}_0^{-1/2} \mathbf{b}_{\mathbf{k}_0}(U) \mathbf{s}_0^{-1/2} - \mathbf{I}_d| + 2 |\mathbf{s}_0^{1/2} \mathbf{s}_*^{-1}(U) \mathbf{s}_0^{1/2} - \mathbf{I}_d| + 2 |\mathbf{s}_*^{-1/2}(U) (\mathbf{k}(U) - \mathbf{k}_0) \mathbf{s}_0^{-1/2}|^2 \\ & \leq 2 |\mathbf{s}_0^{-1} (\mathbf{s} - \mathbf{s}_*)(U)| + 4 |\mathbf{s}_{L,*}^{-1/2}(U) (\mathbf{k}(U) - \mathbf{k}_0) \mathbf{s}_0^{-1/2}|^2 \\ & \quad + 2 |\mathbf{s}_0^{1/2} \mathbf{s}_*^{-1}(U) \mathbf{s}_0^{1/2} - \mathbf{I}_d| + 2 |\mathbf{s}_0^{-1/2} \mathbf{s}_*(U) \mathbf{s}_0^{-1/2} - \mathbf{I}_d| \\ & \leq 2 |\mathbf{s}_0^{-1} (\mathbf{s} - \mathbf{s}_*)(U)| + 4 |\mathbf{s}_*^{-1/2}(U) (\mathbf{k}(U) - \mathbf{k}_0) \mathbf{s}_0^{-1/2}|^2 \\ & \quad + 4 |\mathbf{s}_*^{-1/2}(U) (\mathbf{s}_*(U) - \mathbf{s}_0) \mathbf{s}_0^{-1/2}|^2 + 4 |\mathbf{s}_*^{-1/2}(U) (\mathbf{s}_*(U) - \mathbf{s}_0) \mathbf{s}_0^{-1/2}| \\ & \leq 12 (H(U) + H^2(U)). \end{aligned}$$

Thus, since we assume  $\max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(z + \diamond_n) \leq 1$ , implying that  $H(z + \diamond_n) \leq 1$  as well, we get

$$\left( \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} |\mathbf{A}_0^{1/2} ((X_{n,z})_{z + \diamond_n} - (X_m)_{\diamond_m})|^2 \right)^{1/2} \leq C \left( \max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(z + \diamond_n) \right) |\mathbf{s}_0^{1/2} e|. \quad (6.93)$$

To estimate the third term in (6.91), similarly to (6.61), we obtain

$$|(\mathbf{A}_0^{1/2} (X_m - X_{n,z}))_{z + \diamond_n}|^2 \leq |\mathbf{A}_0^{-1/2} \mathbf{A}(z + \diamond_n) \mathbf{A}_0^{-1/2}| \|\mathbf{A}^{1/2} (X_{n,z} - X_m)\|_{\underline{L}^2(y + \diamond_n)}^2,$$

and hence

$$\begin{aligned} & \left( \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} |(\mathbf{A}_0^{1/2} (X_m - X_{n,z}))_{z + \diamond_n}|^2 \right)^{1/2} \\ & \leq C \left( 1 + \max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(z + \diamond_n) \right) \left( \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} (J(z + \diamond_n, 0, q) - J(\diamond_m, 0, q)) \right)^{1/2}. \end{aligned} \quad (6.94)$$

We further estimate, using (2.13), as

$$\begin{aligned}
& \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} J(z + \diamond_n, 0, q) - J(\diamond_m, 0, q) \\
&= \mathbf{A}_0^{1/2} \begin{pmatrix} 0 \\ q \end{pmatrix} \cdot \left( \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathbf{A}_0^{-1/2} (\mathbf{A}(z + \diamond_n) - \mathbf{A}(\diamond_m)) \mathbf{A}_0^{-1/2} \right) \mathbf{A}_0^{1/2} \begin{pmatrix} 0 \\ q \end{pmatrix} \\
&\leq 2 \left| \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathbf{A}_{0, \mathbf{k}_0}^{-1/2} (\mathbf{A}_{\mathbf{k}_0}(z + \diamond_n) - \mathbf{A}_{\mathbf{k}_0}(\diamond_m)) \mathbf{A}_{0, \mathbf{k}_0}^{-1/2} \right| |\mathbf{s}_0^{1/2} e|. \tag{6.95}
\end{aligned}$$

We then appeal to the following elementary facts. For any positive and symmetric matrix  $B \in \mathbb{R}^{2d \times 2d}$ , we have

$$\begin{aligned}
|B| &= \sup_{|P| \leq 1} P \cdot B P = \sup_{(|e| \vee |e'|)^2 \leq 2} \begin{pmatrix} -e + e' \\ e + e' \end{pmatrix} \cdot B \begin{pmatrix} -e + e' \\ e + e' \end{pmatrix} \\
&\leq 4 \sup_{|e| \leq 1} \begin{pmatrix} -e \\ e \end{pmatrix} \cdot B \begin{pmatrix} -e \\ e \end{pmatrix} + 4 \sup_{|e| \leq 1} \begin{pmatrix} e \\ e \end{pmatrix} \cdot B \begin{pmatrix} e \\ e \end{pmatrix},
\end{aligned}$$

and the identities

$$\begin{aligned}
& \begin{pmatrix} -e \\ e \end{pmatrix} \cdot \mathbf{A}_{0, \mathbf{k}_0}^{-1/2} (\mathbf{A}_{\mathbf{k}_0}(z + \diamond_n) - \mathbf{A}_{\mathbf{k}_0}(\diamond_m)) \mathbf{A}_{0, \mathbf{k}_0}^{-1/2} \begin{pmatrix} -e \\ e \end{pmatrix} \\
&= \begin{pmatrix} -e \\ e \end{pmatrix} \cdot \mathbf{A}_{0, \mathbf{k}_0}^{-1/2} (\mathbf{A}_{\mathbf{k}_0}(z + \diamond_n) - \mathbf{A}_{\mathbf{k}_0}(\diamond_m)) \mathbf{A}_{0, \mathbf{k}_0}^{-1/2} \begin{pmatrix} -e \\ e \end{pmatrix} \\
&\quad - \begin{pmatrix} -e \\ e \end{pmatrix} \cdot \mathbf{A}_{0, \mathbf{k}_0}^{-1/2} (\mathbf{A}_{\mathbf{k}_0}(z + \diamond_n) - \mathbf{A}_{\mathbf{k}_0}(\diamond_m)) \mathbf{A}_{0, \mathbf{k}_0}^{-1/2} \begin{pmatrix} -e \\ e \end{pmatrix} \\
&= 2J_{\mathbf{k}_0}(z + \diamond_n, \mathbf{s}_0^{-1/2} e, \mathbf{s}_0^{1/2} e) - 2J_{\mathbf{k}_0}(\diamond_m, \mathbf{s}_0^{-1/2} e, \mathbf{s}_0^{-1/2} e),
\end{aligned}$$

and similarly for the other term, but using now  $J^*$ . Since we assume that  $\mathcal{E} \leq 1$ , we get, by (6.94) and (6.95), the previous three displays and the subadditivity of  $J$  and  $J^*$ ,

$$\left( \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} |(\mathbf{A}_0^{1/2}(X_m - X_{n,z}))_{z + \diamond_n}|^2 \right)^{1/2} \leq C \max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(z + \diamond_n) |\mathbf{s}_0^{1/2} e|. \tag{6.96}$$

Combining thus (6.91) with (6.92), (6.93) and (6.96) yields

$$\left( \sum_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} |\mathbf{A}_0^{1/2}(X_m - P)_{z + \diamond_n}|^2 \right)^{1/2} \leq C \left( \max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \mathcal{E}_1(z + \diamond_n) \right) |\mathbf{s}_0^{1/2} e|.$$

This, together with (6.89) and (6.90), gives us

$$s 3^{-sm} [\mathbf{A}_0^{1/2}(X_m - P)]_{\tilde{\mathcal{B}}_{2,1}^{-s}(\diamond_m)} \leq C \max_{z \in 3^n \mathbb{L}_0 \cap \diamond_m} \left( \mathcal{E}_1(z + \diamond_n) + 3^{-s(m-n)} (1 + \mathcal{E}_s(z + \diamond_n)) \right) |\mathbf{s}_0^{1/2} e|,$$

which proves (6.86), completing the proof.  $\square$



**6.3. Large-scale regularity.** In view of the statements of harmonic approximation results in Propositions 6.7 and 6.9, as well as the coarse-graining estimates in Lemma 6.1 and Caccioppoli estimate in Lemma 6.6,  $\mathcal{E}_s$  defined in (6.2) plays a fundamental role what comes to the regularity properties of arbitrary solutions. In this subsection we establish many natural estimates assuming suitable smallness conditions.

To quantify the above discussion, we set

$$\delta_k(\mathbf{a}, \mathbf{a}_0) := \min_{j \in \mathbb{N} \cap [1, k]} \inf_{s \in (0, 1]} \max_{z \in 3^j \mathbb{L}_0 \cap \diamond_k} \left( \mathcal{E}_1(z + \diamond_j; \mathbf{a}, \mathbf{a}_0) + 3^{-s(k-j)} (1 + \mathcal{E}_{1-s}(z + \diamond_j; \mathbf{a}, \mathbf{a}_0)) \right). \quad (6.97)$$

Define consequently, for every  $m, n \in \mathbb{N}$  with  $n \leq m$  and  $t \in [1, \infty]$ , the "good" event

$$G_{n,m}^t(\delta; \mathbf{a}, \mathbf{a}_0) := \left\{ \left( \sum_{k=n}^m (\delta_k(\mathbf{a}, \mathbf{a}_0))^t \right)^{1/t} \leq \delta \right\}. \quad (6.98)$$

For  $t = \infty$  the interpretation is that the sum over  $k$  is replaced by the maximum over  $k \in \mathbb{N} \cap [n, m]$ . The motivation for the definition of  $\delta_k$  in (6.97) is that it controls the error terms in the Caccioppoli estimate, Lemma 6.6, the harmonic approximation error in Propositions 6.7 as well as the right-sides of (6.87) and (6.88) provided that  $\delta_k \leq 1$ . This also motivates us to study the good event  $G_{n,m}^t(\delta; \mathbf{a}, \mathbf{a}_0)$  in (6.98). In practice, we will use it by choosing the parameters  $j$  and  $s$  appropriately. For instance, many times it suffices to take  $s' = \rho$  and then  $j = k - h$  with  $h \in \mathbb{N}$  being the smallest integer such that  $3^{-\rho h} \leq c(d)$  for some given small constant  $c(d)$ . Thus bounding  $\delta_k$  becomes easy since the necessary union bounds are taken only over a relatively few scales depending on  $d$ .

The first lemma states that under the event  $G_{n,m}^\infty(\delta)$  we have  $C^{0,\alpha}$ -estimates, and under the event  $G_{n,m}^1(\delta)$  this can be upgraded to  $C^{0,1}$ -type estimate.

**Lemma 6.11.** *There exist constants  $C(d) < \infty$  and  $c(d) \in (0, 1)$  such that for every  $m, n \in \mathbb{N}$  with  $n < m$ ,  $\delta \in (0, c]$  and  $u \in \mathcal{A}(\square_m)$ , we have, for every  $s \in (0, 1)$ , the implications*

$$\|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_n)} \mathbf{1}_{G_{n,m}^\infty(\delta; \mathbf{a}, \mathbf{a}_0)} \leq C \exp(C\delta(m-n)) \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)} \quad (6.99)$$

and

$$\|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_n)} \mathbf{1}_{G_{n,m}^1(c; \mathbf{a}, \mathbf{a}_0)} \leq C \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \quad (6.100)$$

*Proof.* Fix  $m, n \in \mathbb{N}$  with  $n < m$  and  $u \in \mathcal{A}(\square_m)$ . Denote, for short, for every  $k \in \mathbb{N}$ ,

$$\delta_k := \min_{j \in \mathbb{N} \cap [1, k]} \inf_{s \in (0, 1]} \max_{z \in 3^j \mathbb{L}_0 \cap \diamond_k} \left( \mathcal{E}_1(z + \diamond_j; \mathbf{a}, \mathbf{a}_0) + 3^{-s(k-j)} (1 + \mathcal{E}_{1-s}(z + \diamond_j; \mathbf{a}, \mathbf{a}_0)) \right). \quad (6.101)$$

and

$$E_k := \lambda_0^{1/2} \inf_{\ell \text{ affine}} 3^{-k} \|u - \ell\|_{\underline{L}^2(\diamond_k)} \quad \text{and} \quad D_k := \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_k)}.$$

Let  $\ell^{(k)}$  denote the affine function realizing the minimum in  $E_k$ . Throughout the proof we assume the event  $G_{n,m}^\infty(\delta; \mathbf{a}, \mathbf{a}_0)$ , which implies that  $\delta_k \leq \delta$  for every  $k \in \mathbb{N} \cap [n, m]$ .

*Step 1.* We show that there exists  $N(d) \in \mathbb{N}$  and  $C(d) < \infty$  such that, for every  $k \in \mathbb{N}$  with  $n \leq k \leq m$ ,

$$E_{k-N} \leq \frac{1}{8} E_k + C \delta_k D_k. \quad (6.102)$$

By Proposition 6.7 there exist a constant  $C(d) < \infty$  and, for every  $k \in \mathbb{N}$ , an  $\mathbf{a}_0$ -harmonic function  $\bar{u}_k$  in  $\diamond_{k-1}$  such that

$$\lambda_0^{1/2} 3^{-k} \|u - \bar{u}_k\|_{\underline{L}^2(\diamond_{k-1})} \leq C \delta_k \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_k)}.$$

By the regularity of  $\mathbf{a}_0$ -harmonic functions, there exists  $C(d) < \infty$  such that

$$\begin{aligned} E_{k-N} &\leq \lambda_0^{1/2} \inf_{\ell \text{ affine}} 3^{N-k} \|\bar{u}_k - \ell\|_{\underline{L}^2(\diamond_{k-N})} + \lambda_0^{1/2} 3^{N-k} \|u - \bar{u}_k\|_{\underline{L}^2(\diamond_{k-N})} \\ &\leq C \lambda_0^{1/2} 3^{-N-k} \inf_{\ell \text{ affine}} \|\bar{u}_k - \ell\|_{\underline{L}^2(\diamond_{k-1})} + \lambda_0^{1/2} 3^{(d/2+1)N-k} \|u - \bar{u}_k\|_{\underline{L}^2(\diamond_{k-1})} \\ &\leq C 3^{-N} E_k + C \lambda_0^{1/2} \left( 3^{-k-N} + 3^{(d/2+1)N-k} \right) \|u - \bar{u}_k\|_{\underline{L}^2(\diamond_{k-1})} \\ &\leq C 3^{-N} E_k + C 3^{(d/2+1)N} \delta_k D_k. \end{aligned} \quad (6.103)$$

We choose  $N(d) \in \mathbb{N}$  so large that  $C_{(6.103)} 3^{-N} \leq 1/8$ . Thus, the previous display implies (6.102).

*Step 2.* We next show that there exist constants  $C(d) < \infty$  such that, for every  $h \in \mathbb{N}$  with  $h \leq n$ ,

$$D_h \leq C D_m + C \sum_{j=h+1}^m \delta_j D_j. \quad (6.104)$$

To see this, we first sum over  $j$  in (6.102) to get

$$\sum_{j=-1}^{\lfloor N^{-1}(m-h) \rfloor - 1} E_{h+jN} \leq \frac{1}{2} \sum_{j=0}^{\lfloor N^{-1}(m-h) \rfloor} E_{h+jN} + C \sum_{j=h}^m \delta_j D_j.$$

Reorganizing and reabsorbing then leads to

$$\sum_{j=h}^m E_j \leq C E_m + C \sum_{j=h+1}^m \delta_j D_j.$$

We then obtain, for every  $j \in \mathbb{N} \cap [n, m]$ ,

$$|\nabla \ell^{(j)} - \nabla \ell^{(m)}| \leq C_d \sum_{j=n}^{m-1} 3^{-j} \|\ell^{(j+1)} - \ell^{(j)}\|_{\underline{L}^2(\diamond_j)} \leq C_d \sum_{j=n}^m E_j \leq C_d E_m + C \sum_{j=h+1}^m \delta_j D_j. \quad (6.105)$$

We may choose  $\ell^{(m)}(0) = (u)_{\diamond_m}$ , and then use the Poincaré inequality (2.128) and (6.10) to get

$$E_m \leq C D_m.$$

Consequently, by the coarse-grained Caccioppoli estimate, Lemma 6.6, we deduce that

$$D_h \leq C |\nabla \ell^{(h+1)}| + C E_{h+1} + C \delta_{h+1} D_{h+1} \leq C D_m + C \sum_{j=h+1}^m \delta_j D_j,$$

which is (6.104).

*Step 3.* We show (6.99). Assume inductively that there exist constants  $H, K \in [1, \infty)$ , to be determined by means of  $d$ , such that, for some  $h \in \mathbb{N}$  with  $h < k$ ,

$$\sup_{k \in \mathbb{N} \cap [h+1, m]} 3^{-K\delta(m-k)} D_k \leq H D_m. \quad (6.106)$$

The induction assumption together with (6.104) and  $\delta \leq c := K^{-1}$  implies that

$$D_h \leq CD_m + C\delta H \sum_{k=h+1}^m 3^{K\delta(m-k)} D_m \leq H 3^{K\delta(m-h)} D_m \left( CH^{-1} 3^{-K\delta(m-h)} + CK^{-1} \right),$$

and thus (6.106) follows by choosing  $H, K$  large enough.

*Step 4.* We finally show (6.100). Assume that

$$\sum_{k=n}^m \delta_k \leq \delta_0 \quad (6.107)$$

for some  $\delta_0$  to be fixed. Assume inductively that, for a given large constant  $H(d) \in [1, \infty)$ , we have that, for some  $h \in \mathbb{N}$  with  $n \leq h < m$  we have that

$$\sup_{j \in \mathbb{N} \cap [h+1, m]} D_j \leq HD_m. \quad (6.108)$$

By (6.104), (6.107) and (6.108) we then deduce that

$$D_h \leq CD_m + C\delta_0 HD_m.$$

Choosing thus  $\delta_0 = (2C)^{-1}$  and  $H = 2C$ , we obtain the induction step and complete the proof.  $\square$

The next lemma shows that the finite-volume corrector  $v(\cdot, \diamond_m, e)$  constructed in Lemma 6.10, under the good event  $G_{n,m}^\infty(\delta)$ , is also flat at the scale  $n$ . The precise statement is as follows.

**Lemma 6.12** (Flatness at every scale). *There exist constants  $C(d) < \infty$  and  $c(d) \in (0, (2C)^{-1}]$  such that if  $\delta \in (0, c]$ , then, for every  $n, m \in \mathbb{N}$ , under the event  $G_{n,m}^\infty(\delta; \mathbf{a}, \mathbf{a}_0)$  defined by (6.98), there exists a linear map  $e \mapsto Q_n[e; m]$  such that, for  $\ell_e(x) := x \cdot e$ , we have*

$$3^{-n} \|v(\cdot, \diamond_m, Q_n[e; m]) - (v(\cdot, \diamond_n, Q_n[e; m]))_{\diamond_n} - \ell_e\|_{\underline{L}^2(\diamond_n)} \mathbf{1}_{G_{n,m}^\infty(\delta; \mathbf{a}, \mathbf{a}_0)} \leq C\delta |\mathbf{q}_0 e|. \quad (6.109)$$

Moreover, we have under the event  $G_{n,m}^\infty(\delta; \mathbf{a}, \mathbf{a}_0)$  that, for every  $e \in \mathbb{R}^d$ ,

$$C^{-1} |\mathbf{s}_0^{1/2} e| \leq \|\mathbf{s}^{1/2} \nabla v(\cdot, \diamond_m, Q_n[e; m])\|_{\underline{L}^2(\diamond_n)} \leq C |\mathbf{s}_0^{1/2} e|. \quad (6.110)$$

Finally, for every  $k \in \mathbb{N}$  with  $n \leq k \leq m$  and  $e \in \mathbb{R}^d$ , under the event  $G_{n,m}^\infty(\delta; \mathbf{a}, \mathbf{a}_0)$ ,

$$(1 + C\delta)^{-(k-n)} \leq \frac{|\mathbf{s}_0 Q_k[e; m]|}{|\mathbf{s}_0 Q_n[e; m]|} \leq (1 - C\delta)^{k-n} \quad \text{and} \quad |Q_m[\cdot; m] - \text{Id}| \leq C\delta \leq \frac{1}{2}, \quad (6.111)$$

and, under the event  $G_{n,m}^1(\delta; \mathbf{a}, \mathbf{a}_0)$ ,

$$|Q_n[\cdot; m] - \text{Id}| \leq C\delta \leq \frac{1}{2}. \quad (6.112)$$

*Proof.* Define  $\ell^{(n)}[e; m]$  to be the affine function minimizing the following quantity

$$E_n[e; m] := \inf_{\ell \text{ affine}} 3^{-n} \|v(\cdot, \diamond_m, e) - \ell\|_{\underline{L}^2(\diamond_n)} := 3^{-n} \|v(\cdot, \diamond_m, e) - \ell^{(n)}(e; m)\|_{\underline{L}^2(\diamond_n)}.$$

Denote, for short,  $v = v(\cdot, \diamond_m, e)$ . Denote also  $P_n[e; m] := \nabla \ell^{(n)}[e; m]$ . The mapping  $e \mapsto P_n[e; m]$  is linear by the linearity of  $e \mapsto \nabla v(\cdot, \square_m, e)$  and (6.87). We claim that, for  $k \in \mathbb{N} \cap [n, m]$ ,

$$E_k[e; m] \leq K\delta |\mathbf{q}_0 P_k[e; m]| \quad (6.113)$$

for a large constant  $K(d)$  to be fixed. Letting  $N(d) \in \mathbb{N}$  be as in Step 1 of the proof Lemma 6.11, we have, for every  $k \in \mathbb{N} \cap [n \vee (m - N), m]$ ,

$$|\mathbf{q}_0(e - P_k[e; m])| \leq C3^{-k} \|\ell_k[e; m] - \ell_e\|_{\underline{L}^2(\diamond_k)} \leq C3^{(d/2+1)N} 3^{-m} \|v - \ell_e\|_{\underline{L}^2(\diamond_m)} \leq C\delta |\mathbf{q}_0 e|. \quad (6.114)$$

Thus, if  $C_{(6.114)}c \leq 1/2$  and  $K \geq 2C_{(6.114)}$ , we obtain (6.113) for  $k \in \mathbb{N} \cap [n \vee (m - N), m]$ .

Next, if  $n < m - N$ , we then assume inductively that there exists  $h \in \mathbb{N} \cap [n, m - N]$  such that (6.113) is valid for every  $k \in \mathbb{N}$  with  $h + 1 \leq k \leq m$ . Since  $m$  and  $e$  are fixed, we drop them from the notation for both  $\ell$  and  $E$ . By Step 1 of the proof of Lemma 6.11 and the induction assumption, there exists  $N(d) \in \mathbb{N}$  and  $C(d) < \infty$  such that,

$$E_h \leq \frac{1}{8}E_{h+N} + C\delta\lambda_0^{-1/2} \|\mathbf{s}^{1/2}\nabla v\|_{\underline{L}^2(\diamond_{h+N})} \leq \frac{K}{8}\delta |\mathbf{q}_0 \nabla \ell^{(h+N)}| + C\delta\lambda_0^{-1/2} \|\mathbf{s}^{1/2}\nabla v\|_{\underline{L}^2(\diamond_{h+N})}. \quad (6.115)$$

In view of our claim (6.113), we define, for  $k \in \mathbb{N}$ , the composite quantity

$$F_k := E_k - K\delta |\mathbf{q}_0 \nabla \ell^{(k)}|,$$

and our goal is to show that  $F_h \leq 0$ . The induction assumption guarantees that  $F_k \leq 0$  for  $k \in \mathbb{N} \cap (h, m]$ . Notice that we then get by the triangle inequality, for every  $k \in \mathbb{N} \cap [n, m]$ ,

$$|\mathbf{q}_0 \nabla \ell^{(k+1)}| \leq |\mathbf{q}_0 \nabla \ell^{(k)}| + CE_{k+1} \leq |\mathbf{q}_0 \nabla \ell^{(k)}| + CK\delta |\mathbf{q}_0 \nabla \ell^{(k+1)}| + CF_{k+1}, \quad (6.116)$$

and thus, if  $C_{(6.116)}Kc \leq 1/2$ , we have

$$|\mathbf{q}_0 \nabla \ell^{(k+1)}| \leq (1 + \delta) |\mathbf{q}_0 \nabla \ell^{(k)}| + CF_{k+1} \quad \text{with} \quad \delta := \frac{C_{(6.116)}K\delta}{1 - C_{(6.116)}K\delta}. \quad (6.117)$$

By the induction assumption and iteration we then also get, for every  $k \in \mathbb{N}$  with  $k > h$ ,

$$|\mathbf{q}_0 \nabla \ell^{(k)}| \leq (1 + \delta)^{h-k} |\mathbf{q}_0 \nabla \ell^{(h)}| \quad \text{and} \quad |\mathbf{q}_0 e| \leq C(1 + \delta)^{m-h} |\mathbf{q}_0 \nabla \ell^{(h)}|.$$

By iterating the Caccioppoli inequality, Lemma 6.6, and using (6.88), we obtain

$$\begin{aligned} \lambda_0^{-1/2} \|\mathbf{s}^{1/2}\nabla v\|_{\underline{L}^2(\diamond_{h+N})} &\leq C3^{-n} \|v - (v)_{\diamond_{h+N+1}}\|_{\underline{L}^2(\diamond_{h+N+1})} + C\lambda_0^{-1/2} \delta^{1/2} \|\mathbf{s}^{1/2}\nabla v\|_{\underline{L}^2(\diamond_{h+N+1})} \\ &\leq C \sum_{k=h+N+1}^m (C\delta^{1/2})^{k-(h+N+1)} (F_k + |\mathbf{q}_0 \nabla \ell^{(k)}|) + C(C\delta^{1/2})^{m-h} |\mathbf{q}_0 e| \\ &\leq C \sum_{k=h+N+1}^m (C\delta^{1/2})^{k-(h+N+1)} |\mathbf{q}_0 \nabla \ell^{(k)}| + C(C\delta^{1/2})^{m-h} |\mathbf{q}_0 e| \end{aligned} \quad (6.118)$$

since  $F_k \leq 0$  for  $k \in \mathbb{N} \cap [h + 1, m]$ . Taking  $c$  small enough so that  $C_{(6.118)}(1 + \delta)c^{1/2} \leq 1/2$ , we then obtain

$$\|\mathbf{s}^{1/2}\nabla v\|_{\underline{L}^2(\diamond_{h+N})} \leq C(1 + \delta)^N |\mathbf{s}_0 \nabla \ell^{(h)}|. \quad (6.119)$$

Inserting the above three displays into (6.115) yields

$$F_h \leq -K\delta |\mathbf{q}_0 \nabla \ell^{(h)}| + (1 + \delta)^N \left( \frac{K\delta}{8} + C\delta \right) |\mathbf{q}_0 \nabla \ell^{(h)}| \quad (6.120)$$

with  $\delta$  as in (6.117). Taking  $K \geq 8C_{(6.120)}$  and then  $c$  so small that  $(1 + \delta)^N \leq 2$  and  $C_{(6.120)}Kc \leq 1$ , we deduce that  $F_h \leq 0$ , proving the induction step and establishing (6.113).

Next, similarly to (6.116), we obtain

$$|\mathbf{q}_0 \nabla \ell^{(k)} - \mathbf{q}_0 \nabla \ell^{(k+1)}| \leq C E_k + C E_{k+1} \leq C(\delta_k + \delta_{k+1})(|\mathbf{q}_0 \nabla \ell^{(k)}| + |\mathbf{q}_0 \nabla \ell^{(k+1)}|),$$

which gives us by (6.114) that, for every  $k \in \mathbb{N} \cap [n, m-1]$ ,

$$1 - C\delta \leq \frac{|\mathbf{q}_0 \nabla \ell^{(k+1)}|}{|\mathbf{q}_0 \nabla \ell^{(k)}|} \leq 1 + C\delta \implies (1 - C\delta)^{n+1} \leq \frac{|\mathbf{q}_0 P_n[e; m]|}{|\mathbf{q}_0 e|} \leq (1 + C\delta)^{n+1}. \quad (6.121)$$

Thus  $e \mapsto P_n[e; m]$  has full rank and is invertible, and we define  $Q_n[\cdot; m] = P_n^{-1}[\cdot; m]$ . The estimate (6.111) then follows from the above display by increasing the constant  $C$ .

Next, the upper bound in (6.110) follows by (6.118) and (6.121) since  $F_h \leq 0$  for every  $h \in \mathbb{N} \cap [n, m]$ . To see the lower bound, we have by (2.128) and (6.10)

$$|\mathbf{q}_0 P_n[e; m]| \leq C 3^{-n} \|v - (v)_{\diamond_n}\|_{\underline{L}^2(\diamond_n)} + C E_n \leq C \lambda_0^{-1/2} \|\mathbf{s}^{1/2} \nabla v\|_{\underline{L}^2(\diamond_n)} + C \delta |\mathbf{q}_0 P_n[e; m]|, \quad (6.122)$$

and thus the lower bound in (6.110) follows provided that  $C_{(6.122)} c \leq 1/2$ .

Finally, by (6.105), Lipschitz estimate (6.99) and (6.110), we have, under the event  $G_{n,m}^1(\delta)$  with small enough  $\delta_0(d)$  and  $\delta \in (0, \delta_0]$ , that

$$|\mathbf{q}_0(\nabla \ell_n[e; m] - \nabla \ell_m[e; m])| \leq C \inf_{\ell \text{ affine}} 3^{-m} \|v - \ell\|_{\underline{L}^2(\diamond_m)} + C \lambda_0^{-1/2} \delta |\mathbf{s}_0^{1/2} e| \leq C \delta |\mathbf{q}_0^{1/2} e|.$$

By (6.114) we have that  $|\mathbf{q}_0(e - \nabla \ell_m[e; m])| \leq C \delta |\mathbf{q}_0^{1/2} e|$ , and thus

$$|\mathbf{q}_0 P_{n,m}[\mathbf{q}_0^{-1} e; m] - e| \leq C \delta |e| \implies |P_{n,m}[\cdot; m] - \text{Id}| \leq C \delta \leq \frac{1}{2}.$$

Since  $Q_{n,m} = P_{n,m}^{-1}$ , we get (6.112). The proof is complete.  $\square$

We can use the previous lemma to show the large-scale  $C^{1,\gamma}$ -estimate stating that locally any solution is close to some finite volume corrector.

**Lemma 6.13** (Large-scale  $C^{1,\alpha}$  regularity). *For every  $\alpha \in (0, 1)$  there exist constants  $C(\alpha, d) < \infty$  and  $c(\alpha, d) \in (0, 1)$  so that the following statement is valid. For every  $m, n \in \mathbb{N}$  with  $n < m$ ,  $\delta \in (0, c]$  and  $u \in \mathcal{A}(\square_m)$ , there exists  $e \in \mathbb{R}^d$  such that, for every  $k \in \mathbb{N}$  with  $n \leq k \leq m$ , we have*

$$\|\mathbf{s}^{1/2} \nabla(u - v(\cdot, \diamond_m, e))\|_{\underline{L}^2(\diamond_k)} \mathbf{1}_{G_{n,m}^\infty(\delta)} \leq C 3^{-\alpha(m-k)} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \quad (6.123)$$

*Proof.* Throughout the proof, we fix  $m, n \in \mathbb{N}$  with  $n < m$  and assume the good event  $G_{n,m}^\infty(\delta)$  as in (6.98) for  $\delta \in (0, \theta_0]$  with  $\theta_0$  to be fixed.

Let  $u \in \mathcal{A}(\diamond_m)$ , and let  $e_k \in \mathbb{R}^d$  be such that

$$E_k := \|\mathbf{s}^{1/2} \nabla(u - v(\cdot, \diamond_m, e_k))\|_{\underline{L}^2(\diamond_k)} := \inf_{e \in \mathbb{R}^d} \|\mathbf{s}^{1/2} \nabla(u - v(\cdot, \diamond_m, e))\|_{\underline{L}^2(\diamond_k)}.$$

*Step 1.* We first show that, for every  $\alpha \in [1/2, 1)$ , there exists constants  $C(\alpha, d)$  and  $c_0(\alpha, d) \in (0, 1)$  such that if  $\delta \in (0, c_0]$ , then

$$E_n \leq C 3^{-\alpha(m-n)} E_m. \quad (6.124)$$

Set  $u_k := u - v(\cdot, \diamond_m, e_k)$ . By the harmonic approximation, Proposition 6.7, we find a harmonic function  $\bar{u}_k$  such that

$$3^{-k} \|u_k - \bar{u}_k\|_{\underline{L}^2(\diamond_{k-1})} \leq C \delta \lambda_0^{-1/2} \|\mathbf{s}^{1/2} \nabla u_k\|_{\underline{L}^2(\diamond_k)}.$$

Notice that, by the triangle inequality, (2.128), (6.10) and the above display, we get

$$3^{-k} \|\bar{u}_k - (\bar{u}_k)_{\diamond_{k-1}}\|_{\underline{L}^2(\diamond_{k-1})} \leq C \lambda_0^{-1/2} \|\mathbf{s}^{1/2} \nabla u_k\|_{\underline{L}^2(\diamond_k)}.$$

Now, by the regularity of  $\mathbf{a}_0$ -harmonic functions, there is an affine function  $\ell^{(k)}$  such that

$$\|\bar{u}_k - \ell^{(k)}\|_{\underline{L}^2(\diamond_{k-h})} \leq C 3^{-2h} \|\bar{u}_k - (\bar{u}_k)_{\diamond_{k-1}}\|_{\underline{L}^2(\diamond_{k-1})} \leq C \lambda_0^{-1/2} 3^{k-2h} \|\mathbf{s}^{1/2} \nabla u_k\|_{\underline{L}^2(\diamond_k)}.$$

The affine function above satisfies

$$|\mathbf{q}_0 \nabla \ell^{(k)}| \leq \|\mathbf{q}_0 \nabla \bar{u}_k\|_{L^\infty(\diamond_{k-2})} \leq C 3^{-k} \|\bar{u}_k - (\bar{u}_k)_{\diamond_{k-1}}\|_{L^\infty(\diamond_{k-1})} \leq C \lambda_0^{-1/2} \|\mathbf{s}^{1/2} \nabla u_k\|_{\underline{L}^2(\diamond_k)}.$$

Lemma 6.12 provides us  $\tilde{v}_k := v(\cdot, \diamond_m, \tilde{e}_k)$ , which is a good approximant of  $\ell^{(k)}$  so that

$$3^{-k} \|\tilde{v}_k - \ell^{(k)}\|_{\underline{L}^2(\diamond_{k-1})} \leq C \delta |\mathbf{q}_0 \nabla \ell^{(k)}|.$$

Thus, by the triangle inequality,

$$\|u_k - \tilde{v}_k\|_{\underline{L}^2(\diamond_{k+1-h})} \leq C 3^k \lambda_0^{-1/2} (3^{-2h} + C \delta 3^{\frac{d}{2}h}) \|\mathbf{s}^{1/2} \nabla u_k\|_{\underline{L}^2(\diamond_k)}.$$

The Caccioppoli inequality, Lemma 6.6, yields

$$E_{k-h} \leq \|\nabla(u_k - \tilde{v}_k)\|_{\underline{L}^2(\diamond_{k-h})} \leq C(3^{-h} + C \delta 3^{(\frac{d}{2}+1)h}) E_k. \quad (6.125)$$

Choosing  $h_0(\alpha, d)$  be the smallest integer such  $3^{-(1-\alpha)h_0} C_{(6.125)} \leq 1/4$  and then requiring that  $\theta_0$  is so small that  $C_{(6.125)} c_0 3^{(\frac{d}{2}+2)h_0} \leq 1/4$ , we deduce that

$$E_{k-h_0} \leq \frac{1}{2} \cdot 3^{-\alpha h_0} E_k.$$

An iteration argument then proves (6.124).

*Step 2.* We prove (6.123). Fix  $\alpha \in [1/2, 1]$ , and assume that  $c_0$  is small enough so that (6.124) is valid for  $\delta \in (0, c_0]$ . By (6.111) and (6.110), we deduce that there is  $c_0(d) \in (0, 1)$  such that  $\delta \leq c_0$  implies, for every  $k, j$  with  $n \leq j \leq k \leq m$  and  $e \in \mathbb{R}^d$ ,

$$\|\mathbf{s}^{1/2} \nabla v(\cdot, \diamond_m, e)\|_{\underline{L}^2(\diamond_k)} \leq C 3^{\frac{1}{4}(k-j)} \|\mathbf{s}^{1/2} \nabla v(\cdot, \diamond_m, e)\|_{\underline{L}^2(\diamond_j)}.$$

It follows that, since  $\alpha \geq 1/2$ ,

$$\begin{aligned} \|\mathbf{s}^{1/2} \nabla(v(\cdot, \diamond_m, e_j - e_{j+1}))\|_{\underline{L}^2(\diamond_k)} &\leq C 3^{\frac{1}{4}(k-j)} \|\mathbf{s}^{1/2} \nabla v(\cdot, \diamond_m, e_j - e_{j+1})\|_{\underline{L}^2(\diamond_j)} \\ &\leq C 3^{\frac{1}{4}(k-j)} 3^{-\alpha(m-j)} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)} \\ &\leq C 3^{-\frac{1}{4}(k-j)} 3^{-\alpha(m-k)} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \end{aligned}$$

Thus, by the telescope summation and the triangle inequality,

$$\|\mathbf{s}^{1/2} \nabla v(\cdot, \diamond_m, e_n - e_k)\|_{\underline{L}^2(\diamond_k)} \leq C 3^{-\alpha(m-k)} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}.$$

Therefore, (6.123) follows by the triangle inequality, the above display and (6.124) by taking  $e := e_n$ . The proof is complete.  $\square$

**6.4. Estimates for correctors.** In this subsection, we complete the proof of Theorem B. We first construct the global correctors. For this, we take  $\mathbf{a}_0 := \bar{\mathbf{a}}$  in the previous section. By taking  $s := 1 - \frac{1}{2}(\gamma + \kappa)$  and  $\theta := \kappa/32$  in Corollary 4.5, we obtain by (6.7) that

$$\begin{aligned} \delta_k &:= \min_{j \in \mathbb{N} \cap [1, k]} \inf_{s \in (0, 1]} \max_{z \in 3^j \mathbb{L}_0 \cap \diamond_k} \left( \mathcal{E}_1(z + \diamond_j; \mathbf{a}, \bar{\mathbf{a}}) + 3^{-s(k-j)} (1 + \mathcal{E}_{1-s}(z + \diamond_j; \mathbf{a}, \bar{\mathbf{a}})) \right) \\ &\leq 2 \min_{j \in \mathbb{N} \cap [1, k]} \left( \delta 3^{\frac{1}{2}(\gamma + \kappa)(k-j)} \left( \frac{\mathcal{Z}_{\delta, s} \vee \mathcal{S}}{3^k} \right)^\theta + 2 \cdot 3^{-(1-\gamma/2-\kappa/2)(k-j)} \right). \end{aligned}$$

Choosing  $j$  appropriately, we get

$$3^{j-1} \leq \delta 3^{(1-\theta)k} (\mathcal{Z}_{\delta, s} \vee \mathcal{S})^\theta \leq 3^j \implies \delta_k \leq 9\delta^{1/2} \left( \frac{\mathcal{Z}_{\delta, s} \vee \mathcal{S}}{3^k} \right)^{\theta(1-\frac{1}{2}(\gamma+\kappa))}.$$

Relabelling  $\theta(1 - \frac{1}{2}(\gamma + \kappa))$  as  $\theta$  and taking  $\mathcal{X} = \mathcal{Z}_{\delta_1, s} \vee \mathcal{S}$  with  $\delta_1 = (C\theta^{-1})^{-2}\delta^2$  for large enough constant  $C(d) < \infty$ , we obtain that

$$\delta_k \leq 9\delta_1^{1/2} (3^{-k} \mathcal{X})^\theta \quad \text{and} \quad \sum_{k=m}^{\infty} \delta_k \leq C\theta^{-1} \delta_1^{1/2} (3^{-m} \mathcal{X})^\theta \leq \delta (3^{-m} \mathcal{X})^\theta. \quad (6.126)$$

Similar reasoning is valid using adapted simplexes instead of cubes in view of Corollary (4.5). According to the definition of the good event  $G_{n,m}^t(\delta; \mathbf{a}, \bar{\mathbf{a}})$  in (6.98), this yields that

$$G_{m,\infty}^1(\delta(3^{-m} \mathcal{X})^\theta; \mathbf{a}, \bar{\mathbf{a}}) \subseteq \{\mathcal{X} \geq 3^m\}.$$

In particular, if  $\delta \in (0, c(d)]$ , we have, by (6.100),

$$n, m \in \mathbb{N}, \quad m \geq n, \quad 3^n \geq \mathcal{X}, \quad u \in \mathcal{A}(\diamond_m) \implies \|s^{1/2} \nabla u\|_{\underline{L}^2(\diamond_n)} \leq C \|s^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \quad (6.127)$$

By (6.86), the finite volume correctors satisfy

$$\begin{aligned} 3^{-\frac{1}{2}m} \left( [\bar{s}^{1/2}(\nabla v(\cdot, \diamond_m, e) - e)]_{\dot{B}_{2,1}^{-1/2}(\diamond_m)} + [\bar{s}^{-1/2}(\mathbf{a} \nabla v(\cdot, \diamond_m, e) - \bar{\mathbf{a}} e)]_{\dot{B}_{2,1}^{-1/2}(\diamond_m)} \right) \\ \leq C \delta (3^{-m} \mathcal{X})^\theta |\bar{s}^{1/2} e| \end{aligned} \quad (6.128)$$

and, by (6.87),

$$3^{-m} \|v(\cdot, \diamond_m, e) - \ell_e\|_{\underline{L}^2(\diamond_m)} \leq C \bar{\lambda}^{-1/2} \delta (3^{-m} \mathcal{X})^\theta |\bar{s}^{1/2} e|. \quad (6.129)$$

For the energy we have

$$\left| \|s^{1/2} \nabla v(\cdot, \diamond_m, e)\|_{\underline{L}^2(\diamond_m)} - |\bar{s}^{1/2} e| \right| \leq C \delta (3^{-m} \mathcal{X})^\theta |\bar{s}^{1/2} e|. \quad (6.130)$$

Furthermore, by (6.9) and (6.10) we obtain, whenever  $3^m \geq \mathcal{X}$  and  $u \in \mathcal{A}(\diamond_m)$ ,

$$3^{-\frac{1}{2}m} [\bar{s}^{-1/2}(\mathbf{a} \nabla u - \bar{\mathbf{a}} \nabla u)]_{\dot{B}_{2,1}^{-1/2}(\diamond_m)} \leq C \delta (3^{-m} \mathcal{X})^\theta \|s^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)} \quad (6.131)$$

and

$$3^{-\frac{1}{2}m} [\bar{s}^{1/2} \nabla u]_{\dot{B}_{2,1}^{-s}(\diamond_m)} \leq C \|s^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \quad (6.132)$$

Caccioppoli estimate (6.16) implies that

$$\|s^{1/2} \nabla u\|_{\underline{L}^2(\diamond_{m-1})} \leq C \bar{\lambda}^{1/2} 3^{-m} \|u\|_{\underline{L}^2(\diamond_m)} + C \delta (3^{-m} \mathcal{X})^\theta \|s^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}. \quad (6.133)$$

Using (6.127), (6.132), (6.133), (6.129) and (6.130) we get, for  $w_m := \nabla v(\cdot, \diamond_{m+1}, e) - \nabla v(\cdot, \diamond_m, e)$ ,

$$\begin{aligned} \|\mathbf{s}^{1/2} \nabla w_m\|_{\underline{L}^2(\diamond_n)} &\leq C \|\mathbf{s}^{1/2} \nabla w_m\|_{\underline{L}^2(\diamond_{m-1})} \\ &\leq C \lambda_0^{1/2} 3^{-m} \|w_m - (w_m)_{\diamond_m}\|_{\underline{L}^2(\diamond_m)} + C \delta (3^{-m} \mathcal{X})^\theta \|\mathbf{s}^{1/2} \nabla w_m\|_{\underline{L}^2(\diamond_m)} \\ &\leq C \delta (3^{-m} \mathcal{X})^\theta |\bar{\mathbf{s}}^{1/2} e|. \end{aligned}$$

In particular,  $\{w_m\}_m$  converges to zero in  $H_{\mathbf{s}}^1(\diamond_n)$  and we set

$$\psi_e := \lim_{m \rightarrow \infty} v(\cdot, \diamond_m, e) \quad \text{and} \quad \phi_e := \psi_e - \ell_e.$$

Then  $\psi_e \in \mathcal{A}(\mathbb{R}^d)$  and  $e \mapsto \psi_e$  is linear. Telescope summation and the triangle inequality gives us

$$\|\mathbf{s}^{1/2} \nabla(\psi_e - v(\cdot, \diamond_m, e))\|_{\underline{L}^2(\diamond_m)} \leq C \theta^{-1} \delta (3^{-m} \mathcal{X})^\theta |\bar{\mathbf{s}}^{1/2} e|.$$

This, together with (6.128), (6.131) and (6.132), gives us (1.37) in Theorem B by taking smaller  $\delta$ , if necessary. Furthermore, by Lemma 6.13 and linearity of  $e \mapsto \psi_e$  we have that there exists a linear map  $e \mapsto P_m[e] \in \mathbb{R}^d$  such that, for every  $n, m \in \mathbb{N}$  with  $m \geq n$  and  $3^n \geq \mathcal{X}$ ,

$$\begin{aligned} \|\mathbf{s}^{1/2} \nabla(\psi_e - v(\cdot, \diamond_m, P_m[e]))\|_{\underline{L}^2(\diamond_n)} &\leq C_\eta 3^{-\eta(m-n)} \|\mathbf{s}^{1/2} \nabla(\psi_e - v(\cdot, \diamond_m, P_m[e]))\|_{\underline{L}^2(\diamond_m)} \\ &\leq C_\eta 3^{-\eta(m-n)} \sum_{n=m}^{\infty} \|\mathbf{s}^{1/2} \nabla w_n\|_{\underline{L}^2(\diamond_m)} \\ &\leq C_\eta 3^{-\eta(m-n)} \theta^{-1} \delta (3^{-m} \mathcal{X})^\theta |\bar{\mathbf{s}}^{1/2} P_m[e]|. \end{aligned}$$

For  $n = m$  we obtain, by the previous two displays and the triangle inequality,

$$\begin{aligned} |\bar{\mathbf{s}}^{1/2}(e - P_m[e])| &\leq C \|\mathbf{s}^{1/2} \nabla(v(\cdot, \diamond_m, e) - v(\cdot, \diamond_m, P_m[e]))\|_{\underline{L}^2(\diamond_m)} \\ &\leq C \theta^{-1} \delta (3^{-m} \mathcal{X})^\theta (|\bar{\mathbf{s}}^{1/2} P_m[e]| + |\bar{\mathbf{s}}^{1/2} e|). \end{aligned}$$

By taking  $\delta$  small enough we find that  $e \mapsto P_m[e]$  is injective. Therefore we get that for every  $e \in \mathbb{R}^d$  there exists  $e' \in \mathbb{R}^d$  such that

$$\|\mathbf{s}^{1/2} \nabla(\psi_{e'} - v(\cdot, \diamond_m, e))\|_{\underline{L}^2(\diamond_k)} \leq C_\eta 3^{-\eta(m-k)} \theta^{-1} \delta (3^{-m} \mathcal{X})^\theta |\bar{\mathbf{s}}^{1/2} e|.$$

Using this, (6.130) and Lemma 6.13 we then deduce that, for every  $m, n \in \mathbb{N}$  with  $m \geq n$  and  $3^n \geq \mathcal{X}$  and for every  $u \in \mathcal{A}(\diamond_m)$ , there exists  $e \in \mathbb{R}^d$  such that

$$\|\mathbf{s}^{1/2} \nabla(u - \psi_e)\|_{\underline{L}^2(\diamond_n)} \leq C_\eta 3^{-\eta(m-k)} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)}.$$

Moreover, by taking  $\frac{1}{2}(1 + \eta)$  instead of  $\eta$ , this immediately implies that

$$\nabla \mathcal{A}_1(\mathbb{R}^d) := \left\{ \nabla u : u \in \mathcal{A}(\mathbb{R}^d), \lim_{m \rightarrow \infty} 3^{-\eta m} \|\mathbf{s}^{1/2} \nabla u\|_{\underline{L}^2(\diamond_m)} = 0 \right\} = \left\{ \nabla \psi_e : e \in \mathbb{R}^d \right\}.$$

The above two displays, together with (6.132) and (2.128), imply both (1.41) and (1.38) in Theorem B.

Finally, the harmonic approximation properties (1.34) and (1.35) are consequences of Propositions 6.7 and 6.9, respectively, together with (6.126). For (1.35) we also use

$$\begin{aligned} \hat{\mathcal{E}}_{s,n}(\diamond_m; \mathbf{a}, \bar{\mathbf{a}}) \|\bar{\mathbf{s}}^{1/2} \nabla u_{\text{hom}}\|_{\underline{L}^2(\diamond_m)} + 3^n \|\bar{\mathbf{s}} \nabla^2 u_{\text{hom}}\|_{L^2(\diamond_m)} \\ \leq C (\hat{\mathcal{E}}_{s,n}(\diamond_m; \mathbf{a}, \bar{\mathbf{a}}) + 3^{-(m-n)}) 3^{-m} \bar{\lambda}^{1/2} \|u_{\text{hom}}\|_{\underline{L}^2(2\diamond_m)} \leq C \delta (3^{-m} \mathcal{X})^\theta 3^{-m} \bar{\lambda}^{1/2} \|u_{\text{hom}}\|_{\underline{L}^2(2\diamond_m)}. \end{aligned}$$

The proof of Theorem B is now complete.



## A. Besov spaces and functional inequalities

We first present a variant of the multiscale Poincaré which has better integrability. Recall the definition of seminorms of Besov spaces  $B_{p,q}^s$  in (2.109) and (2.110), and the weak norm in (2.112).

**Proposition A.1** (Multiscale Sobolev-Poincaré inequality). *Let  $p \in [1, \infty)$  and let*

$$p^* := \begin{cases} \frac{dp}{d-p}, & p < d, \\ \infty, & p \geq d. \end{cases} \quad (\text{A.1})$$

*There exists  $C(p, d) < \infty$  such that, for every  $q \in [p, p^*)$  and  $f \in L^p(\square_m)$ ,*

$$\begin{aligned} \|f\|_{\widehat{W}^{-1,q}(\square_m)} &:= \sup \left\{ \int_{\square_m} fg : 3^{-m}|(g)_{\square_m}| + \|\nabla g\|_{\underline{L}^{q'}(\square_m)} \leq 1 \right\} \\ &\leq C 3^m \sum_{n=-\infty}^m 3^{(n-m)(\frac{d}{q} - \frac{d}{p^*})} \left( \sum_{y \in 3^n \mathbb{Z}^d \cap \square_m} |(f)_{y+\square_n}|^p \right)^{\frac{1}{p}}. \end{aligned} \quad (\text{A.2})$$

*Consequently, for every  $u \in W^{1,p}(\square_m)$  and  $q \in [p, p^*)$ ,*

$$\|u - (u)_{\square_m}\|_{\underline{L}^q(\square_m)} \leq C 3^m \sum_{n=-\infty}^m 3^{(n-m)(\frac{d}{q} - \frac{d}{p^*})} \left( \sum_{y \in 3^n \mathbb{Z}^d \cap \square_m} |(\nabla u)_{y+\square_n}|^p \right)^{\frac{1}{p}}. \quad (\text{A.3})$$

*Proof.* For every  $n \in \mathbb{Z}$  with  $n \leq m$  and  $z \in 3^n \mathbb{Z}^d \cap \square_m$ ,

$$\begin{aligned} \int_{z+\square_k} f \cdot (g - (g)_{z+\square_k}) &= \sum_{y \in 3^{k-1} \mathbb{Z}^d \cap (z+\square_k)} \int_{y+\square_{k-1}} f \cdot (g - (g)_{y+\square_{k-1}}) \\ &\quad + \sum_{y \in 3^{k-1} \mathbb{Z}^d \cap (z+\square_k)} ((g)_{y+\square_{k-1}} - (g)_{z+\square_k}) \cdot (f)_{y+\square_{k-1}}. \end{aligned}$$

Summing over  $z \in 3^k \mathbb{Z}^d \cap \square_m$ , we get

$$\begin{aligned} \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \int_{z+\square_k} f \cdot (g - (g)_{z+\square_k}) &- \sum_{y \in 3^{k-1} \mathbb{Z}^d \cap \square_m} \int_{y+\square_{k-1}} f \cdot (g - (g)_{y+\square_{k-1}}) \\ &= \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \sum_{y \in 3^{k-1} \mathbb{Z}^d \cap (z+\square_k)} ((g)_{y+\square_{k-1}} - (g)_{z+\square_k}) \cdot (f)_{y+\square_{k-1}}. \end{aligned}$$

Summing over  $k \in \mathbb{Z}$  with  $k \leq m$ , we obtain

$$\int_{\square_m} f \cdot (g - (g)_{\square_m}) = \sum_{k=-\infty}^m \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \sum_{y \in 3^{k-1} \mathbb{Z}^d \cap (z+\square_k)} ((g)_{y+\square_{k-1}} - (g)_{z+\square_k}) \cdot (f)_{y+\square_{k-1}}.$$

By applying the Hölder inequality we get

$$\begin{aligned} &\left| \int_{\square_m} f \cdot (g - (g)_{\square_m}) \right| \\ &\leq \sum_{k=-\infty}^m \left( \sum_{y \in 3^{k-1} \mathbb{Z}^d \cap \square_m} |(f)_{y+\square_{k-1}}|^p \right)^{\frac{1}{p}} \left( \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \sum_{y \in 3^{k-1} \mathbb{Z}^d \cap (z+\square_k)} |(g)_{y+\square_{k-1}} - (g)_{z+\square_k}|^{p'} \right)^{\frac{1}{p'}}. \end{aligned}$$

By the Sobolev inequality, for every  $q' \in ((p')_*, p']$ , if we set  $\beta := d(p')_*^{-1} - d(q')^{-1} > 0$ , we have

$$\begin{aligned}
& \left( \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \sum_{y \in 3^{k-1} \mathbb{Z}^d \cap (z + \square_k)} |(g)_{y + \square_{k-1}} - (g)_{z + \square_k}|^{p'} \right)^{\frac{1}{p'}} \\
& \leq \left( 3^d \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \int_{z + \square_k} |g - (g)_{z + \square_k}|^{p'} \right)^{\frac{1}{p'}} \\
& \leq C 3^k \left( \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \left( \int_{z + \square_k} |\nabla g|^{(p')_*} \right)^{\frac{p'}{(p')_*}} \right)^{\frac{1}{p'}} \\
& \leq C 3^k \left( \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \left( \int_{z + \square_k} |\nabla g|^{q'} \right)^{\frac{p'}{q'}} \right)^{\frac{1}{p'}} \\
& \leq C 3^k \left( \frac{|\square_m|}{|\square_k|} \right)^{\frac{1}{q'} - \frac{1}{p'}} \left( \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \int_{z + \square_k} |\nabla g|^{q'} \right)^{\frac{1}{q'}} = 3^{m - \beta(m-k)} \left( \int_{\square_m} |\nabla g|^{q'} \right)^{\frac{1}{q'}}.
\end{aligned}$$

Using also that  $\left| \oint_{\square_m} f(g)_{\square_m} \right| = |(f)_{\square_m}| |(g)_{\square_m}|$ , we obtain

$$\left| \oint_{\square_m} fg \right| \leq C \left( 3^{-m} |(g)_{\square_m}| + \|\nabla g\|_{\underline{L}^{q'}(\square_m)} \right) 3^m \sum_{k=-\infty}^m 3^{\beta(k-m)} \left( \sum_{y \in 3^k \mathbb{Z}^d \cap \square_m} |(f)_{y + \square_k}|^p \right)^{\frac{1}{p}}.$$

By duality, noting that  $(p')_* = (p^*)'$ , we obtain (A.2).

*Step 2.* Proof of (A.3). Assume that  $(u)_{\square_m} = 0$ . Let  $w \in W^{2,q'}(\square_m)$  solve  $-\Delta w = |u|^{q-2}u - (|u|^{q-2}u)_{\square_m}$  in  $\square_m$  with zero Neumann boundary data. Then, by the Calderón-Zygmund estimate and the triangle inequality,

$$\|\nabla w\|_{\underline{W}^{1,q'}(\square_m)} \leq C \| |u|^{q-2}u - (|u|^{q-2}u)_{\square_m} \|_{\underline{L}^{q'}(\square_m)} \leq C \|u\|_{\underline{L}^q(\square_m)}^{q-1}.$$

Using this and testing the equation of  $w$  with  $u$ , we deduce by  $(u)_{\square_m} = 0$  that

$$\begin{aligned}
\|u\|_{\underline{L}^q(\square_m)}^q &= \oint_{\square_m} \nabla w \cdot \nabla u \\
&\leq \|\nabla w\|_{\underline{W}^{1,q'}(\square_m)} \|\nabla u\|_{\widehat{W}^{-1,q}(\square_m)} \\
&\leq C 3^m \|u\|_{\underline{L}^q(\square_m)}^{q-1} \sum_{n=-\infty}^m 3^{(n-m)(\frac{d}{q} - \frac{d}{p^*})} \left( \sum_{y \in 3^n \mathbb{Z}^d \cap \square_m} |(\nabla u)_{y + \square_n}|^p \right)^{\frac{1}{p}}.
\end{aligned}$$

This concludes the proof.  $\square$

The following lemma gives an estimate for duality pairing between  $\underline{B}_{q',p'}^s(\square_m)$  and  $\underline{B}_{q,p}^{-s}(\square_m)$ .

**Lemma A.2.** *Let  $q \in [1, \infty]$ . Let also  $p \in (1, \infty)$  and  $s \in (0, 1)$ , or  $p = 1$  and  $s \in (0, 1]$ . Then, with  $p' = \frac{p}{p-1}$  for  $p < \infty$  and with  $p' = \infty$  for  $p = 1$ , and similarly for  $q'$ , we have that*

$$[f]_{\widehat{\underline{B}}_{p,q}^{-s}(\square_m)} \leq 3^{d+s} \left( \sum_{k=-\infty}^m \left( 3^{spk} \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} |(f)_{z + \square_k}|^p \right)^{q/p} \right)^{1/q}. \quad (\text{A.4})$$

*Proof.* We follow the proof of Proposition A.1 until the identity

$$\int_{\square_m} f \cdot (g - (g)_{\square_m}) = \sum_{k=-\infty}^m \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \sum_{y \in 3^{k-1} \mathbb{Z}^d \cap (z + \square_k)} ((g)_{y + \square_{k-1}} - (g)_{z + \square_k}) \cdot (f)_{y + \square_{k-1}}.$$

After this we deviate from the proof and estimate, for  $p \in (1, \infty)$  using Hölder's inequality, as

$$\begin{aligned} & \sum_{k=-\infty}^m \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \sum_{y \in 3^{k-1} \mathbb{Z}^d \cap (z + \square_k)} |((g)_{y + \square_{k-1}} - (g)_{z + \square_k}) \cdot (f)_{y + \square_{k-1}}| \\ & \leq 3^d \sum_{k=-\infty}^m \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \|g - (g)_{z + \square_k}\|_{\underline{L}^1(z + \square_k)} \sum_{y \in 3^{k-1} \mathbb{Z}^d \cap (z + \square_k)} |(f)_{y + \square_{k-1}}| \\ & \leq 3^d \sum_{k=-\infty}^m \left( \sum_{z \in 3^k \mathbb{Z}^d \cap \square_m} \|g - (g)_{z + \square_k}\|_{\underline{L}^{p'}(z + \square_k)}^{p'} \right)^{1/p'} \left( \sum_{z \in 3^{k-1} \mathbb{Z}^d \cap \square_m} |(f)_{z + \square_{k-1}}|^p \right)^{1/p}. \end{aligned}$$

The first average  $|(g)_{\square_m}| |(f)_{\square_m}|$  is then added to the sum trivially as the term  $k = m$ , and (A.4) follows by Hölder's inequality. The case  $p = 1$  is similar. The proof is complete.  $\square$

The next lemma provides a Poincaré type inequality between Besov spaces.

**Lemma A.3.** *There exists  $C(d) < \infty$  such that, for every  $n \in \mathbb{N}$ ,  $s \in [0, 1)$  and  $u \in B_{2,\infty}^s(\square_n)$ , we have that*

$$\|u - (u)_{\square_n}\|_{\underline{B}_{2,\infty}^s(\square_n)} \leq C[\nabla u]_{\dot{\underline{B}}_{2,1}^{s-1}(\square_n)}. \quad (\text{A.5})$$

Moreover, if  $\varphi \in C_c^\infty(\square_n)$  satisfies  $3^n \|\nabla \varphi\|_{L^\infty(\square_n)} + 3^{2n} \|\nabla^2 \varphi\|_{L^\infty(\square_n)} \leq 1$ , then

$$\|(u - (u)_{\square_n}) \nabla \varphi\|_{\underline{B}_{2,\infty}^s(\square_n)} \leq C 3^{-n} [\nabla u]_{\dot{\underline{B}}_{2,1}^{s-1}(\square_n)}. \quad (\text{A.6})$$

*Proof.* Fix  $s \in [0, 1)$  and  $n \in \mathbb{N}$ . Without loss of generality, assume that  $(u)_{\square_n} = 0$ . Throughout the proof we denote the lattice appearing in (2.109) by  $\mathcal{Z}_k := \{z \in 3^{k-1} \mathbb{Z}^d : z + \square_k \subseteq \square_n\}$  for each  $k \in \mathbb{Z}$  with  $k \leq n$ . Fix also  $\phi \in C_c^\infty(\square_n)$  satisfying  $3^n \|\nabla \phi\|_{L^\infty(\square_n)} + 3^{2n} \|\nabla^2 \phi\|_{L^\infty(\square_n)} \leq 1$ .

*Step 1.* The proof of (A.5). Apply (A.3) and Hölder's inequality to get that

$$\|u - (u)_{z + \square_k}\|_{\underline{L}^2(z + \square_k)} \leq C \sum_{j=-\infty}^k 3^j \left( \sum_{z' \in z + 3^j \mathbb{Z}^d \cap \square_k} |(\nabla u)_{z' + \square_j}|^2 \right)^{1/2}.$$

Therefore, we obtain, again by Hölder's inequality, that

$$\begin{aligned} & \sup_{k \in (-\infty, n] \cap \mathbb{Z}} 3^{-sk} \left( \sum_{z \in \mathcal{Z}_k} \|u - (u)_{z + \square_k}\|_{\underline{L}^2(z + \square_k)}^2 \right)^{1/2} \\ & \leq \sup_{k \in (-\infty, n] \cap \mathbb{Z}} 3^{-sk} \sum_{z \in \mathcal{Z}_k} \|u - (u)_{z + \square_k}\|_{\underline{L}^2(z + \square_k)} \\ & \leq \sup_{k \in (-\infty, n] \cap \mathbb{Z}} 3^{-sk} \sum_{z \in \mathcal{Z}_k} \sum_{j=-\infty}^k 3^j \left( \sum_{z' \in z + 3^j \mathbb{Z}^d \cap \square_k} |(\nabla u)_{z' + \square_j}|^2 \right)^{1/2} \\ & \leq C \sup_{k \in (-\infty, n] \cap \mathbb{Z}} \sum_{j=-\infty}^k 3^{(1-s)j} \left( \sum_{z' \in z + 3^j \mathbb{Z}^d \cap \square_n} |(\nabla u)_{z' + \square_j}|^2 \right)^{1/2}. \end{aligned}$$

This yields (A.5).

*Step 2.* The proof of (A.6). We first show that there exists a constant  $C(d) < \infty$  such that, for every  $v \in B_{2,\infty}^s(\square_n)$ ,

$$[v \nabla \varphi]_{\underline{B}_{2,\infty}^s(\square_n)} \leq C 3^{(1-s)n} \|\nabla^2 \varphi\|_{L^\infty(\square_n)} \|v\|_{\underline{L}^2(\square_n)} + C \|\nabla \varphi\|_{L^\infty(\square_n)} [v]_{\underline{B}_{2,\infty}^s(\square_n)}. \quad (\text{A.7})$$

Recall the definition of the seminorm from (2.109):

$$[v \nabla \varphi]_{\underline{B}_{2,\infty}^s(\square_n)} = \sup_{k \in (-\infty, n] \cap \mathbb{Z}} 3^{-sk} \left( \sum_{z \in \mathcal{Z}_k} \|v \nabla \varphi - (v \nabla \varphi)_{z+\square_k}\|_{\underline{L}^2(z+\square_k)}^2 \right)^{1/2}.$$

By the triangle inequality,

$$\begin{aligned} & \left( \sum_{z \in \mathcal{Z}_k} \|v \nabla \varphi - (v \nabla \varphi)_{z+\square_k}\|_{\underline{L}^2(z+\square_k)}^2 \right)^{1/2} \\ & \leq 2 \left( \sum_{z \in \mathcal{Z}_k} \left( \|\nabla \varphi - (\nabla \varphi)_{z+\square_k}\|_{\underline{L}^2(z+\square_k)}^2 |(v)_{z+\square_k}|^2 + \|(v - (v)_{z+\square_k}) \nabla \varphi\|_{\underline{L}^2(z+\square_k)}^2 \right) \right)^{1/2} \\ & \leq C 3^k \|\nabla^2 \varphi\|_{L^\infty(\square_n)} \|v\|_{\underline{L}^2(\square_n)} + C \|\nabla \varphi\|_{L^\infty(\square_n)} \left( \sum_{z \in \mathcal{Z}_k} \|v - (v)_{z+\square_k}\|_{\underline{L}^2(z+\square_k)}^2 \right)^{1/2}. \end{aligned}$$

Thus (A.7) follows. Since  $|(u \nabla \varphi)_{\square_n}| \leq \|\nabla \varphi\|_{L^\infty(\square_n)} \|u\|_{\underline{L}^2(\square_n)}$ , we may apply (A.7) to obtain

$$\|u \nabla \varphi\|_{\underline{B}_{2,\infty}^s(\square_n)} \leq C 3^{-sn} (3^n \|\nabla^2 \varphi\|_{L^\infty(\square_n)} + \|\nabla \varphi\|_{L^\infty(\square_n)}) \|u\|_{\underline{L}^2(\square_n)} + C \|\nabla \varphi\|_{L^\infty(\square_n)} [u]_{\underline{B}_{2,\infty}^s(\square_n)}. \quad (\text{A.8})$$

An application of (A.3) and (A.5) then concludes the proof.  $\square$

## B. Geometric means for positive matrices

Throughout, we denote the set of  $N$ -by- $M$  matrices with real entries by  $\mathbb{R}^{N \times M}$ . If  $A \in \mathbb{R}^{N \times M}$ , then the transpose of  $A$  is denoted by  $A^t$ . We let  $\mathbb{R}_{\text{sym}}^{N \times N} := \{A \in \mathbb{R}^{N \times N} : A = A^t\}$  be the set of symmetric  $N$ -by- $N$  matrices, and  $\mathbb{R}_{\text{skew}}^{N \times N} := \{A \in \mathbb{R}^{N \times N} : A = -A^t\}$  the set of anti-symmetric matrices. We use the Loewner partial order on  $\mathbb{R}_{\text{sym}}^{N \times N}$ ; that is, if  $A, B \in \mathbb{R}_{\text{sym}}^{N \times N}$  then we write  $A \leq B$  if  $B - A$  has nonnegative eigenvalues.

We use the spectral norm for matrices. For each  $A \in \mathbb{R}^{N \times N}$ , we write

$$|A| := \max_{e \in \mathbb{R}^N, |e|=1} |Ae|.$$

Note that  $|A|$  is the largest eigenvalue of  $(A^t A)^{1/2}$ . It follows that, for any pair of square matrices  $A$  and  $B$ , we have that  $|A| = |A^t|$ . The spectral norm is also submultiplicative: for all square matrices  $A$  and  $B$ ,

$$|AB| \leq |A| |B|.$$

Recall that if  $A, B > 0$  are positive real numbers, then the minimum of the map

$$x \mapsto \frac{1}{2} x^{-1/2} A x^{-1/2} + \frac{1}{2} x^{1/2} B^{-1} x^{1/2}$$

is attained uniquely at  $x = A\#B$ , where  $A\#B := (AB)^{1/2}$  is the geometric mean of  $A$  and  $B$ , and the minimum is equal to  $A\#B^{-1}$ . It turns out that this fact can be generalized to positive definite matrices.

There are two different notions of geometric mean for positive definite matrices. The first one, introduced by Ando [And78], is called the *metric geometric mean*. It is defined for any pair of positive definite matrices  $A$  and  $B$  by

$$A\#B = A^{1/2}(A^{-1/2}BA^{-1/2})^{1/2}A^{1/2}.$$

The matrix  $A\#B$  is the unique positive definite matrix solution  $X$  of the equation

$$XA^{-1}X = B. \tag{B.1}$$

We see from this characterization that the metric geometric mean is symmetric in  $A$  and  $B$ , that is,  $A\#B = B\#A$ .

A second notion of geometric mean for positive definite matrices was introduced later by Fiedler and Pták [FP97], is the *spectral geometric mean*  $A\sharp B$  of two matrices  $A$  and  $B$  is defined by

$$A\sharp B := (A^{-1}\#B)^{1/2}A(A^{-1}\#B)^{1/2}.$$

It is also characterized by the following identity that relates it to the metric geometric mean:

$$A^{-1}\#(A\sharp B) = B\#(A\sharp B)^{-1}.$$

It gets its name from the fact that  $(A\sharp B)^2$  is positively similar to  $AB$ . In fact, there exists a positive definite matrix  $C$  such that

$$A\sharp B = CAC = C^{-1}BC^{-1},$$

and this property characterizes  $A\sharp B$ . It also follows from this characterization that the spectral geometric means is also symmetric in  $A$  and  $B$ , that is,  $A\sharp B = B\sharp A$ . Since  $(A\sharp B)^2$  is similar to  $AB$ , the eigenvalues of  $A\sharp B$  are the square roots of those of  $AB$  or  $BA$  or  $A^{1/2}BA^{1/2}$  or  $B^{1/2}AB^{1/2}$ . This property gives it its name.

The largest eigenvalue of  $A\sharp B$  is larger than the largest eigenvalue of  $A\#B$ , while this relation is reversed for the smallest eigenvalue. In particular,  $|A\#B| \leq |A\sharp B|$ . It turns out that  $A\#B = A\sharp B$  if and only if  $A$  and  $B$  commute. If they do not commute, there is no relation between the two in the Loewner partial order. All of the facts asserted above can be found in [FP97].

It turns out that the map

$$X \mapsto \frac{1}{2}X^{-1/2}AX^{-1/2} + \frac{1}{2}X^{1/2}B^{-1}X^{1/2}$$

is minimized by  $X = A\#B$  and the minimum is equal to  $A\sharp B^{-1}$ .

## C. Orlicz quasi-norms and concentration inequalities

**C.1. The  $\mathcal{O}_\Psi$  notation for weak Orlicz quasi-norms.** If  $\Psi : \mathbb{R}_+ \rightarrow [1, \infty)$  is an increasing function satisfying the mild growth condition

$$\lim_{t \rightarrow \infty} \frac{1}{t} \Psi(t) = +\infty,$$

and  $X$  is a random variable, then we use the notation “ $X \leq \mathcal{O}_\Psi(A)$ ” for  $A \geq 0$  as a shorthand for the statement

$$\mathbb{P}[X > tA] \leq \frac{1}{\Psi(t)}, \quad \forall t \in [1, \infty).$$

For example, we could write (1.16) as  $S \leq \mathcal{O}_{\Psi_S}(1)$  and, similarly, (1.31) can be written as

$$\left| \sum_{z \in 3^n \mathbb{Z}^d \cap \square_m} X_z \right| \leq \mathcal{O}_\Psi(3^{-\frac{d}{2}(m-n)}).$$

Thus the notation “ $X \leq \mathcal{O}_\Psi(A)$ ” is an alternative way of writing that a weak Orlicz quasi-norm of  $X$  is bounded by  $A$ . We note however that we do *not* require  $\Psi$  is be convex here. We also write  $X = \mathcal{O}_\Psi(A)$  to mean that  $|X| \leq \mathcal{O}_\Psi(A)$ .

If  $\Psi_1$  and  $\Psi_2$  are two such functions, then we write  $X \leq \mathcal{O}_{\Psi_1}(A_i)$  to mean that  $X$  can be written as the sum  $X = Y_1 + Y_2$  of two random variables  $Y_1$  and  $Y_2$  satisfying  $Y_i = \mathcal{O}_{\Psi_i}(A_i)$  for  $i \in \{1, 2\}$ .

For this  $\mathcal{O}_\Psi$  notation to be useful, we need to have other properties such as a (generalized) triangle inequality for infinite sums, and sufficient growth of  $\Psi$  that  $X = \mathcal{O}_\Psi(A)$  implies bounds on finite moments of  $X$ . In the next lemma, we prove such properties under the growth condition (1.29) used in our assumptions.

**Lemma C.1.** *Suppose that  $\Psi : \mathbb{R}_+ \rightarrow [1, \infty)$  is an increasing function and  $K_\Psi \in [2, \infty)$  such that*

$$t\Psi(t) \leq \Psi(K_\Psi t), \quad \forall t \in [1, \infty). \quad (\text{C.1})$$

*Then we have the following:*

- For every  $p \in [1, \infty)$ ,

$$\frac{t^p}{\Psi(t)} \leq \frac{1}{\Psi(K_\Psi^{-4[p]} t)}, \quad \forall t \in [K_\Psi^{4[p]}, \infty), \quad (\text{C.2})$$

- $\Psi$  satisfies the following minimal growth bound:

$$\Psi(t) \geq \exp\left(\frac{\log^2 t}{9 \log K_\Psi}\right), \quad \forall t \in [K_\Psi^2, \infty). \quad (\text{C.3})$$

- For every  $p \in [2, \infty)$ ,

$$s^p \leq K_\Psi^{3p^2} \frac{\Psi(ts)}{\Psi(t)}, \quad \forall t, s \in [1, \infty). \quad (\text{C.4})$$

- For any random variable  $X$  and  $a \in (0, \infty)$  and  $p \in [1, \infty)$ ,

$$X = \mathcal{O}_\Psi(a) \implies \mathbb{E}[X^p] \leq a^p \left(1 + 2p K_\Psi^{\lfloor \frac{1}{2} p(p+1) \rfloor} (1 + \log K_\Psi)\right). \quad (\text{C.5})$$

- For any sequence  $\{X_k\}_{k \in \mathbb{N}}$  of random variables,

$$X_k \leq \mathcal{O}_\Psi(a_k) \implies \sum_{k \in \mathbb{N}} X_k \leq \mathcal{O}_\Psi\left(4K_\Psi^7 \sum_{k \in \mathbb{N}} a_k\right). \quad (\text{C.6})$$

*Proof. Step 1.* We begin with the observation that (1.29) allows us to absorb powers of  $t$  in front of  $\Psi$  by a dilation of  $\Psi$ . By induction, it implies that, for every power  $k \in \mathbb{N}$ ,

$$t^k \Psi(t) \leq K_\Psi^{-\frac{1}{2}k(k-1)} \Psi(K_\Psi^k t), \quad \forall t \in [1, \infty). \quad (\text{C.7})$$

In particular, for every  $p \in [1, \infty)$ ,

$$t^p \Psi(t) \leq \Psi(K_\Psi^{[p]} t), \quad \forall t \in [1, \infty). \quad (\text{C.8})$$

This implies that, for every  $p \in [1, \infty)$ ,

$$\frac{t^p}{\Psi(t)} \leq (K_\Psi^{2[2p]} t^{-1})^p \frac{1}{\Psi(K_\Psi^{[2p]} t)}, \quad \forall t \geq K_\Psi^{[2p]}.$$

If we restrict to  $t \geq K_\Psi^{4[p]}$  then we can ignore the first factor. This yields (C.2).

*Step 2.* The proof of (C.3). We observe that (C.7) implies, for every  $p \in \mathbb{N} \cap [1, \infty)$  and  $t \in [K_\Psi^p, K_\Psi^{p+1}]$ ,

$$\Psi(t) \geq \frac{\Psi(t)}{\Psi(1)} \geq K_\Psi^{\frac{1}{2}p(p-1)} \geq t^{\frac{p(p-1)}{2(p+1)}} \geq \exp\left(\frac{p(p-1)}{2(p+1)} \log t\right) \geq \exp\left(\frac{p(p-1)}{2(p+1)^2} \frac{\log^2 t}{\log K_\Psi}\right). \quad (\text{C.9})$$

Specializing to  $p \geq 2$  yields (C.3).

*Step 3.* The proof of (C.4). As in (C.9), we use (C.7) to find that, for every  $p \in \mathbb{N}$ , we have

$$\frac{\Psi(ts)}{\Psi(t)} \geq s^{\frac{p(p-1)}{2(p+1)}}, \quad \forall t \in [1, \infty), \quad s \in [K_\Psi^p, K_\Psi^{p+1}].$$

The upper bound restriction on  $s$  can clearly be removed, and so we obtain

$$\frac{\Psi(ts)}{\Psi(t)} \geq s^{\frac{p(p-1)}{2(p+1)}}, \quad \forall t \in [1, \infty), \quad s \in [K_\Psi^p, \infty),$$

and this implies

$$\frac{\Psi(ts)}{\Psi(t)} \geq K_\Psi^{-\frac{p^2(p-1)}{2(p+1)}} s^{\frac{p(p-1)}{2(p+1)}}, \quad \forall t \in [1, \infty), \quad s \in [1, \infty),$$

Restricting to  $p \geq 5$ , we get

$$\frac{\Psi(ts)}{\Psi(t)} \geq K_\Psi^{-p^2/3} s^{p/3}, \quad \forall t \in [1, \infty), \quad s \in [1, \infty),$$

This implies (C.4).

*Step 4.* We prove (C.5). For any random variable  $X$  satisfying  $X = \mathcal{O}_\Psi(a)$ , we have

$$a^{-n} \mathbb{E}[|X|^n] = n \int_0^\infty t^{n-1} \mathbb{P}[|X| > at] dt \leq 1 + n \int_1^\infty \frac{t^{n-1}}{\Psi(t)} dt.$$

Using (C.7), we find that, for every  $j \in \mathbb{N}$  and  $t \in [K_\Psi^j, \infty)$ ,

$$\Psi(t) \geq K_\Psi^{-\frac{1}{2}j(j+1)} t^j \Psi(K_\Psi^{-j} t) \geq K_\Psi^{-\frac{1}{2}j(j+1)} t^j.$$

Using this, we obtain

$$\begin{aligned}
\int_1^\infty \frac{t^{n-1}}{\Psi(t)} dt &\leq \sum_{j=0}^n \int_{K_\Psi^j}^{K_\Psi^{j+1}} \frac{t^{n-1}}{\Psi(t)} dt + \int_{K_\Psi^{n+1}}^\infty \frac{t^{n-1}}{\Psi(t)} dt \\
&\leq \sum_{j=0}^n K_\Psi^{\frac{1}{2}j(j+1)} \int_{K_\Psi^j}^{K_\Psi^{j+1}} t^{n-j-1} dt + n K_\Psi^{\frac{1}{2}(n+2)(n+1)} \int_{K_\Psi^{n+1}}^\infty t^{-2} dt \\
&\leq \sum_{j=0}^{n-1} \frac{1}{n-j} K_\Psi^{\frac{1}{2}j(j+1)+(n-j)(j+1)} + (1 + \log K_\Psi) K_\Psi^{\frac{1}{2}n(n+1)} \\
&\leq 2(1 + \log K_\Psi) K_\Psi^{\frac{1}{2}n(n+1)}.
\end{aligned}$$

This yields, for every  $n \in \mathbb{N}$ ,

$$\mathbb{E}[|X|^n] \leq a^n \left( 1 + 2n K_\Psi^{\frac{1}{2}n(n+1)} (1 + \log K_\Psi) \right),$$

completing the proof of (C.5).

*Step 5.* We show that (C.4) implies the generalized triangle inequality (C.6). We assume only that  $\Psi$  satisfies, for some  $p, C_0 \in (1, \infty)$ ,

$$s^p \leq C_0 \frac{\Psi(ts)}{\Psi(t)}, \quad \forall t, s \in [1, \infty). \quad (\text{C.10})$$

We argue that this condition (C.10) implies the following generalized triangle inequality: there exists a constant  $C(p, C_0) < \infty$  (given explicitly below in (C.12)), such that, if  $\{X_k\}_{k \in \mathbb{N}}$  is any sequence of random variables, then

$$X_k \leq \mathcal{O}_\Psi(a_k) \implies \sum_{k \in \mathbb{N}} X_k \leq \mathcal{O}_\Psi\left(C \sum_{k \in \mathbb{N}} a_k\right). \quad (\text{C.11})$$

To see this, we set  $X := \sum_k X_k$ ,  $a := \sum_k a_k$ , fix  $t > 0$  and compute

$$\mathbb{P}[X > t] \leq \mathbb{P}\left[\sum_{k \in \mathbb{N}} X_k \mathbf{1}_{\{X_k > ta_k/2a\}} > \frac{1}{2}t\right] \leq \frac{2}{t} \sum_{k \in \mathbb{N}} \mathbb{E}[X_k \mathbf{1}_{\{X_k > ta_k/2a\}}].$$

We then observe that

$$\begin{aligned}
\mathbb{E}[X_k \mathbf{1}_{\{X_k > ta_k/2a\}}] &= \int_0^\infty \mathbb{P}[X_k \mathbf{1}_{\{X_k > ta_k/2a\}} > s] ds \\
&\leq \frac{ta_k}{2a} \mathbb{P}\left[X_k > \frac{ta_k}{2a}\right] + \int_{ta_k/2s}^\infty \mathbb{P}[X_k > s] ds \\
&\leq \frac{ta_k}{2a} \frac{1}{\Psi(t/2a)} + \int_{ta_k/2a}^\infty \frac{1}{\Psi(s/a_k)} ds \\
&\leq \frac{1}{\Psi(t/2a)} \left( \frac{ta_k}{2a} + \int_{ta_k/2a}^\infty \frac{\Psi(t/2a)}{\Psi(s/a_k)} ds \right) \\
&\leq \frac{1}{\Psi(t/2a)} \left( \frac{ta_k}{2a} + C_0 \int_{ta_k/2a}^\infty \left( \frac{2as}{a_k t} \right)^{-p} ds \right) = \frac{1}{\Psi(t/2a)} \left( \frac{ta_k}{2a} \right) \left( 1 + \frac{C_0}{p-1} \right).
\end{aligned}$$



Here we used (C.10) in the last line. Inserting this into the previous display after summing it over  $k \in \mathbb{N}$  gives

$$\mathbb{P}[X > t] \leq \frac{1 + C_0(p-1)^{-1}}{\Psi(t/2a)}.$$

Using (C.10) again, we can bound the right side by  $\Psi(t/Ca)^{-1}$  with

$$C(p, C_0) := 2(C_0 + C_0^2(p-1)^{-1})^{1/p}, \quad (\text{C.12})$$

which then implies  $X = \mathcal{O}_\Psi(Ca)$ , as claimed. If we have (C.4), then we may take  $p = 2$  and  $C_0 = K_\Psi^7$  and we find that  $C(p, C_0) \leq 4K_\Psi^7$ , as claimed in (C.6). This completes the proof of the lemma.  $\square$

We may also use (C.10) in the equivalent form

$$\frac{s}{\Psi(t)} \leq \frac{1}{\Psi(t(sC_0)^{-1/p})}, \quad \forall s \in [C_0^{-1}, \infty), \quad t \in [(sC_0)^{1/p}, \infty). \quad (\text{C.13})$$

**C.2. Examples of functions satisfying the growth condition.** We introduce, for each  $\sigma \in (0, \infty)$ , the function

$$\Gamma_\sigma(t) := \exp(t^\sigma). \quad (\text{C.14})$$

If  $\sigma \geq 1$ , then the function  $\Gamma_\sigma$  is nonnegative, increasing on  $\mathbb{R}_+$ , convex and satisfies  $t\Gamma_\sigma(t) \leq \Gamma_\sigma(2t)$  for every  $t \geq 1$  and  $\Gamma_\sigma$  satisfies the generalized triangle inequality (C.6) with constant  $C = 1$ . In the case  $\sigma \in (0, 1)$ ,  $\Gamma_\sigma$  is convex on the interval  $((\frac{1-\sigma}{\sigma})^{1/\sigma}, \infty)$ , the growth condition is satisfied for

$$K_{\Gamma_\sigma} = \left(\frac{\sigma+1}{\sigma}\right)^{1/\sigma}. \quad (\text{C.15})$$

and the generalized triangle inequality is valid with constant  $C = 2\sigma^{-1}$ . For every  $\sigma_1, \sigma_2 \in (0, \infty)$  and random variables  $X_1$  and  $X_2$ ,

$$X_1 \leq \mathcal{O}_{\Gamma_{\sigma_1}}(A_1) \quad \text{and} \quad X_2 \leq \mathcal{O}_{\Gamma_{\sigma_2}}(A_2) \implies X_1 X_2 \leq \mathcal{O}_{\Gamma_{\frac{\sigma_1 \sigma_2}{\sigma_1 + \sigma_2}}}(A_1 A_2). \quad (\text{C.16})$$

For any  $\sigma, p, K \in (0, \infty)$  and random variable  $X$ ,

$$X \leq \Gamma_\sigma(K) \iff X^p \leq \Gamma_{\sigma/p}(K^p).$$

A normal random variable  $X$  with zero mean and variance  $\gamma^2 > 0$  satisfies

$$X = \mathcal{O}_{\Gamma_2}(\gamma).$$

If  $\sigma, A > 0$  and  $X_1, \dots, X_N$  is a sequence of random variables satisfying  $X_i = \mathcal{O}_{\Gamma_\sigma}(A)$  with  $N \geq 2$ , then

$$\max_{1 \leq i \leq N} X_i = \mathcal{O}_{\Gamma_\sigma}((3 \log N)^{1/\sigma} A). \quad (\text{C.17})$$

To see this, we observe that, for every  $t \geq 1$  we use a union bound to estimate

$$\begin{aligned} \mathbb{P}\left[\max_{1 \leq i \leq N} X_i > A(3 \log N)^{1/\sigma} t\right] &\leq \sum_{i=1}^N \mathbb{P}[X_i > A(3 \log N)^{1/\sigma} t] \\ &\leq N \exp(-3t^\sigma \log N) \leq \exp(-t^\sigma (3 \log N - \log N)) \leq \exp(-t^\sigma). \end{aligned}$$

The indicator function  $\mathbf{1}_E$  of an event  $E$  with  $0 < \mathbb{P}[E] < 1$  satisfies, for every  $\sigma \in (0, \infty)$ ,

$$\mathbf{1}_E \leq \mathcal{O}_{\Gamma_\sigma}(|\log \mathbb{P}[E]|^{-1/\sigma}). \quad (\text{C.18})$$

This is immediate from the definitions.

We have seen in (C.3) that any admissible  $\Psi$  grows at least like  $t \mapsto c \exp(c \log^2 t)$ . Conversely, this function satisfies the growth condition, and it is an important example since it characterizes the integrability of log-normal random variables. For instance,

$$\Psi_1(t) = \exp(\log^2(1+t))$$

is nonnegative, increasing on  $[0, \infty)$  and satisfies (1.29) with  $K_{\Psi_1} = 10$ . More generally, for each  $\sigma \in (0, \infty)$ , we define

$$\Psi_\sigma(t) := \exp\left(\frac{1}{\sigma^2} \log^2(1 + \sigma t)\right), \quad t \in [0, \infty). \quad (\text{C.19})$$

This class of functions captures the stochastic integrability of log-normal random variables in the sense that, for every random variable  $X$ ,

$$X \leq \mathcal{O}_{\Gamma_2}(\sigma) \iff \exp(X) - 1 \leq \mathcal{O}_{\Psi_\sigma}(\sigma). \quad (\text{C.20})$$

If  $\sigma \leq 1$ , then

$$\frac{1}{\sigma^2} \log^2(1 + \sigma t) \leq \log^2(1 + t), \quad \forall t \in [0, \infty)$$

by the concavity of the logarithm, and so  $Y \leq \mathcal{O}_{\Psi_\sigma}(a)$  implies that  $Y \leq \mathcal{O}_{\Psi_1}(a)$ . For this reason we will generally use  $\Psi_\sigma$  only for  $\sigma \geq 1$ . For all such  $\sigma$ , the function  $\Psi_\sigma$  is admissible since

$$\text{for every } \sigma \in [1, \infty), \quad \Psi_\sigma \text{ satisfies (1.29) with constant } K = K_{\Psi_\sigma} := 2 \exp(2\sigma^2). \quad (\text{C.21})$$

To prove (C.21), observe that, for every  $\sigma, K \geq 1$ ,

$$\begin{aligned} t\Psi_\sigma(t) \leq \Psi_\sigma(Kt), \quad \forall t \in [1, \infty) &\iff \sigma^2 \log t \leq \log^2(1 + K\sigma t) - \log^2(1 + \sigma t), \quad \forall t \in [1, \infty) \\ &\iff \sigma^2 \log t \leq (\log(1 + K\sigma t)) \log\left(\frac{1 + K\sigma t}{1 + \sigma t}\right), \quad \forall t \in [1, \infty), \end{aligned}$$

where in the last line we used that  $A^2 - B^2 = (A + B)(A - B) \geq A(A - B)$  for all  $A \geq B > 0$ . Moreover, the last statement on the right is valid with the choice of  $K = 2 \exp(\sigma^2)$  since it makes the second logarithm factor on the right side larger than  $\sigma^2$ , while the first is clearly larger than  $\log t$ . This completes the proof of (C.21).

**C.3. Concentration inequalities with respect to Orlicz quasi-norms.** We first present a simple concentration inequality for random variables with at least exponential integrability.

**Lemma C.2** (Concentration for exponential random variables). *Let  $\sigma \in [1, 2]$  and  $m \in \mathbb{N}$ . Suppose that  $X_1, \dots, X_m$  is a sequence of independent random variables satisfying*

$$X_k = \mathcal{O}_{\Gamma_\sigma}(1) \quad \text{and} \quad \mathbb{E}[X_k] = 0, \quad \forall k \in \{1, \dots, m\}. \quad (\text{C.22})$$

*Then, for every  $t \geq 1$ ,*

$$\mathbb{P}\left[\sum_{k=1}^m X_k > t\right] \leq \begin{cases} \max\left\{\exp\left(-\frac{t^2}{40m}\right), \exp\left(-\frac{1}{2}t\right)\right\} & \text{if } \sigma = 1, \\ \max\left\{\exp\left(-\frac{t^2}{40m}\right), \frac{128}{(\sigma-1)^3} \exp\left(-\frac{1}{2\sigma}t^\sigma\right)\right\} & \text{if } \sigma \in (1, 2]. \end{cases} \quad (\text{C.23})$$

In particular,

$$\sum_{k=1}^m X_k = \begin{cases} \mathcal{O}_{\Gamma_1}(40m^{1/2}) & \text{if } \sigma = 1, \\ \mathcal{O}_{\Gamma_\sigma}(\max\{40m^{1/2}, 20|\log(\sigma-1)|^{1/\sigma}\}) & \text{if } \sigma \in (1, 2]. \end{cases} \quad (\text{C.24})$$

*Proof.* Denote  $S_m := X_1 + \dots + X_m$ . We start from the Chernoff bound: for every  $a > 0$ ,

$$\mathbb{P}[S_m \geq a] \leq \inf_{\lambda \in (0, \infty)} \exp(-\lambda a) [\exp(\lambda S_m)] = \inf_{\lambda \in (0, \infty)} \exp(-\lambda a) \prod_{k=1}^N \mathbb{E}[\exp(\lambda X_k)]. \quad (\text{C.25})$$

To estimate  $\mathbb{E}[\exp(\lambda X_k)]$ , we use the elementary inequality

$$|\exp(\lambda x) - (1 + \lambda x)| \leq \frac{1}{2} \lambda^2 |x|^2 \exp(\lambda \max\{x, 0\}), \quad \forall x \in \mathbb{R}, \quad (\text{C.26})$$

and the centering assumption that  $\mathbb{E}[X_k] = 0$  to get

$$\begin{aligned} |\mathbb{E}[\exp(\lambda X_k)] - 1| &\leq \frac{1}{2} \lambda^2 \mathbb{E}[|X_k|^2 \exp(\lambda \max\{X_k, 0\})] \\ &\leq \frac{1}{2} \lambda^2 (\mathbb{E}[|X_k|^2] + \exp(\lambda)) + \frac{1}{2} \lambda^2 \int_1^\infty (2t + \lambda t^2) \exp(\lambda t) \mathbb{P}[X_k > t] dt \\ &\leq \frac{1}{2} \lambda^2 (3 + \exp(\lambda)) + \frac{1}{2} \lambda^2 \int_1^\infty (2t + \lambda t^2) \exp(\lambda t) \mathbb{P}[X_k > t] dt, \end{aligned} \quad (\text{C.27})$$

where in the last line we estimated the second moment by

$$\mathbb{E}[|X_k|^2] \leq 1 + 2 \int_1^\infty t \mathbb{P}[X_k > t] dt \leq 1 + 2 \int_1^\infty t \exp(-t^\sigma) dt \leq 5.$$

We split the estimate of the integral on the right side of (C.27) into two cases:  $\sigma = 1$  and  $\sigma \in (1, 2]$ .

In the case  $\sigma = 1$ , we impose the additional restriction  $\lambda \leq 1/2$ , and then apply the assumption  $X_k = \mathcal{O}_{\Gamma_\sigma}(1)$  to get

$$\begin{aligned} \int_1^\infty (2t + \lambda t^2) \exp(\lambda t) \mathbb{P}[X_k > t] dt &\leq \int_1^\infty (2t + \lambda t^2) \exp(\lambda t - t) dt \\ &\leq \int_1^\infty (2t + \lambda t^2) \exp\left(-\frac{1}{2}t\right) dt = 8(1 + 2\lambda) \leq 16. \end{aligned} \quad (\text{C.28})$$

Combining the above displays, we deduce that, in the case  $\sigma = 1$ , for every  $\lambda \in (0, 1/2]$ ,

$$\mathbb{E}[\exp(\lambda X_k)] \leq 1 + \frac{1}{2} \lambda^2 (5 + \exp(1/2) + 16) \leq 1 + 12\lambda^2.$$

Returning then to (C.25) and using the bound  $1 + x \leq \exp(x)$ , we get

$$\mathbb{P}[S_m \geq a] \leq \exp\left(-\lambda a + 12\lambda^2 m\right).$$

Taking  $\lambda := \min\{1/2, \frac{a}{20m}\}$  yields, for every  $a > 0$ ,

$$\mathbb{P}[S_m \geq a] \leq \exp\left(-\min\left\{\frac{1}{5}a, \frac{a^2}{40m}\right\}\right).$$

This is (C.23) in the case  $\sigma = 1$ . In particular, for every  $a \geq 1$ ,

$$\mathbb{P}[S_m \geq m^{1/2}a] \leq \exp\left(-\min\left\{\frac{1}{5}m^{1/2}a, \frac{a^2}{40}\right\}\right) \leq \exp\left(-\frac{a}{40}\right),$$

which implies that  $S_m = \mathcal{O}_{\Gamma_1}(40m^{1/2})$ .

We next consider the case  $\sigma \in (1, 2]$ . If  $a \leq 10m$ , we may impose the restriction  $\lambda \leq 1/2$  and then we may follow the computation leading to (C.28) in the case  $\sigma = 1$  to get

$$\int_1^\infty (2t + \lambda t^2) \exp(\lambda t) \mathbb{P}[X_k > t] dt \leq \int_1^\infty (2t + \lambda t^2) \exp(\lambda t - t^\sigma) dt \leq 16$$

and then select  $\lambda = \frac{a}{20m}$  to obtain

$$\mathbb{P}[S_m \geq a] \leq \exp\left(-\frac{a^2}{40m}\right), \quad \forall a \in (0, 10m].$$

In the case  $a > 10m$ , we may need to select  $\lambda > 1/2$  and therefore we must estimate the integral differently. Using the assumption that  $X_k = \mathcal{O}_{\Gamma_\sigma}(1)$ , we have that, for every  $\lambda > 0$ ,

$$\begin{aligned} \int_1^\infty (2t + \lambda t^2) \exp(\lambda t) \mathbb{P}[X_k > t] dt &\leq \int_1^\infty (2t + \lambda t^2) \exp(\lambda t - t^\sigma) dt \\ &\leq \exp\left(\frac{\sigma-1}{\sigma} \lambda^{\frac{\sigma}{\sigma-1}}\right) \int_1^\infty (2t + \lambda t^2) \exp\left(-\frac{\sigma-1}{\sigma} t^\sigma\right) dt, \end{aligned}$$

where in the last line we used Cauchy's inequality in the form

$$\lambda t \leq \frac{1}{\sigma} t^\sigma + \frac{\sigma-1}{\sigma} \lambda^{\frac{\sigma}{\sigma-1}}.$$

Continuing the computation, we find by a change of variables that

$$\begin{aligned} \int_1^\infty (2t + \lambda t^2) \exp\left(-\frac{\sigma-1}{\sigma} t^\sigma\right) dt &\leq \frac{1}{\sigma} \int_0^\infty \left(2\left(\frac{\sigma}{\sigma-1}\right)^{\frac{2}{\sigma}} s^{\frac{2}{\sigma}-1} + \lambda\left(\frac{\sigma}{\sigma-1}\right)^{\frac{3}{\sigma}} s^{\frac{3}{\sigma}-1}\right) \exp(-s) ds \\ &= \frac{2}{\sigma} \left(\frac{\sigma}{\sigma-1}\right)^{\frac{2}{\sigma}} \Gamma\left(\frac{2}{\sigma}\right) + \frac{\lambda}{\sigma} \left(\frac{\sigma}{\sigma-1}\right)^{\frac{3}{\sigma}} \Gamma\left(\frac{3}{\sigma}\right) \leq \frac{2+\lambda}{\sigma} \left(\frac{\sigma}{\sigma-1}\right)^{\frac{3}{\sigma}}. \end{aligned}$$

Inserting these bounds into (C.27) yields

$$\mathbb{E}[\exp(\lambda X_k)] \leq 1 + \lambda^2 \left(2 + \frac{8+4\lambda}{(\sigma-1)^3} \exp\left(\frac{\sigma-1}{\sigma} \lambda^{\frac{\sigma}{\sigma-1}}\right)\right).$$

Assuming  $\lambda \geq 1/2$ , we get

$$\mathbb{E}[\exp(\lambda X_k)] \leq \left(12 + \frac{20}{(\sigma-1)^3}\right) \lambda^3 \exp\left(\frac{\sigma-1}{\sigma} \lambda^{\frac{\sigma}{\sigma-1}}\right) \leq \frac{32}{(\sigma-1)^3} \lambda^3 \exp\left(\frac{\sigma-1}{\sigma} \lambda^{\frac{\sigma}{\sigma-1}}\right).$$

Inserting this into (C.25) and selecting  $\lambda = a^{\sigma-1}$ —which we note satisfies the constraint  $\lambda \geq 1/2$  since we have assumed  $a > 10m$ —we obtain

$$\mathbb{P}[S_m \geq a] \leq \frac{32a^{3(\sigma-1)}}{(\sigma-1)^3} \exp\left(-\frac{1}{\sigma} a^\sigma\right) \leq \frac{128}{(\sigma-1)^3} \exp\left(-\frac{1}{2\sigma} a^\sigma\right), \quad \forall a \in [10m, \infty).$$

This completes the proof of (C.23) in the case  $\sigma \in (1, 2]$ , which also implies (C.24).  $\square$

We next present a variant of the concentration argument above for random variables which may have weaker, sub-exponential integrability quantified by a general Orlicz quasinorms. The proof is similar to the arguments of the recent paper [BMdlP23], using a truncation method

**Proposition C.3** (Concentration for sums of independent and  $\mathcal{O}_\Psi$ -bounded random variables).  
*Let  $\Psi : \mathbb{R}_+ \rightarrow [1, \infty)$  be an increasing function satisfying*

$$C_\Psi := \int_1^\infty \frac{t}{\Psi(t)} dt < \infty \quad (\text{C.29})$$

*Let  $m \in \mathbb{N}$  and  $X_1, \dots, X_m$  be a sequence of independent random variables satisfying*

$$X_k = \mathcal{O}_\Psi(1) \quad \text{and} \quad \mathbb{E}[X_k] = 0, \quad \forall k \in \{1, \dots, m\}. \quad (\text{C.30})$$

*Suppose that  $M \in [0, \infty)$ ,  $\lambda \in (0, 1]$  and  $L \in [1, \infty)$  satisfy the relation*

$$\lambda t \leq \log \Psi(t) - 4 \log t + \log M, \quad \forall t \in [1, L]. \quad (\text{C.31})$$

*Then, for every  $t > 0$ ,*

$$\mathbb{P}\left[\sum_{k=1}^m X_k > t\right] \leq \frac{m}{\Psi(L)} + \exp(-\lambda t + \lambda^2 m(2 + M + C_\Psi)). \quad (\text{C.32})$$

*Proof.* Denote  $S_m := X_1 + \dots + X_m$ . Fix  $a > 0$  and  $L \in [1, \infty)$  and define the truncations

$$Y_k := \min\{X_k, L\} \quad \text{and} \quad T_k := \sum_{k=1}^m Y_k.$$

Either  $S_m = T_m$ , or else  $\max_{1 \leq k \leq m} X_k > L$ . We deduce therefore that

$$\mathbb{P}[S_m > a] \leq \mathbb{P}[T_m > a] + \mathbb{P}\left[\max_{1 \leq k \leq m} X_k > L\right]. \quad (\text{C.33})$$

By a union bound and the assumption  $X_k \leq \mathcal{O}_\Psi(1)$ , the second term on the right side of (C.33) is bounded by

$$\mathbb{P}\left[\max_{1 \leq k \leq m} X_k > L\right] \leq \sum_{k=1}^m \mathbb{P}[X_k > L] \leq \frac{m}{\Psi(L)}. \quad (\text{C.34})$$

For the first term, we use the Markov inequality and independence to obtain, for every  $\lambda > 0$ ,

$$\mathbb{P}[T_m > a] \leq \exp(-\lambda a) \mathbb{E}[\exp(\lambda T_m)] \leq \exp(-\lambda a) \prod_{k=1}^m \mathbb{E}[\exp(\lambda Y_k)]. \quad (\text{C.35})$$

In order to estimate  $\mathbb{E}[\exp(\lambda Y_k)]$ , we use (C.26), which implies

$$|\mathbb{E}[\exp(\lambda Y_k)] - (1 + \lambda \mathbb{E}[Y_k])| \leq \frac{1}{2} \lambda^2 \mathbb{E}[|Y_k|^2 \exp(\lambda \max\{Y_k, 0\})].$$

Due to the truncation, the random variable  $Y_k$  is not centered. However,  $\mathbb{E}[Y_k] \leq \mathbb{E}[X_k] \leq 0$ , which suffices for our purposes. We deduce therefore that

$$\mathbb{E}[\exp(\lambda Y_k)] \leq 1 + \frac{1}{2} \lambda^2 \mathbb{E}[Y_k^2 \exp(\lambda \max\{Y_k, 0\})]. \quad (\text{C.36})$$

To estimate the right side of (C.36), we use the hypothesis that  $X_k = \mathcal{O}_\Psi(1)$  to compute

$$\begin{aligned}\mathbb{E}\left[Y_k^2 \exp(\lambda \max\{Y_k, 0\})\right] &= \exp(\lambda) + \int_1^L (2t + \lambda t^2) \exp(\lambda t) \mathbb{P}[X_k > t] dt \\ &\leq \mathbb{E}[X_k^2] + \int_0^1 (2t + \lambda t^2) \exp(\lambda t) dt + \int_1^L \frac{(2t + \lambda t^2) \exp(\lambda t)}{\Psi(t)} dt \\ &= \mathbb{E}[X_k^2] + \exp(\lambda) + \int_1^L (2t + \lambda t^2) \exp(\lambda t - \log \Psi(t)) dt.\end{aligned}$$

We continue now under the assumption that  $\lambda \in (0, 1]$  and  $L \in [1, \infty)$  satisfy (C.31). We have

$$\int_1^L (2t + \lambda t^2) \exp(\lambda t - \log \Psi(t)) dt \leq M \int_1^L (2t + \lambda t^2) t^{-4} dt \leq M \int_1^\infty (2t + \lambda t^2) t^{-4} dt = \left(\frac{2}{3} + \frac{\lambda}{3}\right) M.$$

For the second moments of  $X_k$ , we use the condition (C.29), which implies that

$$\mathbb{E}[X_k^2] \leq 1 + 2C_\Psi.$$

We therefore obtain that

$$\mathbb{E}[\exp(\lambda Y_k)] \leq 1 + \frac{1}{2} \lambda^2 (\mathbb{E}[X_k^2] + \exp(\lambda) + M) \leq 1 + \frac{1}{2} \lambda^2 (4 + M + 2C_\Psi) \leq \exp\left(\frac{1}{2} \lambda^2 (4 + M + 2C_\Psi)\right).$$

Inserting this result into (C.35), we obtain, for every  $\lambda \in (0, 1]$  and  $L \geq 1$  satisfying (C.31),

$$\mathbb{P}[T_m > a] \leq \exp\left(-\lambda a + \frac{1}{2} \lambda^2 m (4 + M + 2C_\Psi)\right). \quad (\text{C.37})$$

Inserting (C.34) and (C.37) into (C.33) yields (C.32), completing the proof.  $\square$

We next exhibit consequences of Proposition C.3 for some particular heavy-tailed distributions, including stretched exponentials (Weibull distributions) and those with log-normal-type tails.

**Corollary C.4** (Concentration for stretched exponential tails). *Let  $\sigma \in (0, 1)$  and define*

$$\Gamma_\sigma(t) := \exp(t^\sigma), \quad t \in [1, \infty). \quad (\text{C.38})$$

*Let  $m \in \mathbb{N}$  and  $X_1, \dots, X_m$  be a sequence of independent random variables satisfying*

$$X_k = \mathcal{O}_{\Gamma_\sigma}(1) \quad \text{and} \quad \mathbb{E}[X_k] = 0, \quad \forall k \in \{1, \dots, m\}. \quad (\text{C.39})$$

*Then, for every  $t \geq 1$ ,*

$$\mathbb{P}\left[\sum_{k=1}^m X_k > t\right] \leq (m+1) \exp(-4t^\sigma) + \exp\left(-\frac{t^2}{4((8^{2/\sigma} \Gamma(2/\sigma))^4 + 1)m}\right). \quad (\text{C.40})$$

*In particular,*

$$\sum_{k=1}^m X_k \leq \mathcal{O}_{\Gamma_\sigma}\left(2((8^{2/\sigma} \Gamma(2/\sigma))^2 + 1)m^{1/2}\right). \quad (\text{C.41})$$

**Corollary C.5** (Concentration for log-normal tails). *As in (C.19), we define, for  $\sigma \in [1, \infty)$ ,*

$$\Psi_\sigma(t) := \exp\left(\frac{1}{\sigma^2} \log^2(1 + \sigma t)\right), \quad t \in [1, \infty). \quad (\text{C.42})$$

*Let  $m \in \mathbb{N}$  and  $X_1, \dots, X_m$  be a sequence of independent random variables satisfying*

$$X_k = \mathcal{O}_{\Psi_\sigma}(1) \quad \text{and} \quad \mathbb{E}[X_k] = 0, \quad \forall k \in \{1, \dots, m\}. \quad (\text{C.43})$$

*Then, for every  $t \geq 4\sigma$ ,*

$$\mathbb{P}\left[\sum_{k=1}^m X_k > t\right] \leq (m+1) \exp\left(-\frac{1}{\sigma^2} \log^2 \frac{\sigma t}{1600}\right) + \exp\left(-\frac{t^2}{(2 \exp(32\sigma^2) + 20)m}\right). \quad (\text{C.44})$$

*In particular,*

$$\sum_{k=1}^m X_k \leq \mathcal{O}_{\Psi_\sigma}(32 \exp(16\sigma^2) m^{1/2}). \quad (\text{C.45})$$

*Proof of Corollary C.4.* The constant  $C_{\Gamma_\sigma}$  defined in (C.29) is given by

$$C_{\Gamma_\sigma} = \int_1^\infty \frac{t}{\Gamma_\sigma(t)} dt = \int_1^\infty t \exp(-t^\sigma) dt = \frac{1}{\sigma} \int_1^\infty s^{\frac{2}{\sigma}-1} \exp(-s) ds \leq \frac{1}{\sigma} \Gamma\left(\frac{2}{\sigma}\right),$$

where  $\Gamma$  denotes the gamma function. We also have<sup>13</sup>

$$\sigma \in (0, 1), \quad M = M_\sigma := (8^{2/\sigma} \Gamma(2/\sigma))^4, \quad \lambda \in (0, 1], \quad 1 \leq L \leq (2\lambda)^{-\frac{1}{1-\sigma}} \implies (\text{C.31}). \quad (\text{C.46})$$

By applying Proposition C.3, we therefore obtain, for every  $\lambda \in (0, 1]$  and  $t \geq 1$ ,

$$\mathbb{P}\left[\sum_{k=1}^m X_k > t\right] \leq m \exp\left(-(2\lambda)^{-\frac{\sigma}{1-\sigma}}\right) + \exp(-\lambda t + A_\sigma \lambda^2 m). \quad (\text{C.47})$$

where we set  $A_\sigma := 2((\Gamma(2/\sigma) 8^{2/\sigma})^4 + 1) \geq (2 + M_\sigma + C_{\Gamma_\sigma})$ . Given  $t \in [1, \infty)$ , we apply the above inequality with  $\lambda$  selected by

$$\lambda := \min\left\{\frac{t}{2A_\sigma m}, \frac{1}{2}t^{\sigma-1}\right\}.$$

Note that this ensures  $\lambda \leq 1$  and, with this choice, the first and second terms on the right side of (C.47) are estimated, respectively, by

$$\exp(-\lambda t + A_\sigma \lambda^2 m) \leq \exp\left(-\frac{1}{2}\lambda t\right) = \max\left\{\exp\left(-\frac{t^2}{2A_\sigma m}\right), \exp\left(-\frac{1}{4}t^\sigma\right)\right\}$$

and

$$m \exp\left(-(2\lambda)^{-\frac{\sigma}{1-\sigma}}\right) \leq m \exp(-t^\sigma).$$

Combining the above displays yields (C.40). □

---

<sup>13</sup>The proof of (C.46) can be found in the latex file for this paper (available on arXiv), commented out below this sentence.

*Proof of Corollary C.5.* We check that the constant in (C.29) satisfies

$$C_{\Psi_\sigma} \leq \exp(4\sigma^2) \quad (\text{C.48})$$

and that the condition (C.31) is satisfied with  $M = M_\sigma := \exp(32\sigma^2)$  and every  $\lambda \in (0, 1]$  and  $L$  satisfying<sup>14</sup>

$$1 \leq L \leq \frac{1}{32\sigma^2\lambda} \log^2\left(1 + \frac{1}{\sigma\lambda}\right). \quad (\text{C.49})$$

Applying Proposition C.3 therefore yields, for every  $t > 0$  and  $\lambda \in (0, 1]$ ,

$$\mathbb{P}\left[\sum_{k=1}^m X_k > t\right] \leq \frac{m}{\Psi_\sigma\left(\frac{1}{32\sigma^2\lambda} \log^2\left(1 + \frac{1}{\sigma\lambda}\right)\right)} + \exp(-\lambda t + A_\sigma \lambda^2 m), \quad (\text{C.50})$$

where we have set

$$A_\sigma := 2\exp(32\sigma^2) + 5 \geq (2 + M_\sigma + C_{\Psi_\sigma}).$$

We apply the above inequality to each  $t \geq 4\sigma$  with  $\lambda$  chosen in terms of  $t$  by

$$\lambda := \min\left\{\frac{t}{2A_\sigma m}, \frac{2}{\sigma^2 t} \log^2(1 + \sigma t)\right\}.$$

This choice of  $\lambda$  implies that

$$\exp(-\lambda t + A_\sigma \lambda^2 m) \leq \exp\left(-\frac{1}{2}\lambda t\right) = \max\left\{\exp\left(-\frac{t^2}{4A_\sigma m}\right), \exp\left(-\frac{1}{\sigma^2} \log^2(1 + \sigma t)\right)\right\}$$

as well as

$$\frac{1}{32\sigma^2\lambda} \log^2\left(1 + (\sigma\lambda)^{-1}\right) \geq \frac{t}{64} \frac{\log^2(1 + (\sigma\lambda)^{-1})}{\log^2(1 + \sigma t)} \geq \frac{t}{64} \left(\frac{\log(1 + \frac{1}{2}\sigma t \log^{-2}(1 + \sigma t))}{\log(1 + \sigma t)}\right)^2 \geq \frac{t}{1600}.$$

In the last inequality of the previous display, we used that

$$\log\left(1 + \frac{1}{2}s \log^{-2}(1 + s)\right) \geq \frac{1}{5} \log(1 + s), \quad \forall s \in [1, \infty). \quad (\text{C.51})$$

The validity of the inequality (C.51) for some (universal)  $c > 0$  in place of the  $1/5$  on the right side is clear, due to the fact that the ratio of the left side and  $\log(1 + s)$  is positive for all  $s \geq 1$  and tends to one as  $s \rightarrow \infty$ . That the inequality is valid as stated with this constant equal to  $1/5$  can be confirmed by either Mathematica or Wolfram Alpha. We used the command

$$\text{Reduce}[\text{Log}[1 + s/(2 \text{Log}[1 + s]^2)] > 1/5 \text{Log}[1 + s], s, \text{Reals}]$$

in Mathematica to validate (C.51), which instantly reported its validity for all  $s > 0$ . Combining the above displays, we obtain (C.44).

To obtain (C.45), we use (C.44) with  $4A_\sigma^{1/2} m^{1/2} t$  in place of  $t$ , to obtain, for every  $t \geq 1$  (note that  $A_\sigma^{1/2} \geq \sigma$ ),

$$\mathbb{P}\left[\sum_{k=1}^m X_k > 4A_\sigma^{1/2} m^{1/2} t\right] \leq \frac{m+1}{\Psi_\sigma\left(\frac{1}{400} A_\sigma^{1/2} m^{1/2} t\right)} + \exp(-4t^2).$$

---

<sup>14</sup>For completeness we have included full demonstrations of these assertions in a commented-out part of the latex file for this paper available from the arXiv.



By an easy exercise, it can be checked<sup>15</sup> that, for every  $\sigma, t, m \geq 1$ ,

$$\frac{m+1}{\Psi_\sigma\left(\frac{1}{400}A_\sigma^{1/2}m^{1/2}t\right)} \leq \frac{1}{2\Psi_\sigma(t)} \quad \text{and} \quad \exp(-4t^2) \leq \frac{1}{2\Psi_\sigma(t)}.$$

We also observe that  $4A_\sigma^{1/2} \leq 8\exp(16\sigma^2)$ , for every  $\sigma \geq 1$ . This completes the proof of (C.45).  $\square$

## D. Examples of random fields satisfying the assumptions

**D.1. Poisson inclusions.** In this subsection we prove Proposition 1.1. We consider two Poisson point clouds  $\omega_1$  and  $\omega_2$  on  $\mathbb{R}^d$  with intensities  $\rho_1 \geq 0$  and  $\rho_2 \geq 0$ , respectively. Let  $\lambda \in (0, 1]$ ,  $\Lambda \in [1, \infty)$  and define the scalar matrix-valued field

$$\mathbf{a} := (1 + (\Lambda - 1)\mathbf{1}_{B_{1/3}} * \omega_1 + (\lambda - 1)\mathbf{1}_{B_{1/3}} * \omega_2)\mathbf{I}_d. \quad (\text{D.1})$$

Notice that our inclusions are balls of radius  $1/3$  rather than unit radius like in (1.42). We have introduced this extra dilation for notational convenience. We also denote  $\rho := \rho_1 + \rho_2$  and  $\omega = \omega_1 + \omega_2$ .

We adapt some arguments and notation from classical percolation theory. We view  $\mathbb{Z}^d$  as an undirected graph with vertices  $x, y \in \mathbb{Z}^d$  connected by an edge if and only if  $\max_{i \in \{1, \dots, d\}} |x_i - y_i| = 1$ . When we speak of *connected* subsets of  $\mathbb{Z}^d$ , we mean those which are connected with respect to this graph structure. A *lattice animal* is a finite, connected subset of  $\mathbb{Z}^d$ . As is well-known, a crude combinatorial counting argument gives that, for each fixed  $z_0 \in \mathbb{Z}^d$ , the number of distinct lattice animals which contains  $z_0$  and has exactly  $\ell$  elements is at most  $\exp(C\ell)$  for some  $C(d) < \infty$ . For each lattice animal  $A \subseteq \mathbb{Z}^d$  with  $|A| = \ell$ , the probability that every unit cell  $z + \square_0$  with  $z \in A$  has nonempty intersection with  $\omega$  is estimated by

$$\mathbb{P}\left[\forall z \in A, (z + \square_0) \cap \omega \neq \emptyset\right] \leq \rho^\ell = \exp(-\ell \log \rho).$$

Let  $E_\ell(z)$  denote the event that there exists *any* lattice animal which has length at least  $\ell$ , has nontrivial intersection with  $z + \ell\square_0$ , and overlaps with points of  $\omega$  in each of its cells:

$$E_\ell(z) := \left\{ \begin{array}{l} \text{there exists a lattice animal } A \subseteq \mathbb{R}^d \text{ with } |A| \geq \ell \text{ and} \\ A \cap (z + \ell\square_0) \neq \emptyset \text{ such that, for every } z' \in A, (z' + \square_0) \cap \omega \neq \emptyset \end{array} \right\}.$$

A union bound yields that the probability of  $E_\ell(0)$  is at most

$$\mathbb{P}[E_\ell(0)] \leq \sum_{k=\ell}^{\infty} \ell^d \exp(Ck - k \log \rho).$$

If we restrict the intensity  $\rho$  of the Poisson cloud  $\omega$  by requiring  $\rho \leq c$  for sufficiently small  $c(d) > 0$ , then we obtain

$$\mathbb{P}[E_\ell] \leq \exp\left(d \log \ell - \frac{1}{3}\ell \log \rho\right) \leq \exp(-c\ell \log \rho). \quad (\text{D.2})$$

By (C.18), this implies that, for every  $\sigma \in (0, \infty)$ ,

$$\mathbf{1}_{E_\ell} \leq \mathcal{O}_{\Gamma_{1/\sigma}}(C^\sigma(\ell \log \rho)^{-\sigma}) \quad (\text{D.3})$$

---

<sup>15</sup>The proof of these inequalities are routine, but for completeness we have also commented them out in the latex file after this sentence.

We next discretize  $\omega$  by setting

$$\hat{\omega} := \{z \in \mathbb{Z}^d : (z + \square_0) \cap \omega \neq \emptyset\}.$$

For each  $m \in \mathbb{N}$ , we define  $\mathbb{N}$ -valued stationary random fields  $N$  and  $N_m$  on  $\mathbb{Z}^d$  by

$$\begin{cases} N(z) := \text{the number of elements in the connected component of } \hat{\omega} \text{ containing } z, \\ N_m(z) := \text{the number of elements in the connected component of } \hat{\omega} \cap (z + \square_m) \text{ containing } z. \end{cases}$$

Note that  $N(z) = N_m(z) = 0$  if  $z \notin \hat{\omega}$ . It is clear that  $N_m$  is nondecreasing as a function of  $m$ . In view of (D.2), for every  $m \in \mathbb{N}$ ,

$$N_m(0) \leq N(0) \leq \mathcal{O}_{\Gamma_1}(C|\log \rho|^{-1}). \quad (\text{D.4})$$

It is also clear that  $N_{m+1}(z)$  differs from  $N_m(z)$  for  $z \in \square_m$  only if  $N_m(z) > 3^m$ , and thus only on the event  $E_{3^m}$ . More generally, for every  $n, m \in \mathbb{N}$  with  $n < m$ ,

$$\begin{aligned} \mathbb{P}[\exists z \in \square_m, N_m(z) \neq N_n(z)] &\leq \mathbb{P}[\exists z \in \square_m, E_{3^n}(z) > 3^n] \leq 3^{md} \exp(-c3^n |\log \rho|) \\ &= \exp(Cm - c3^n |\log \rho|). \end{aligned} \quad (\text{D.5})$$

This implies that  $N_m$  has an approximate finite range of dependence property, which allows us to use concentration inequalities.

**Lemma D.1.** *Let  $\mathbf{a}(\cdot)$  be the field defined in (D.1), with parameters  $\rho, \lambda$  and  $\Lambda$  as introduced there. There exist constants  $c(d) > 0$  and  $C(d) < \infty$  such that, if  $\rho \leq c$  and  $\gamma \in (0, 1)$ , then there exists a minimal scale  $\mathcal{S}$  satisfying  $\mathcal{S} \leq \mathcal{O}_{\Psi_{\mathcal{S}}}(1)$  and*

$$3^m \geq \mathcal{S} \implies \mathbf{A}(z + \square_n) \leq 3^{\gamma(m-n)} \mathbf{E}_0 \quad \forall n \in \mathbb{Z} \cap (-\infty, m], \quad \forall z \in 3^n \mathbb{Z}^d \cap \square_m$$

with

$$\mathbf{E}_0 := (1 + C|\log \rho|^{-2}) \mathbf{I}_{2d} \quad \text{and} \quad \Psi_{\mathcal{S}}(t) := \exp\left(c(\Lambda \vee \lambda^{-1})^{-\frac{1}{d+2} - \frac{\gamma}{d}} t^{\frac{\gamma}{d+2}} - 1\right).$$

Moreover,  $\Psi_{\mathcal{S}}$  satisfies

$$t\Psi_{\mathcal{S}}(t) \leq \Psi_{\mathcal{S}}(K_{\Psi_{\mathcal{S}}} t) \quad \forall t \in [1, \infty) \quad \text{with} \quad K_{\Psi_{\mathcal{S}}} := (C\gamma^{-1})^{\frac{d+2}{\gamma}} (\Lambda \vee \lambda^{-1})^{\frac{1}{\gamma} + 1 + \frac{2}{d}}.$$

*Proof.* We construct, for each  $m \in \mathbb{N}$ , a partition of the cube  $\square_m$ . We let  $\mathcal{C}_m(\omega)$  denote the collection of connected components of  $\hat{\omega} \cap \square_m$ . For each  $A \in \mathcal{C}_m(\omega)$ , we associate the continuum set

$$\tilde{A} := \bigcup_{z \in A} (z + \square_0)$$

and slightly enlarge this set by defining

$$\check{A} := \{x \in \square_m : \text{dist}(x, \tilde{A}) < 1/3\}.$$

Observe that  $\check{A}_1 \cap \check{A}_2 = \emptyset$  if  $A_1$  and  $A_2$  are distinct connected components of  $\hat{\omega}$ . If  $\tilde{A}$  does not touch the boundary  $\partial \square_m$ , we write  $A \in \mathcal{C}_m^\circ(\omega)$ . We also let  $\mathcal{C}_m^b(\omega) := \mathcal{C}_m(\omega) \setminus \mathcal{C}_m^\circ(\omega)$  be those connected components for which  $\tilde{A}$  does touch  $\partial \square_m$ .

We let  $\mathcal{P}$  be the collection of all subsets of  $\square_m$  of the form: (i)  $\tilde{A}$ , for  $A \in \mathcal{C}_m(\omega)$ ; (ii)  $\check{A} \setminus \tilde{A}$ , for  $A \in \mathcal{C}_m(\omega)$ ; and (iii)  $\square_m \setminus (\cup \{\check{A} : A \in \mathcal{C}_m(\omega)\})$ . It is clear that  $\mathcal{P}$  is a partition of  $\square_m$ , up to a zero Lebesgue measure set.

Given  $p \in \mathbb{R}^d$ , we define a gradient field  $\nabla \phi_p$  on  $\square_m$  with the following properties:

- $\nabla\phi_p$  vanishes on  $\tilde{A}$ , for every  $A \in \mathcal{C}_m^\circ(\omega)$ .
- $\nabla\phi_p = p$  in  $\square_m \setminus \cup \{\tilde{A} : A \in \mathcal{C}_m^\circ(\omega)\}$ .
- For each  $A \in \mathcal{C}_m^\circ(\omega)$ , we have  $|\nabla\phi_p| \leq 20|p||A|$  in  $\tilde{A} \setminus \tilde{A}$ .

In particular, since  $\nabla\phi_p = p$  in a neighborhood of the boundary  $\partial\square_m$ , we have that

$$p \cdot \mathbf{a}(\square_m)p \leq \min_{u \in \ell_p + H_0^1(\square_m)} \int_{\square_m} \nabla u \cdot \mathbf{a} \nabla u \leq \int_{\square_m} \nabla\phi_p \cdot \mathbf{a} \nabla\phi_p.$$

Since  $\nabla\phi_p$  vanishes in  $\tilde{A}$  for each  $A \in \mathcal{C}_m^\circ(\omega)$ , we have that

$$\sum_{A \in \mathcal{C}_m^\circ(\omega)} \int_{\tilde{A}} \nabla\phi_p \cdot \mathbf{a} \nabla\phi_p = 0.$$

Since  $\mathbf{a}(x) = \mathbf{I}_d$  in  $B := \square_m \setminus \cup \{\tilde{A} : A \in \mathcal{C}_m(\omega)\}$  and  $\mathbf{a}(x) \leq \Lambda \mathbf{I}_d$  otherwise, we deduce that

$$\int_{\square_m} \nabla\phi_p \cdot \mathbf{a} \nabla\phi_p = \frac{1}{|\square_m|} \int_B |\nabla\phi_p|^2 + \frac{\Lambda}{|\square_m|} \sum_{A \in \mathcal{C}_m^b(\omega)} \int_{\tilde{A}} |\nabla\phi_p|^2.$$

Using the properties of  $\nabla\phi_p$ , we have

$$\begin{aligned} \frac{1}{|\square_m|} \int_B |\nabla\phi_p|^2 &\leq |p|^2 + \frac{1}{|\square_m|} \sum_{A \in \mathcal{C}_m(\omega)} 400|p|^2|A|^2|\tilde{A} \setminus \tilde{A}| \leq |p|^2 + \frac{800|p|^2}{|\square_m|} \sum_{A \in \mathcal{C}_m(\omega)} |A|^3 \\ &\leq |p|^2 \left( 1 + 800 \sum_{z \in \square_m} N_{m+1}(z)^2 \right) \end{aligned}$$

and, similarly,

$$\begin{aligned} \frac{\Lambda}{|\square_m|} \sum_{A \in \mathcal{C}_m^b(\omega)} \int_{\tilde{A}} |\nabla\phi_p|^2 &\leq \frac{400\Lambda|p|^2}{|\square_m|} \sum_{A \in \mathcal{C}_m^b(\omega)} |A|^2 \leq \frac{400\Lambda|p|^2}{|\square_m|} \sum_{z \in \partial\square_m} N_{m+1}(z)^2 \\ &\leq 400|p|^2 3^{-m} \Lambda \sum_{z \in \partial\square_m} N_{m+1}(z)^2. \end{aligned}$$

Next, let  $n \in \mathbb{N}$  with  $n < m$  and observe that

$$\sum_{z \in \square_m} N_{m+1}(z)^2 \leq \sum_{z \in \square_m} N_n(z)^2 + \sum_{z \in \square_m} \mathbf{1}_{E_{3^n}(z)} N_{m+1}(z)^2.$$

Using (D.4) and (C.40), and noting that the mean of  $N_n(z)$  is at most  $C|\log \rho|^{-2}$  by (D.4), we find that, for every  $t \geq C|\log \rho|^{-2}$ ,

$$\mathbb{P} \left[ \sum_{z \in \square_m} N_n(z)^2 > t \right] \leq 3^{d(m-n)} \exp(-c3^{\frac{d}{2}(m-n)} t^{\frac{1}{2}}) + \exp(-c3^{d(m-n)} t^2).$$

This yields, for  $t \geq C|\log \rho|^{-2}$  and  $\rho$  sufficiently small,

$$\mathbb{P} \left[ \sum_{z \in \square_m} N_n(z)^2 > t \right] \leq \exp(C(m-n) - ct^{\frac{1}{2}} 3^{\frac{d}{2}(m-n)}) + \exp(-ct^2 3^{d(m-n)}) \leq \exp(-ct^{\frac{1}{2}} 3^{\frac{d}{2}(m-n)}).$$

By (D.5), we have that

$$\mathbb{P}\left[\sum_{z \in \square_m} \mathbf{1}_{E_{3^n}(z)} N_{m+1}(z)^2 \neq 0\right] \leq \mathbb{P}[\exists z \in \square_m, E_{3^n}(z) > 3^n] \leq \exp(Cm - c3^n |\log \rho|).$$

Combining the above yields, for every  $t \geq C|\log \rho|^{-2}$ ,

$$\mathbb{P}\left[\sum_{z \in \square_m} N_{m+1}(z)^2 > t\right] \leq \exp(-ct^{\frac{1}{2}} 3^{\frac{d}{2}(m-n)}) + \exp(Cm - c3^n |\log \rho|).$$

By a very similar computation, we find that

$$\mathbb{P}\left[\sum_{z \in \partial \square_m} N_{m+1}(z)^2 > t\right] \leq \exp(-ct^{\frac{1}{2}} 3^{\frac{d-1}{2}(m-n)}) + \exp(Cm - c3^n |\log \rho|).$$

Putting these together, we obtain, for every  $t \geq C|\log \rho|^{-2}$  and  $m, n \in \mathbb{N}$  with  $n < m$  and  $3^n \geq \Lambda$ ,

$$\mathbb{P}[\mathbf{a}(\square_m) \not\leq (1+t)\mathbf{I}_d] \leq \exp(-ct^{\frac{1}{2}} 3^{\frac{d}{2}(m-n)}) + \exp(Cm - c3^n |\log \rho|). \quad (\text{D.6})$$

By essentially the same argument, we also obtain an estimate for  $\mathbf{a}_*^{-1}(\square_m)$ , which states that, for every  $t \geq C|\log \rho|^{-2}$  and  $m, n \in \mathbb{N}$  with  $n < m$  and  $3^n \geq \Lambda \vee \lambda^{-1}$ ,

$$\mathbb{P}[\mathbf{a}_*^{-1}(\square_m) \not\leq (1+t)\mathbf{I}_d] \leq \exp(-ct^{\frac{1}{2}} 3^{\frac{d}{2}(m-n)}) + \exp(Cm - c3^n |\log \rho|). \quad (\text{D.7})$$

To prove (D.7), we start from the variational formula

$$q \cdot \mathbf{a}_*^{-1}(\square_m) q = \min_{\mathbf{g} \in q + L_{\text{sol},0}^2(\square_m)} \int_{\square_m} \mathbf{g} \cdot \mathbf{a}^{-1} \mathbf{g}.$$

We test this formula with a divergence-free field  $\mathbf{h}_q$  on  $\square_m$  which has the following properties: for a constant  $C(d) < \infty$ ,

- $\mathbf{h}_q$  vanishes on  $\tilde{A}$ , for every  $A \in \mathcal{C}_m^\circ(\omega)$ .
- $\mathbf{h}_q = q$  in  $\square_m \setminus \cup \{\check{A} : A \in \mathcal{C}_m^\circ(\omega)\}$ .
- For each  $A \in \mathcal{C}_m^\circ(\omega)$ , we have  $|\mathbf{h}_q| \leq C|q||A|$  in  $\check{A} \setminus \tilde{A}$ .

Such a divergence-free field is relatively straightforward to construct (even if it is less obvious than for the analogous gradient field above). The argument for (D.7) then follows nearly identically to that of (D.6), with  $\mathbf{h}_q$  in place of  $\nabla \phi_p$ .

By combining (D.6) and (D.7) we obtain, for every  $t \geq C|\log \rho|^{-2}$  and  $m, n \in \mathbb{N}$  with  $n < m$  and  $3^n \geq \Lambda \vee \lambda^{-1}$ ,

$$\mathbb{P}[\mathbf{A}(\square_m) \not\leq (1+t)\mathbf{I}_{2d}] \leq \exp(-ct^{\frac{1}{2}} 3^{\frac{d}{2}(m-n)}) + \exp(Cm - c3^n |\log \rho|).$$

Optimizing the parameter  $n$  leads to choose  $n$  so that  $3^{(\frac{d}{2}+1)n} \simeq t^{\frac{1}{2}} 3^{\frac{d}{2}m}$ . We obtain, for every  $t \geq C|\log \rho|^{-2}$  and  $m \in \mathbb{N}$  with  $3^m \geq (\Lambda \vee \lambda^{-1})^{1+\frac{2}{d}}$ ,

$$\mathbb{P}[\mathbf{A}(\square_m) \not\leq (1+t)\mathbf{I}_{2d}] \leq \exp(-ct^{\frac{1}{d+2}} 3^{\frac{d}{d+2}m}).$$

Let  $n_0 \in \mathbb{N}$  be the smallest integer such that  $3^{n_0} \geq (\Lambda \vee \lambda^{-1})^{1+\frac{2}{d}} \vee \exp(C\gamma^{-1})$ . A union bound and the above display now give us

$$\begin{aligned}
& \mathbb{P}\left[\exists n \in \mathbb{N} \cap [n_0, m], \ z \in 3^n \mathbb{Z}^d \cap \square_m, \ \mathbf{A}(z + \square_n) \not\leq 3^{\gamma(m-n)}(1 + C|\log \rho|^{-2})\mathbf{I}_{2d}\right] \\
& \leq \sum_{n=n_0}^m \sum_{z \in 3^n \mathbb{Z}^d \cap \square_m} \mathbb{P}\left[\mathbf{A}(z + \square_n) \not\leq 3^{\gamma(m-n)}(1 + C|\log \rho|^{-2})\mathbf{I}_{2d}\right] \\
& \leq \sum_{n=n_0}^m \exp(C(m-n) - c3^{\frac{\gamma}{d+2}(m-n)}3^{\frac{d}{d+2}n}) \\
& \leq \exp(-c3^{\frac{\gamma}{d+2}m + \frac{d-\gamma}{d+2}n_0}).
\end{aligned} \tag{D.8}$$

On the other hand, we have the quenched bound

$$3^{-\gamma(m-n_0)}\mathbf{A}(z + \square_n) \leq C3^{\gamma n_0}(\Lambda \vee \lambda^{-1})3^{-\gamma m}\mathbf{I}_{2d} \leq C(\Lambda \vee \lambda^{-1})^{1+\gamma(1+\frac{2}{d})}3^{-\gamma m}\mathbf{I}_{2d},$$

so that if  $m_0 \in \mathbb{N}$  is the smallest integer such that

$$3^{\gamma m_0} \geq C(\Lambda \vee \lambda^{-1})^{1+\gamma(1+\frac{2}{d})},$$

we have

$$m \geq m_0, \quad n \leq n_0 \quad \implies \quad \mathbf{A}(z + \square_n) \leq 3^{\gamma(m-n)}\mathbf{I}_{2d} \quad \forall z \in 3^n \mathbb{Z}^d \cap \square_m. \tag{D.9}$$

Defining now, for  $\mathbf{E}_0 := (1 + C|\log \rho|^{-2})\mathbf{I}_{2d}$ , the minimal scale  $\mathcal{S}$  by

$$\mathcal{S} := \sup \left\{ 3^{m+1} : m \in \mathbb{N} \cap [m_0, \infty), \sup_{n \in \mathbb{N} \cap (-\infty, m]} \sup_{z \in 3^n \mathbb{Z}^d \cap \square_m} \mathbf{A}(z + \square_n) \not\leq 3^{\gamma(m-n)}\mathbf{E}_0 \right\},$$

we obtain

$$3^m \geq \mathcal{S} \implies \mathbf{A}(z + \square_n) \leq 3^{\gamma(m-n)}\mathbf{E}_0 \quad \forall n \in \mathbb{Z} \cap (-\infty, m], \quad \forall z \in 3^n \mathbb{Z}^d \cap \square_m.$$

By (D.8), (D.9) and a union bound we have

$$\mathcal{S} \leq \mathcal{O}_{\Psi_{\mathcal{S}}}(1) \quad \text{with} \quad \Psi_{\mathcal{S}}(t) = \exp\left(c(3^{-m_0}t)^{\frac{\gamma}{d+2}} - 1\right).$$

By a direct computation we also deduce that

$$K_{\Psi_{\mathcal{S}}} := \left(1 + C\gamma^{-1}3^{\frac{\gamma}{d+2}m_0}\right)^{\frac{d+2}{\gamma}} \implies t\Psi_{\mathcal{S}}(t) \leq \Psi(K_{\Psi_{\mathcal{S}}}t) \quad \forall t \in [1, \infty),$$

which yields the statement.  $\square$

**D.2. Fractional Gaussian fields.** In this section, we review some basic facts about fractional Gaussian fields and verify the claim made in the introduction that these fields give rise to examples of random elliptic coefficient fields satisfying our hypotheses. Many of the general

*D.2.1. Definition of fractional Gaussian fields.* We begin with the definition and basic properties of fractional Gaussian fields. Many of the facts presented here can be found in [LSSW16], but we include proofs and full details of the computations for the reader's convenience.

We denote by  $W$  a standard Gaussian white noise process on  $\mathbb{R}^d$ . It is a random distribution on  $\mathbb{R}^d$ , that is, a random element of  $\mathcal{S}'(\mathbb{R}^d)$ , the dual of the space  $\mathcal{S}(\mathbb{R}^d)$  of Schwarz functions on  $\mathbb{R}^d$ . The field  $W$  is characterized by two properties: first, that  $W(\psi)$  is a Gaussian random variable for each  $\psi \in \mathcal{S}(\mathbb{R}^d)$ ; and second, that the following covariance formula is satisfied:

$$\text{cov}[W(\psi_1), W(\psi_2)] = \int_{\mathbb{R}^d} \psi_1(x) \psi_2(x) dx, \quad \forall \psi_1, \psi_2 \in \mathcal{S}(\mathbb{R}^d). \quad (\text{D.10})$$

The distribution  $W$  almost surely belongs to  $H_{\text{loc}}^{-d/2-\varepsilon}(\mathbb{R}^d)$ , for every  $\varepsilon > 0$ , but not  $H_{\text{loc}}^{-d/2}(\mathbb{R}^d)$ . Immediate from the covariance formula is the following scaling invariance for  $W$ : for every  $\lambda > 0$ ,

$$\lambda^{d/2} W(\lambda \cdot) \quad \text{has the same law as } W. \quad (\text{D.11})$$

A proof of these facts, as well as an explicit construction of  $W$ , can be found in [AKM19, Chapter 5]. In what follows, we abuse notation by informally writing  $\int_{\mathbb{R}^d} \psi(x) W(x) dx$  in place of  $W(\psi)$ .

White noise is an example of a self-similar fractional Gaussian process. These fields are typically indexed by the *Hurst parameter*, which is roughly the regularity of the field. The white noise field  $W$  has Hurst parameter  $-d/2$ .

We denote the self-similar fractional Gaussian field with Hurst parameter  $-\sigma$ , with  $\sigma \in (0, d/2)$ , by  $F_\sigma$ . It is characterized by the fact that  $F_\sigma(\psi)$  is a Gaussian random variable for each fixed test function  $\psi \in \mathcal{S}(\mathbb{R}^d)$ , and the covariance formula (cf. [LSSW16, Theorem 3.3])

$$\text{cov}[F_\sigma(\psi_1), F_\sigma(\psi_2)] = C(\sigma, d) \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |x - y|^{-2\sigma} \psi_1(x) \psi_2(y) dx dy, \quad \forall \psi_1, \psi_2 \in \mathcal{S}(\mathbb{R}^d), \quad (\text{D.12})$$

where  $C(\sigma, d) > 0$  is the special constant defined by

$$C(\sigma, d) := \frac{2^{2\sigma-d} \pi^{-d/2} \Gamma(\sigma)}{\Gamma(d/2 - \sigma)}. \quad (\text{D.13})$$

Note that  $\Gamma(\sigma)$  is of order  $\sigma^{-1}$  as  $\sigma \rightarrow 0$ , and thus so is  $C(\sigma, d)$ .

The field  $F_\sigma$  can be constructed explicitly in terms of  $W$ , in fact, as a deterministic function of  $W$ . It is (in)formally the convolution

$$F_\sigma := 2^\sigma (2\pi)^{-d/2} |x|^{-(\frac{d}{2}+\sigma)} * W.$$

This convolution is however not well-defined. Making of sense of it can be accomplished in various ways. Here we use the following integral identity: for every  $q \in (0, \infty)$  and  $x \in \mathbb{R}^d \setminus \{0\}$ ,

$$|x|^{-q} = \frac{(4\pi)^{d/2}}{2^q \Gamma(q/2)} \int_0^\infty t^{-\frac{1}{2}(2-d+q)} \Phi(t, x) dt. \quad (\text{D.14})$$

Here  $\Gamma$  is the gamma function defined by  $\Gamma(\alpha) := \int_0^\infty t^{-1+\alpha} \exp(-t) dt$ , and  $\Phi$  denotes the standard heat kernel on  $\mathbb{R}^d$ , defined by

$$\Phi(t, x) := (4\pi t)^{-d/2} \exp\left(-\frac{|x|^2}{4t}\right), \quad (t, x) \in (0, \infty) \times \mathbb{R}^d.$$

We define  $F_\sigma$  for every  $\sigma \in (0, d/2)$  as

$$F_\sigma(\psi) := \frac{1}{\Gamma(d/4 - \sigma/2)} \int_0^\infty t^{-1+\frac{1}{2}(\frac{d}{2}-\sigma)} \int_{\mathbb{R}^d} (\Phi(t, \cdot) * \psi)(x) W(x) dx dt, \quad \psi \in \mathcal{S}(\mathbb{R}^d). \quad (\text{D.15})$$

It is clear that (D.15) defines  $F_\sigma$  as a Gaussian random distribution. To check that this definition yields a fractional Gaussian field with Hurst parameter  $\sigma$ , it therefore suffices to check the covariance formula. By polarization, we just need to check the variance formula

$$\text{var}[F_\sigma(\psi)] = C(\sigma, d) \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |x - y|^{-2\sigma} \psi(x) \psi(y) dx dy, \quad \forall \psi \in \mathcal{S}(\mathbb{R}^d), \quad (\text{D.16})$$

where  $C(\sigma, d)$  is as defined in (D.13).

To check (D.16) we straightforwardly compute

$$\begin{aligned} & \Gamma(d/4 - \sigma/2)^2 \text{var}[F_\sigma(\psi)] \\ &= \mathbb{E} \left[ \int_0^\infty \int_0^\infty \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (st)^{-1+\frac{1}{2}(\frac{d}{2}-\sigma)} (\Phi(t, \cdot) * \psi)(x) (\Phi(s, \cdot) * \psi)(y) W(x) W(y) dx dy dt ds \right] \\ &= \int_0^\infty \int_0^\infty \int_{\mathbb{R}^d} (st)^{-1+\frac{1}{2}(\frac{d}{2}-\sigma)} (\Phi(t, \cdot) * \psi)(x) (\Phi(s, \cdot) * \psi)(x) dx dt ds \\ &= \int_0^\infty \int_0^\infty \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (st)^{-1+\frac{1}{2}(\frac{d}{2}-\sigma)} \Phi(t+s, x-y) \psi(x) \psi(y) dx dy dt ds. \end{aligned} \quad (\text{D.17})$$

In the above display, we used (D.10) to get the second equality, and the semigroup property of the heat kernel to get the third equality.

To evaluate the expression on the last line of (D.17) side, we change variables by setting  $t = \frac{1}{4T} - s$  for a new variable  $T$ , and then reverse the order of integration between the variables  $s$  and  $T$  then set  $s := S/4T$ . After some computations, we get that the last line of (D.17) is equal to

$$(\text{D.17}) = 4^{s-d} \pi^{-d/2} \int_0^1 (S - S^2)^{-1+\frac{1}{2}(\frac{d}{2}-\sigma)} dS \int_0^\infty T^{\sigma-1} \exp(-T|x-y|^2) dT \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \psi(x) \psi(y) dx dy. \quad (\text{D.18})$$

For the first integral factor, we have that

$$\int_0^1 (S - S^2)^{-1+\frac{1}{2}(\frac{d}{2}-\sigma)} dS = \frac{\Gamma(\frac{d}{4} - \frac{\sigma}{2})^2}{\Gamma(\frac{d}{2} - \sigma)}. \quad (\text{D.19})$$

Indeed, the integral on the right side is equal (by definition) to  $B(\frac{1}{2}(\frac{d}{2} - \sigma), \frac{1}{2}(\frac{d}{2} - \sigma))$ , where  $B(\cdot, \cdot)$  is the *beta function*. The beta function can be written in terms of the gamma function by the formula  $B(s_1, s_2) = \Gamma(s_1)\Gamma(s_2)/\Gamma(s_1 + s_2)$ , a proof of which can be found in [Art64, p. 18-19] or in the wikipedia on the beta function. This yields (D.19). By a simple change of variables, we can relate the second integral factor on the right side of (D.18) to the gamma function: we have

$$\int_0^\infty T^{\sigma-1} \exp(-T|x-y|^2) dT = \Gamma(\sigma) |x-y|^{-2\sigma}.$$

We therefore obtain

$$\Gamma(d/4 - \sigma/2)^2 \text{var}[F_\sigma(\psi)] = 4^{\sigma-d/2} \pi^{-d/2} \frac{\Gamma(\frac{d}{4} - \frac{\sigma}{2})^2}{\Gamma(\frac{d}{2} - \sigma)} \Gamma(\sigma) \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |x-y|^{-2\sigma} \psi(x) \psi(y) dx dy.$$

This completes the proof of (D.16).

*D.2.2. Finite range decomposition of fractional Gaussian fields.* In this subsection we provide an explicit decomposition of the fractional Gaussian field  $F_\sigma$  defined in (D.15) of the form

$$F_\sigma = \sum_{n \in \mathbb{Z}} F_{\sigma,n},$$

where  $\{F_{\sigma,n}\}_{n \in \mathbb{Z}}$  is a sequence of Gaussian random fields such that each  $F_{\sigma,n}$  which is defined pointwise, is locally smooth and has range of dependence proportional to  $3^n$ .

We select a partition of unity  $\{\eta_n\}_{n \in \mathbb{Z}}$  on  $\mathbb{R}^d \setminus \{0\}$  satisfying the following:

- For every  $n \in \mathbb{Z}$ , the function  $\eta_n$  belongs to  $C_c^\infty(\mathbb{R}^d)$ ,

$$\mathbf{1}_{B_{3^n} \setminus B_{3^{n-1}}} \leq \eta_n \leq \mathbf{1}_{B_{\frac{3}{2} \cdot 3^n} \setminus B_{\frac{2}{3} \cdot 3^{n-1}}} \quad (\text{D.20})$$

and

$$3^n \|\nabla \eta_n\|_{L^\infty(\mathbb{R}^d)} + 3^{2n} \|\nabla^2 \eta_n\|_{L^\infty(\mathbb{R}^d)} \leq 100. \quad (\text{D.21})$$

- For every  $n \in \mathbb{Z}$  and  $x \in \mathbb{R}^d$ ,

$$\eta_n(x) = \eta_0(3^{-n}x). \quad (\text{D.22})$$

- For every  $x \in \mathbb{R}^d \setminus \{0\}$ ,

$$\sum_{n \in \mathbb{Z}} \eta_n(x) = 1. \quad (\text{D.23})$$

We can construct  $\{\eta_n\}_{n \in \mathbb{N}}$  by taking the indicator function  $\mathbf{1}_{B_1 \setminus B_{1/3}}$  and mollifying the near the inner boundary  $\partial B_{1/3}$  with a smooth, radial function  $\zeta$  which is supported in  $B_{1/9}$  and has unit mass, and then mollifying near the outer boundary  $\partial B_1$  with  $3^{-d}\zeta(3^{-1}\cdot)$ . This defines  $\eta_0$ , and we can then define  $\eta_n$  for  $n \in \mathbb{Z} \setminus \{0\}$  using the scaling relation (D.22). The other properties of  $\eta_n$  are then immediate from the construction.

With an eye toward (D.15), we define  $F_{\sigma,n}$  as follows:

$$F_{\sigma,n}(x) := \frac{1}{\Gamma(d/4 - \sigma/2)} \int_0^\infty t^{-1 + \frac{1}{2}(\frac{d}{2} - \sigma)} \int_{\mathbb{R}^d} (\Phi(t, \cdot) * \eta_n(\cdot - x))(y) W(y) dy dt, \quad x \in \mathbb{R}^d. \quad (\text{D.24})$$

By (D.14), we have that

$$F_{\sigma,n}(x) = 2^\sigma \pi^{-d/2} \int_{\mathbb{R}^d} \eta_n(y - x) |y - x|^{-(\frac{d}{2} + \sigma)} W(y) dy. \quad (\text{D.25})$$

It follows that  $F_{\sigma,n}$  is an  $\mathbb{R}^d$ -stationary Gaussian field with zero mean, and

$$\begin{aligned} \text{var}[F_{\sigma,n}(0)] &\leq 4^\sigma \pi^{-d} \int_{\mathbb{R}^d} \eta_n^2(x) |x|^{-(d+2\sigma)} dx \\ &\leq 4^\sigma \pi^{-d} \int_{B_{\frac{3}{2} \cdot 3^n} \setminus B_{\frac{2}{3} \cdot 3^{n-1}}} |x|^{-(d+2\sigma)} dx \\ &= 4^\sigma \pi^{-d} |\partial B_1| \int_{\frac{2}{3} \cdot 3^{n-1}}^{\frac{3}{2} \cdot 3^n} r^{-(1+2\sigma)} dr \leq \frac{162^\sigma}{2\sigma} \pi^{-d} |\partial B_1| 3^{-2n\sigma}. \end{aligned}$$

Since  $\sigma < d/2$ , we obtain that, for a constant  $C(d) < \infty$ ,

$$\text{var}[F_{\sigma,n}(0)] \leq C \sigma^{-1} 3^{-2n\sigma}.$$



Since  $F_{\sigma,n}(0)$  is Gaussian, we have that

$$|F_{\sigma,n}(0)| = \mathcal{O}_{\Gamma_2}(C\sigma^{-1/2}3^{-n\sigma}). \quad (\text{D.26})$$

By a similar computation, using (D.21), we also have that

$$|\nabla F_{\sigma,n}(0)| = \mathcal{O}_{\Gamma_2}(C\sigma^{-1/2}3^{-n(1+\sigma)}). \quad (\text{D.27})$$

We observe next from (D.20), (D.25) and the independence properties of white noise that

$$F_{\sigma,n} \text{ has range of dependence at most } \frac{3}{2} \cdot 3^n \quad (\text{D.28})$$

and

for every  $m, n \in \mathbb{N}$  with  $|m - n| \geq 2$ , the fields  $F_{\sigma,n}$  and  $F_{\sigma,m}$  are independent.

By (D.22) and the scaling invariance of white noise in (D.11), it is immediate that, for every  $n \in \mathbb{Z}$ , the field  $F_{\sigma,n}$  has the same law as  $3^{-\sigma} F_{\sigma,0}(3^{-n}\cdot)$ .

Finally, by (D.23), we obtain that  $F_\sigma$  defined in (D.15) satisfies

$$F_\sigma(\psi) = \sum_{n \in \mathbb{Z}} \int_{\mathbb{R}^d} F_{\sigma,n}(x) \psi(x) dx, \quad \psi \in \mathcal{S}(\mathbb{R}^d). \quad (\text{D.29})$$

What needs to be justified is that the sum on the right side is convergent, but this is straightforward to obtain from the bounds (D.26) and (D.27), above.

*D.2.3. Proof of Proposition 1.2.* We next present the proof of Proposition 1.2. In fact, we will obtain the following more general statement.<sup>16</sup>

**Proposition D.2.** *Consider the case in which*

$$\mathbf{a}(x) = \lambda \mathbf{I}_d + \mathbf{k}(x),$$

where the  $\mathbf{k}(\cdot)$  is a  $\mathbb{Z}^d$ -stationary random field valued in the  $d$ -by- $d$  anti-symmetric matrices with real entries, and which admits the following decomposition:

$$\mathbf{k}(x) = \sum_{j=0}^{\infty} \mathbf{k}_j(x),$$

where the sequence  $\{\mathbf{k}_j\}_{j \in \mathbb{N}}$  satisfies the following:

- For each  $j \in \mathbb{N}$ , the field  $\mathbf{k}_j$  is a  $\mathbb{Z}^d$ -stationary random field valued in the  $d$ -by- $d$  anti-symmetric matrices;
- For each  $j \in \mathbb{N}$ , the range of dependence of  $\mathbf{k}_j$  is at most  $3^j$ ;
- There exists  $K_0 \in (0, \infty)$  and  $\sigma \in (0, d/2)$  such that, for each  $j \in \mathbb{N}$ ,

$$3^j \|\nabla \mathbf{k}_j\|_{L^\infty(\square_j)} + \|\mathbf{k}_j\|_{L^\infty(\square_j)} \leq \mathcal{O}_{\Gamma_2}(K_0 3^{-\sigma j}). \quad (\text{D.30})$$

---

<sup>16</sup>While Proposition D.2 is mostly about checking the ellipticity assumption (P2), it also implies that (P3) is satisfied with  $\beta = 1 - \frac{2\sigma}{d}$  and  $\Psi(t) = \Gamma_2(c(\frac{d}{2} - \sigma)t)$ , see [AK24, Chapter 3].

Then there exists  $C(d) < \infty$  such that, for every  $\gamma \in (0, \sigma \wedge 1)$ , the ellipticity condition (P2) is satisfied with the parameters

$$\mathbf{E}_0 = \begin{pmatrix} 2(\lambda + C\lambda^{-1}K_0^2\sigma^{-2})\mathbf{I}_d & 0 \\ 0 & 2\lambda^{-1}\mathbf{I}_d \end{pmatrix} \quad \text{and} \quad \Psi_S(t) = (\sigma - \gamma) \exp(C^{-1}t^\gamma - C\gamma^{-1}|\log \gamma|).$$

In view of the discussion in Section D.2.2 above, Proposition D.2 indeed implies Proposition 1.2, since the assumptions of the latter imply those of the former with  $K_0 = C(d)\sigma^{-1/2}$  in (D.30).

*Proof of Proposition D.2.* Fix  $\gamma \in (0, 1)$  and  $\rho \in (0, \alpha)$  with  $\gamma < 2\rho$ . Let  $j, n, m \in \mathbb{N}$  with  $j \leq n \leq m$ . By (D.30) and Lemma C.2, there exists a constant  $C(d) < \infty$  such that, for every  $z \in \mathbb{R}^d$ ,

$$\begin{aligned} \|\mathbf{k}_j\|_{\underline{L}^2(z+\square_n)}^2 &= \sum_{z' \in z+3^j\mathbb{Z}^d \cap \square_n} \|\mathbf{k}_j\|_{\underline{L}^2(z'+\square_n)}^2 \\ &= \mathbb{E}[\|\mathbf{k}_j\|_{\underline{L}^2(\square_n)}^2] + \sum_{z' \in z+3^j\mathbb{Z}^d \cap \square_n} (\|\mathbf{k}_j\|_{\underline{L}^2(z'+\square_n)}^2 - \mathbb{E}[\|\mathbf{k}_j\|_{\underline{L}^2(\square_n)}^2]) \\ &\leq CK_0^2 3^{-2\sigma j} + \mathcal{O}_{\Gamma_1}(CK_0^2 3^{-2\sigma j} 3^{-\frac{d}{2}(n-j)}). \end{aligned}$$

By the Cauchy-Schwarz inequality, we have that

$$\|\mathbf{k}_0 + \dots + \mathbf{k}_n\|_{\underline{L}^2(\square_n)}^2 \leq \frac{C}{\sigma} \sum_{j=0}^n 3^{\sigma j} \|\mathbf{k}_j\|_{\underline{L}^2(\square_n)}^2.$$

Combining these, we obtain

$$\begin{aligned} \|\mathbf{k}_0 + \dots + \mathbf{k}_n\|_{\underline{L}^2(\square_n)}^2 &\leq \frac{CK_0^2}{\sigma} \sum_{j=0}^n C 3^{-\sigma j} + \mathcal{O}_{\Gamma_1}\left(\frac{CK_0^2}{\sigma} \sum_{j=0}^n 3^{-\sigma j} 3^{-\frac{d}{2}(n-j)}\right) \\ &\leq \frac{CK_0^2}{\sigma^2} + \mathcal{O}_{\Gamma_1}\left(\frac{CK_0^2 3^{-\sigma n}}{\sigma}\right). \end{aligned}$$

On the other hand, by (D.30) and the generalized triangle inequality,

$$\left\| \sum_{j=n+1}^{\infty} \mathbf{k}_j \right\|_{\underline{L}^2(\square_n)} \leq \sum_{j=n+1}^{\infty} \|\mathbf{k}_j\|_{L^\infty(\square_j)} \leq \mathcal{O}_{\Gamma_2}\left(\frac{CK_0}{\sigma} 3^{-n\sigma}\right).$$

Combining the previous two displays yields, for every  $n \in \mathbb{N}$ ,

$$\|\mathbf{k}\|_{\underline{L}^2(\square_n)}^2 \leq \frac{CK_0^2}{\sigma^2} + \mathcal{O}_{\Gamma_1}\left(\frac{CK_0^2 3^{-\sigma n}}{\sigma}\right).$$

For  $n \in \mathbb{Z}$  with  $n < 0$ , the assumption (D.30) already gives a better bound; combining this with the above estimate yields, for every  $n \in \mathbb{Z}$ ,

$$\|\mathbf{k}\|_{\underline{L}^2(\square_n)}^2 \leq \frac{CK_0^2}{\sigma^2} + \mathcal{O}_{\Gamma_1}\left(\frac{CK_0^2 3^{-\sigma(n \vee 0)}}{\sigma}\right).$$

By the Markov inequality and a union bound, we deduce, for every  $m, n \in \mathbb{N}$  with  $n \leq m$ ,

$$\begin{aligned} \mathbb{P}\left[\max_{z \in 3^n\mathbb{Z}^d \cap \square_m} \|\mathbf{k}\|_{\underline{L}^2(z+\square_n)}^2 > \frac{CK_0^2}{\sigma^2} 3^{\gamma(m-n)}\right] &\leq \sum_{z \in 3^n\mathbb{Z}^d \cap \square_m} \mathbb{P}\left[\|\mathbf{k}\|_{\underline{L}^2(z+\square_n)}^2 > \frac{CK_0^2}{\sigma^2} 3^{\gamma(m-n)}\right] \\ &\leq 3^{d(m-n)} \mathbb{P}\left[\|\mathbf{k}\|_{\underline{L}^2(\square_n)}^2 > \frac{CK_0^2}{\sigma^2} 3^{\gamma(m-n)}\right] \\ &\leq 3^{d(m-n)} \exp\left(-c\sigma^{-1} 3^{\gamma(m-n)+\sigma(n \vee 0)}\right). \end{aligned}$$

By another union bound, we obtain, for every  $m \in \mathbb{N}$ ,

$$\begin{aligned} & \mathbb{P} \left[ \exists n \in \mathbb{Z} \cap (-\infty, m], \max_{z \in 3^n \mathbb{Z}^d \cap \square_m} \|\mathbf{k}\|_{\underline{L}^2(z+\square_n)}^2 > \frac{CK_0^2}{\sigma^2} 3^{\gamma(m-n)} \right] \\ & \leq \sum_{n=-\infty}^m \mathbb{P} \left[ \max_{z \in 3^n \mathbb{Z}^d \cap \square_m} \|\mathbf{k}\|_{\underline{L}^2(z+\square_n)}^2 > \frac{CK_0^2}{\sigma^2} 3^{\gamma(m-n)} \right] \\ & \leq \sum_{n=-\infty}^m 3^{d(m-n)} \exp \left( -c\sigma^{-1} 3^{\gamma(m-n)+\sigma(n \vee 0)} \right) \leq \frac{1}{\sigma - \gamma} \exp \left( \frac{C|\log \gamma|}{\gamma} \right) \exp(-c3^{\gamma m}). \end{aligned}$$

We next define

$$\mathcal{S} := \sup \left\{ 3^{m+1} : m \in \mathbb{N}, \exists n \in \mathbb{Z} \cap (-\infty, m], \max_{z \in 3^n \mathbb{Z}^d \cap \square_m} \|\mathbf{k}\|_{\underline{L}^2(z+\square_n)}^2 > \frac{CK_0^2}{\sigma^2} 3^{\gamma(m-n)} \right\}.$$

The previous display says that  $\mathcal{S}$  has the integrability claimed in the statement. Its definition implies that (1.17) holds for  $\mathbf{E}_0$  given in the statement of the proposition, since by the definition of  $\mathcal{S}$  we have

$$\begin{aligned} 3^m \geq \mathcal{S} & \implies \max_{n \in \mathbb{Z} \cap (-\infty, m]} \max_{z \in 3^n \mathbb{Z}^d \cap \square_m} \|\mathbf{k}\|_{\underline{L}^2(z+\square_n)}^2 \leq \frac{CK_0^2}{\sigma^2} 3^{\gamma(m-n)} \\ & \implies \begin{cases} \mathbf{A}(z + \square_n) \leq \begin{pmatrix} (2\lambda + C\lambda^{-1}K_0^2\sigma^{-2}3^{\gamma(m-n)})\mathbf{I}_d & 0 \\ 0 & 2\lambda^{-1}\mathbf{I}_d \end{pmatrix}, \\ \forall n \in \mathbb{Z} \cap (-\infty, m], z \in 3^n \mathbb{Z}^d \cap \square_m. \end{cases} \end{aligned}$$

This completes the proof.  $\square$

**D.3. Log-normal fields.** In this section we prove Proposition 1.3. We first consider the case of a log-normal random field with a finite range of dependence. Notice that we make no symmetry assumption on  $\mathbf{g}$ , nor do we assume that the symmetric part of  $\mathbf{g}$  is nonnegative. (The symmetric part of  $\mathbf{a}(\cdot)$  will however be positive.)

**Proposition D.3** (Log-normal field with finite range of dependence). *Suppose that  $\mathbf{a}(\cdot)$  is given by*

$$\mathbf{a}(x) = \exp(\mathbf{g}(x)), \quad (\text{D.31})$$

where  $L \geq 1$ ,  $h > 0$  and  $\mathbf{g}(\cdot)$  is a random field valued in the  $\mathbb{R}^{d \times d}$  matrices satisfying

$$\begin{cases} \mathbf{g} \text{ is } \mathbb{Z}^d\text{-stationary,} \\ \mathbf{g} \text{ has range of dependence at most } L, \\ \|\mathbf{g}\|_{L^\infty(\square_0)} \leq \mathcal{O}_{\Gamma_2}(h). \end{cases} \quad (\text{D.32})$$

Then there exists  $C(d) < \infty$  such that, for every  $\gamma \in (0, 1)$ , the field  $\mathbf{a}(\cdot)$  satisfies assumption (P2) with parameters  $\gamma$ ,

$$\mathbf{E}_0 = C \exp(18h^2) \mathbf{I}_{2d},$$

and minimal scale  $\mathcal{S}$  satisfying

$$\mathbb{P}[\mathcal{S} > Lt] \leq \exp(C h^2 \gamma^{-2} - c h^{-2} \log^2 t).$$

The assumption (D.32) is satisfied for example, if  $\mathbf{g}(\cdot)$  is a  $\mathbb{R}^{d \times d}$ -valued stationary Gaussian field with a compactly supported covariance function. Another example is if  $\mathbf{g}(\cdot)$  is given by the convolution of a bounded, deterministic and compactly supported  $\mathbb{R}^{d \times d}$ -valued function on  $\mathbb{R}^d$  and a Poisson point process on  $\mathbb{R}^d$ . Note that, in both of these cases, the distributions of  $\|\mathbf{a}\|_{L^\infty(\square_0)}$  and  $\|\mathbf{a}^{-1}\|_{L^\infty(\square_0)}$  have tails which are as fat as those of a log-normal random variable. More generally, under the assumption (D.32), we have

$$\mathbb{P}[\|\mathbf{a}\|_{L^\infty(\square_0)} > t] \leq \exp\left(-\frac{1}{2}\left(\frac{\log t}{h}\right)^2\right), \quad \forall t > 0,$$

with the same bound holding also for  $\mathbf{a}^{-1}$  in place of  $\mathbf{a}$ .

It is clear that the field  $\mathbf{a}(\cdot)$  satisfies the stationarity assumption (P1). The validity of (P3) with  $\beta = 0$  and  $\Psi = \Gamma_2(\cdot/CL)$  for some  $C(d) < \infty$  follows from the finite range of dependence assumption; see [AK24, Section 3.2.1].

*Proof of Proposition D.3.* We will assume without loss of generality that  $L = 1$ . Let us decompose  $\mathbf{g}$  into pieces by writing

$$\mathbf{g} = \sum_{j=0}^{\infty} \mathbf{g}_j, \quad \text{where} \quad \mathbf{g}_0(x) := \mathbf{g}(x)\mathbf{1}_{|\mathbf{g}(x)| < 1} \quad \text{and} \quad \mathbf{g}_j(x) := \mathbf{g}(x)\mathbf{1}_{\{2^{j-1} \leq |\mathbf{g}(x)| < 2^j\}}, \quad \forall j \geq 1.$$

Note that for each  $x$ , exactly one of the  $\mathbf{g}_j(x) = \mathbf{g}(x)$  and the rest are zero. For every  $m \in \mathbb{N}$  and  $\lambda \geq 1$ ,

$$\int_{\square_m} |\mathbf{a}(x)|^{\lambda/h} dx = \int_{\square_m} |\exp(\lambda h^{-1} \mathbf{g}(x))| dx = 1 + \sum_{j=0}^{\infty} \int_{\square_m} (|\exp(\lambda h^{-1} \mathbf{g}_j(x))| - 1) dx.$$

Let  $N_{m,j}$  denote the number of distinct  $z \in \mathbb{Z}^d \cap \square_m$  such that  $\mathbf{g}_j$  does not vanish in  $z + \square_0$ . The distribution of  $N_{m,j}$  is essentially a binomial with parameters  $3^{dm}$  (the number of unit cubes) and  $p_j$ , the probability of  $\mathbf{g}_j \neq 0$  in  $\square_0$ , which satisfies the upper bound

$$p_j = \mathbb{P}[\mathbf{g}_j \neq 0 \text{ in } \square_0] \leq \mathbb{P}[h^{-1}\|\mathbf{g}\|_{L^\infty(\square_0)} \geq 2^{j-1}] \leq \exp(-2^{2(j-1)}).$$

This is not quite right, since neighboring unit cubes are not independent; it would be more accurate to say that  $N_{m,j}$  is bounded by  $3^d$  many identically distributed copies of a binomial distribution with parameters  $3^{(d-1)m}$  and  $p_j$ . To see this, we partition the collection of unit cubes in  $\square_m$  into  $3^d$  different subcollections, each of which contains cubes that are separated by a distance of at least one. From standard tail estimates on the binomial distribution (or just apply Hoeffding's inequality), we have the bound

$$\mathbb{P}[N_{m,j} \geq 3^{md} \exp(-2^{2(j-1)}) + 3^d t] \leq \exp(-2 \cdot 3^{-(d-1)m} t^2).$$

That is,

$$N_{m,j} \leq 3^{md} \exp(-2^{2(j-1)}) + \mathcal{O}_{\Gamma_2}(C 3^{\frac{d}{2}m}). \quad (\text{D.33})$$

Alternatively, when  $j$  is relatively large, it is better to use a the crude estimate, obtained from a simple union bound,

$$\mathbb{P}[N_{m,j} \neq 0] \leq 3^{md} \mathbb{P}[h^{-1}\|\mathbf{g}\|_{L^\infty(\square_0)} > 2^{j-1}] \leq 3^{md} \exp(-2^{2(j-1)}). \quad (\text{D.34})$$

Applying (D.33) yields the estimate

$$\begin{aligned} \int_{\square_m} \exp(\lambda h^{-1} \mathbf{g}_j(x)) dx &\leq 1 + \frac{N_{m,j}}{|\square_m|} \exp(\lambda 2^j) \\ &\leq 1 + \exp(\lambda 2^j - 2^{2(j-1)}) + \mathcal{O}_{\Gamma_2}(C 3^{-\frac{d}{2}m} \exp(\lambda 2^j)). \end{aligned} \quad (\text{D.35})$$

This inequality will be used for small values of  $j$ , namely those satisfying  $\exp(\lambda 2^j) \leq t 3^{\delta m}$  for parameters  $\delta \in (0, \frac{1}{2}]$  and  $t \geq 1$ , to be selected. Summing over this range of  $j$ 's, we get

$$\begin{aligned} &\sum_{j \in \mathbb{N} : \exp(\lambda 2^j) \leq t 3^{\delta m}} \int_{\square_m} (|\exp(\lambda h^{-1} \mathbf{g}_j(x))| - 1) dx \\ &\leq \sum_{j=0}^{\infty} \exp(\lambda 2^j - 2^{2(j-1)}) + \sum_{j \in \mathbb{N} : \exp(\lambda 2^j) \leq t 3^{\delta m}} \mathcal{O}_{\Gamma_2}(C 3^{-\frac{d}{2}m} \exp(\lambda 2^j)) \\ &\leq \sum_{j=0}^{\infty} \exp(2\lambda^2 + 2^{2j-3} - 2^{2j-2}) + \mathcal{O}_{\Gamma_2}(C t 3^{-(\frac{d}{2}-\delta)m}) \\ &\leq C \exp(2\lambda^2) + \mathcal{O}_{\Gamma_2}(C t 3^{-(\frac{d}{2}-\delta)m}). \end{aligned}$$

For  $j$ 's larger than this, we simply hope that  $N_{m,j} = 0$ , or else we give up. Using (D.34), we have

$$\begin{aligned} \mathbb{P}\left[\exists j \in \mathbb{N}, \exp(\lambda 2^j) > t 3^{\delta m}, N_{m,j} \neq 0\right] &\leq \sum_{j \in \mathbb{N} : \exp(\lambda 2^j) > t 3^{\delta m}} 3^{md} \exp(-2^{2j}) \\ &\leq C 3^{md} \exp(-(\log 3^{\frac{\delta m}{\lambda}})^2) \\ &\leq \exp(-(\lambda^{-1} \log c t 3^{\delta m})^2). \end{aligned}$$

We deduce that

$$\mathbb{P}\left[\int_{\square_m} |\exp(\lambda h^{-1} \mathbf{g}(x))| dx \geq C \exp(2\lambda^2) + t\right] \leq \exp(-c 3^{(d-2\delta)m} t) + \exp(-(\lambda^{-1} \log c t 3^{\delta m})^2).$$

Since  $t \geq 1$ , the second term on the right is larger than the first. Taking  $\delta = 1/2$ , we obtain, for constants  $C(d) < \infty$  and  $c(d) > 0$ ,

$$\mathbb{P}\left[\int_{\square_m} |\exp(\lambda h^{-1} \mathbf{g}(x))| dx \geq C \exp(2\lambda^2) + t\right] \leq C \exp(-c \lambda^{-2} (\log t 3^m)^2).$$

Fix now a parameter  $\gamma \in (0, 1]$ . By performing a union bound over a mesoscale represented by  $n \in \mathbb{N}$ ,  $n \leq m$ , we obtain

$$\begin{aligned} &\mathbb{P}\left[\sup_{z \in 3^n \mathbb{Z}^d \cap \square_m} \int_{z + \square_n} |\exp(\lambda h^{-1} \mathbf{g}(x))| dx \geq C 3^{\gamma(m-n)} \exp(2\lambda^2)\right] \\ &\leq 3^{d(m-n)} \mathbb{P}\left[\int_{\square_n} |\exp(\lambda h^{-1} \mathbf{g}(x))| dx \geq C 3^{\gamma(m-n)} \exp(4\lambda^2)\right] \\ &\leq C 3^{d(m-n)} \exp(-c \lambda^{-2} (\log 3^{\gamma(m-n)} + \log 3^m)^2) \\ &\leq C 3^{d(m-n)} \exp(-c \lambda^{-2} (\log 3^{\gamma(m-n)})^2) \exp(-c \lambda^{-2} (\log 3^m)^2). \end{aligned}$$

Taking another union bound, we obtain

$$\begin{aligned}
\mathbb{P} \left[ \sup_{n \in \{0, \dots, m\}} \sup_{z \in 3^n \mathbb{Z}^d \cap \square_m} \int_{z + \square_n} |\exp(\lambda h^{-1} \mathbf{g}(x))| dx \geq C 3^{\gamma(m-n)} \exp(2\lambda^2) \right] \\
\leq C \sum_{n=0}^m \left( 3^{d(m-n)} \exp(-c\lambda^{-2}(\log 3^{\gamma(m-n)})^2) \right) \exp(-c\lambda^{-2}(\log 3^m)^2) \\
\leq C \exp(C(\lambda\gamma^{-1})^2) \exp(-c\lambda^{-2}(\log 3^m)^2).
\end{aligned}$$

Taking yet another union bound, we get, for every  $k \in \mathbb{N}$ ,

$$\begin{aligned}
\mathbb{P} \left[ \sup_{m \geq k} \sup_{n \in \{0, \dots, m\}} 3^{-\gamma(m-n)} \sup_{z \in 3^n \mathbb{Z}^d \cap \square_m} \int_{z + \square_n} |\exp(\lambda h^{-1} \mathbf{g}(x))| dx \geq C \exp(2\lambda^2) \right] \\
\leq \exp(C(\lambda\gamma^{-1})^2) \sum_{m=k}^{\infty} \exp(-c\lambda^{-2}(\log 3^m)^2) \\
\leq \exp(C(\lambda\gamma^{-1})^2) \exp(-c\lambda^{-2}(\log 3^k)^2). \tag{D.36}
\end{aligned}$$

For the cubes smaller than the unit cubes, we use the bound

$$\begin{aligned}
\mathbb{P}[\|\exp(\lambda h^{-1} \mathbf{g})\|_{L^\infty(\square_m)} > 1 + t] &\leq 3^{dm} \mathbb{P}[\|\exp(\lambda h^{-1} \mathbf{g})\|_{L^\infty(\square_0)} > 1 + t] \\
&\leq 3^{dm} \exp(-\lambda^{-2} \log^2(1 + t)).
\end{aligned}$$

From this we obtain

$$\begin{aligned}
\mathbb{P} \left[ \sup_{m \geq k} \sup_{n \in \mathbb{N}} 3^{-\gamma(m-n)} \sup_{z \in 3^n \mathbb{Z}^d \cap \square_m} \int_{z + \square_n} |\exp(\lambda h^{-1} \mathbf{g}(x))| dx \geq C \right] \\
\leq \mathbb{P} \left[ \sup_{m \geq k} 3^{-\gamma m} \|\exp(\lambda h^{-1} \mathbf{g})\|_{L^\infty(\square_m)} \geq C \right] \\
\leq \exp(C\lambda^2) \sum_{m=k}^{\infty} \exp(-c\lambda^{-2}(\log 3^{\gamma m})^2) \\
\leq \exp(C(\lambda\gamma^{-1})^2) \exp(-c\lambda^{-2}(\log 3^k)^2), \tag{D.37}
\end{aligned}$$

as above. If we define the minimal scale  $\mathcal{S}$  by

$$\mathcal{S}_\lambda := \sup \left\{ 3^m : \sup_{n \in \mathbb{Z} \cap (-\infty, m]} \sup_{z \in 3^n \mathbb{Z}^d \cap \square_m} \int_{z + \square_n} |\exp(\lambda h^{-1} \mathbf{g}(x))| dx \geq C \exp(2\lambda^2) 3^{\gamma(m-n)} \right\}, \tag{D.38}$$

then (D.36) and (D.37) imply that

$$\mathbb{P}[\mathcal{S}_\lambda > 3^k] \leq \exp(C(\lambda\gamma^{-1})^2) \exp(-c\lambda^{-2}(\log 3^k)^2), \tag{D.39}$$

and hence

$$k \geq C\lambda^2\gamma^{-2} \implies \mathbb{P}[\mathcal{S}_\lambda > 3^k] \leq \exp(-c\lambda^{-2}(\log 3^k)^2).$$

In other words,

$$\mathcal{S}_\lambda = \exp(C\lambda^2\gamma^{-2}) + \mathcal{O}_{\Psi_{C\lambda}}(C\lambda).$$

It is clear from its definition in (D.38) that

$$3^m \geq \mathcal{S}_\lambda \implies \int_{z + \square_n} |\mathbf{a}(x)|^{\lambda/h} dx \leq C 3^{\gamma(m-n)} \exp(2\lambda^2), \quad \forall n \in \mathbb{Z} \cap (-\infty, m], \quad z \in 3^n \mathbb{Z}^d \cap \square_m.$$

The same argument gives a similar minimal scale for  $\mathbf{a}^{-1}$  in place of  $\mathbf{a}$ , so by taking the maximum of these we may suppose that  $\mathcal{S}$  satisfies (D.39) and

$$3^m \geq \mathcal{S}_\lambda \implies \begin{cases} \int_{z+\square_n} (|\mathbf{a}(x)|^{\lambda/h} + |\mathbf{a}^{-1}(x)|^{\lambda/h}) dx \leq C3^{\gamma(m-n)} \exp(2\lambda^2), \\ \forall n \in \mathbb{Z} \cap (-\infty, m], z \in 3^n \mathbb{Z}^d \cap \square_m. \end{cases} \quad (\text{D.40})$$

Since  $|\mathbf{A}(x)| \leq (1 + |\mathbf{a}(x)|^2)|\mathbf{a}^{-1}(x)|$ , we deduce from (D.40) and the Hölder inequality that

$$\begin{aligned} 3^m \geq \mathcal{S}_{3h} &\implies \int_{z+\square_n} |\mathbf{A}(x)| dx \leq C3^{\gamma(m-n)} \exp(18h^2), \quad \forall n \in \mathbb{Z} \cap (-\infty, m], z \in 3^n \mathbb{Z}^d \cap \square_m \\ &\implies \mathbf{A}(z + \square_n) \leq C3^{\gamma(m-n)} \exp(18h^2) \mathbf{I}_{2d}, \quad \forall n \in \mathbb{Z} \cap (-\infty, m], z \in 3^n \mathbb{Z}^d \cap \square_m. \end{aligned}$$

Note that scales below the unit scale can be taken care of immediately from the third line of (D.32) and a union bound. This completes the proof.  $\square$

We turn to the proof of Proposition 1.3. We will prove the following more general statement.<sup>17</sup>

**Proposition D.4.** *Suppose that  $\mathbf{a}(\cdot)$  is given by*

$$\mathbf{a}(x) = \exp(\mathbf{g}(x)), \quad (\text{D.41})$$

where  $\mathbf{g}(\cdot)$  is an  $\mathbb{R}^{d \times d}$ -valued random field which admits the decomposition

$$\mathbf{g}(x) = \sum_{j=0}^{\infty} \mathbf{g}_j(x),$$

where the sequence  $\{\mathbf{g}_j\}_{j \in \mathbb{N}}$  satisfies the following:

- For each  $j \in \mathbb{N}$ , the field  $\mathbf{g}_j$  is a  $\mathbb{Z}^d$ -stationary random field valued in the  $d$ -by- $d$  matrices;
- For each  $j \in \mathbb{N}$ , the range of dependence of  $\mathbf{g}_j$  is at most  $3^j$ ;
- There exists  $h \in (0, \infty)$  and  $\sigma \in (0, d/2)$  such that, for each  $j \in \mathbb{N}$ ,

$$3^j \|\nabla \mathbf{g}_j\|_{L^\infty(\square_j)} + \|\mathbf{g}_j\|_{L^\infty(\square_j)} \leq \mathcal{O}_{\Gamma_2}(h3^{-\sigma j}). \quad (\text{D.42})$$

Then there exists  $C(d) < \infty$  such that, for every  $\gamma \in (0, 1)$ , the field  $\mathbf{a}(\cdot)$  satisfies assumption (P2) with parameters  $\gamma$ ,

$$\mathbf{E}_0 = \exp(Ch^2\sigma^{-2})\mathbf{I}_{2d},$$

and minimal scale  $\mathcal{S}$  satisfying

$$\mathbb{P}[\mathcal{S} > t] \leq \exp(Ch^2\sigma^{-2}\gamma^{-2} - ch^{-2}\sigma^2 \log^2 t).$$

*Proof.* Here we approximate by finite range fields and apply the previous result. Denote, for each  $k \in \mathbb{N}$ ,

$$\hat{\mathbf{g}}_k := \sum_{j=0}^k \mathbf{g}_j$$

---

<sup>17</sup>While Proposition D.4 is mostly about checking the ellipticity assumption (P2), it also implies that (P3) is satisfied with  $\beta = 1 - \frac{2\sigma}{d}$  and  $\Psi(t) = \Gamma_2(c(\frac{d}{2} - \sigma)t)$ , see [AK24, Chapter 3].)

and observe that, for each  $k, m \in \mathbb{N}$  with  $k \leq m$ ,

$$\|\mathbf{g} - \hat{\mathbf{g}}_k\|_{L^\infty(\square_k)} \leq \mathcal{O}_{\Gamma_2}(Ch\sigma^{-1}3^{-\sigma k})$$

and hence

$$\mathbb{P}[\|\mathbf{g} - \hat{\mathbf{g}}_k\|_{L^\infty(\square_m)} > t] \leq 3^{d(m-k)} \exp(-ch^{-2}\sigma^2 3^{2\sigma k} t^2).$$

Taking  $k = \lceil m/2 \rceil$  yields, for every  $t \geq Ch\sigma^{-2}$ ,

$$\mathbb{P}[\|\mathbf{g} - \hat{\mathbf{g}}_{\lceil m/2 \rceil}\|_{L^\infty(\square_m)} > t] \leq \exp(-ch^{-2}\sigma^2 3^{\sigma m} t^2). \quad (\text{D.43})$$

Applying the above result for finite range fields to  $\hat{\mathbf{a}}_{\lceil m/2 \rceil} := \exp(h\hat{\mathbf{g}}_{\lceil m/2 \rceil})$ , using that

$$\hat{\mathbf{g}}_{\lceil m/2 \rceil} = \mathcal{O}_{\Gamma_2}(Ch\sigma^{-1})$$

gives, for every  $m \geq Ch^2\sigma^{-2}\gamma^{-2}$ ,

$$\begin{aligned} \mathbb{P}\left[\exists n \in \mathbb{Z} \cap (-\infty, m], z \in 3^n \mathbb{Z}^d \cap \square_m, \int_{z+\square_n} |\hat{\mathbf{A}}_{\lceil m/2 \rceil}(x)| dx \geq 3^{\gamma(m-n)} \exp(Ch^2\sigma^{-2}h^2)\right] \\ \leq \exp(-ch^{-2}\sigma^2(\log 3^m)^2). \end{aligned} \quad (\text{D.44})$$

Combining (D.43) and (D.44) yields, for every  $m \in \mathbb{N}$  with  $m \geq Ch^2\sigma^{-2}\gamma^{-2}$ ,

$$\begin{aligned} \mathbb{P}\left[\exists n \in \mathbb{Z} \cap (-\infty, m], z \in 3^n \mathbb{Z}^d \cap \square_m, \int_{z+\square_n} |\mathbf{A}(x)| dx \geq 3^{\gamma(m-n)} \exp(Ch^2\sigma^{-2})\right] \\ \leq \exp(-ch^{-2}\sigma^2(\log 3^m)^2). \end{aligned}$$

This completes the proof.  $\square$

**Acknowledgments.** S.A. was supported by NSF grants DMS-1954357 and DMS-2000200 and by the Simons Programme at IHES. T.K. was supported by the Academy of Finland and by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 818437).

## References

- [ABRK23] S. Armstrong, A. Bou-Rabee, and T. Kuusi. Superdiffusive central limit theorem for a Brownian particle in a critically-correlated incompressible random drift, 2023. arXiv:2404.01115.
- [AD18] S. Armstrong and P. Dario. Elliptic regularity and quantitative homogenization on percolation clusters. *Comm. Pure Appl. Math.*, 71(9):1717–1849, 2018.
- [AK24] S. Armstrong and T. Kuusi. Elliptic homogenization from qualitative to quantitative, 2024. preprint.
- [AKM16] S. Armstrong, T. Kuusi, and J.-C. Mourrat. Mesoscopic higher regularity and subadditivity in elliptic homogenization. *Comm. Math. Phys.*, 347(2):315–361, 2016.
- [AKM17] S. Armstrong, T. Kuusi, and J.-C. Mourrat. The additive structure of elliptic homogenization. *Invent. Math.*, 208(3):999–1154, 2017.
- [AKM19] S. Armstrong, T. Kuusi, and J.-C. Mourrat. *Quantitative stochastic homogenization and large-scale regularity*, volume 352 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Cham, 2019.



- [AM16] S. N. Armstrong and J.-C. Mourrat. Lipschitz regularity for elliptic equations with random coefficients. *Arch. Ration. Mech. Anal.*, 219(1):255–348, 2016.
- [And78] T. Ando. *Topics on operator inequalities*. Hokkaido University, Research Institute of Applied Electricity, Division of Applied Mathematics, Sapporo, 1978.
- [Art64] E. Artin. *The gamma function*. Holt, Rinehart and Winston, New York-Toronto-London, 1964.
- [AS16] S. N. Armstrong and C. K. Smart. Quantitative stochastic homogenization of convex integral functionals. *Ann. Sci. Éc. Norm. Supér. (4)*, 49(2):423–481, 2016.
- [BMdIP23] M. Bakhshizadeh, A. Maleki, and V. H. de la Pena. Sharp concentration results for heavy-tailed distributions. *Inf. Inference*, 12(3):Paper No. iaad011, 31, 2023.
- [CGQ24] N. Clozeau, A. Gloria, and S. Qi. Quantitative homogenization for log-normal coefficients, 2024. arXiv:2403.00168.
- [Dar21] P. Dario. Optimal corrector estimates on percolation cluster. *Ann. Appl. Probab.*, 31(1):377–431, 2021.
- [DCKT21] H. Duminil-Copin, G. Kozma, and V. Tassion. Upper bounds on the percolation correlation length. In *In and out of equilibrium 3. Celebrating Vladas Sidoravicius*, volume 77 of *Progr. Probab.*, pages 347–369. Birkhäuser/Springer, Cham, [2021] ©2021.
- [DG21] P. Dario and C. Gu. Quantitative homogenization of the parabolic and elliptic Green’s functions on percolation clusters. *Ann. Probab.*, 49(2):556–636, 2021.
- [DG22] M. Duerinckx and A. Gloria. Quantitative homogenization theory for random suspensions in steady Stokes flow. *J. Éc. polytech. Math.*, 9:1183–1244, 2022.
- [FMM98] E. Fabes, O. Mendez, and M. Mitrea. Boundary layers on Sobolev-Besov spaces and Poisson’s equation for the Laplacian in Lipschitz domains. *J. Funct. Anal.*, 159(2):323–368, 1998.
- [FP97] M. Fiedler and V. Pták. A new positive definite geometric mean of two positive definite matrices. *Linear Algebra Appl.*, 251:1–20, 1997.
- [GO11] A. Gloria and F. Otto. An optimal variance estimate in stochastic homogenization of discrete elliptic equations. *Ann. Probab.*, 39(3):779–856, 2011.
- [GO12] A. Gloria and F. Otto. An optimal error estimate in stochastic homogenization of discrete elliptic equations. *Ann. Appl. Probab.*, 22(1):1–28, 2012.
- [Har90] T. Hara. Mean-field critical behaviour for correlation length for percolation in high dimensions. *Probab. Theory Related Fields*, 86(3):337–385, 1990.
- [JK95] D. Jerison and C. E. Kenig. The inhomogeneous Dirichlet problem in Lipschitz domains. *J. Funct. Anal.*, 130(1):161–219, 1995.
- [KO84] A. Kufner and B. Opic. How to define reasonably weighted Sobolev spaces. *Comment. Math. Univ. Carolin.*, 25(3):537–554, 1984.
- [LSSW16] A. Lodhia, S. Sheffield, X. Sun, and S. S. Watson. Fractional Gaussian fields: A survey. *Probab. Surv.*, 13:1 – 56, 2016.
- [Sho97] R. E. Showalter. *Monotone operators in Banach space and nonlinear partial differential equations*, volume 49 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.
- [SW01] S. Smirnov and W. Werner. Critical exponents for two-dimensional percolation. *Math. Res. Lett.*, 8(5-6):729–744, 2001.
- [Wei83] S. Weinberg. Why the renormalization group is a good thing. In *Asymptotic Realms of Physics*, pages 1–19. MIT Press, 1983.